A MODULE-THEORETIC APPROACH TO CLIFFORD THEORY FOR BLOCKS

S. J. WITHERSPOON

(Communicated by Ronald M. Solomon)

ABSTRACT. This work concerns a generalization of Clifford theory to blocks of group-graded algebras. A module-theoretic approach is taken to prove a one-to-one correspondence between the blocks of a fully group-graded algebra covering a given block of its identity component, and conjugacy classes of blocks of a twisted group algebra. In particular, this applies to blocks of a finite group covering blocks of a normal subgroup.

1. INTRODUCTION

In 1973 Dade developed the block theory of group-graded algebras, in particular extending the Clifford correspondence to blocks of group-graded algebras [9]. The Clifford correspondence for representations is a one-to-one correspondence between the irreducible representations of a group G having a given constituent when restricted to a normal subgroup, and the irreducible projective representations (with respect to a particular 2-cocycle) of a certain subgroup of the corresponding quotient group [4]. Dade has recently used his results to make progress towards solving conjectures of Alperin and Dade [12, 13, 15]. Ellers has used Dade's work in making progress on a generalization of Alperin's Conjecture [16]. In view of this recent activity, it seems useful to provide an alternative approach to Dade's results. In this work we treat blocks as bimodules to provide a new proof of the Clifford correspondence for blocks of group-graded algebras. This approach clarifies the connection between the Clifford correspondence for blocks and that for representations.

Specifically, let G be a finite group and A a *fully G -graded algebra* of finite dimension over an algebraically closed field k . That is, $A = \sum_{g \in G} A_g$ is a direct sum of subspaces A_g with $A_g A_h = A_{gh}$ for all $g, h \in G$. Here $A_g A_h$ denotes the set of all finite sums of products xy , where $x \in A_g$ and $y \in A_h$. The first example of a fully G -graded algebra is $A = k\Gamma$ for a group Γ with normal subgroup N and $G = \Gamma/N$. In this case the subspace A_g is spanned by the elements of the coset g . As the identity component is $A_1 = kN$, comparing blocks of A_1 to blocks of A amounts to comparing blocks of N to blocks of Γ .

We consider A to be a right $A^{op} \otimes A$ -module, where A^{op} is the opposite algebra to A , via left and right multiplication by elements of A . A block of A is an indecomposable direct summand of the $A^{op} \otimes A$ -module A . A block $\tilde{B}$ of A *covers* a block

Received by the editors April 20, 1998.

1991 *Mathematics Subject Classification.* Primary 20C20, 20C25.

Research supported in part by NSERC grant # OGP0170281.

B of A_1 if B is a direct summand of the module $\tilde{B}$ restricted to $A_1^{op} \otimes A_1$. That this is equivalent to $\tilde{B}$ lying over B [9, p. 218] follows from the proof of Lemma 2.1 below. The group G acts by conjugation on the blocks of A_1 , an element $g \in G$ sending B to $A_{g^{-1}}BA_g$. In Section 4 we assume B is G -invariant and derive a one-to-one correspondence between the blocks of A covering a given block B of A_1 and G -conjugacy classes of blocks of a twisted group algebra for a subgroup of G (Theorem 4.5 below). A reduction to this invariant case is given in Section 2.

A starting point for the Clifford correspondence for blocks is provided by a more general correspondence for indecomposable modules discussed in Section 3. In this context, we let $B \uparrow^{A_1^{op} \otimes A}$ denote the induced $A_1^{op} \otimes A$ -module $B \otimes_{A_1^{op} \otimes A_1} (A_1^{op} \otimes A)$ and $\mathcal{E} = \text{End}_{A_1^{op} \otimes A}(B \uparrow^{A_1^{op} \otimes A})$. Then $\mathcal{E}$ is a (not necessarily fully) G -graded algebra; that is, $\mathcal{E} = \sum_{g \in G} \mathcal{E}_g$ is a direct sum of subspaces $\mathcal{E}_g$ with $\mathcal{E}_g \mathcal{E}_h \subseteq \mathcal{E}_{gh}$ for all $g, h \in G$. In Section 3 we describe the Miyashita action of G on $\mathcal{E}$ which permutes the blocks. Let $J_G(\mathcal{E})$ denote the graded Jacobson radical of $\mathcal{E}$, the intersection of all maximal graded right ideals. We show in Section 4 that $\mathcal{E}/J_G(\mathcal{E})$ is a twisted group algebra for a subgroup of G , and prove the following theorem.

Theorem 4.5 (Clifford correspondence). *Let B be a G -invariant block of A_1 . The blocks of A covering B correspond one-to-one with G -conjugacy classes of blocks of the twisted group algebra $\mathcal{E}/J_G(\mathcal{E})$.*

In order to prove the theorem, in Section 3 we relate blocks of A covering B to blocks of the endomorphism algebra $\mathcal{E}$ in the following way. Let Δ be the diagonal subalgebra $\sum_{g \in G} (A_{g^{-1}})^{op} \otimes A_g$ of $A^{op} \otimes A$. The identity component A_1 of A is naturally a Δ -module; we denote this Δ -module by $(A_1)_\Delta$. The $A^{op} \otimes A$ -module $(A_1)_\Delta \uparrow^{A^{op} \otimes A}$ induced from Δ is isomorphic to A . If the block B of A_1 is G -invariant, then B is naturally a Δ -module, and so we obtain an ideal direct summand $B_\Delta \uparrow^{A^{op} \otimes A} \cong ABA$ of A . We show in Section 2 that the blocks of A covering B are precisely the indecomposable direct summands of the $A^{op} \otimes A$ -module $B_\Delta \uparrow^{A^{op} \otimes A}$. In turn, these direct summands are in one-to-one correspondence with the blocks of another endomorphism algebra, $\text{End}_{A^{op} \otimes A}(B_\Delta \uparrow^{A^{op} \otimes A})$. This endomorphism algebra is isomorphic to the algebra $\mathcal{E}^G$ of fixed points of $\mathcal{E} = \text{End}_{A_1^{op} \otimes A}(B \uparrow^{A_1^{op} \otimes A})$, and blocks of $\mathcal{E}^G$ correspond to G -conjugacy classes of blocks of $\mathcal{E}$, as we show in Section 3.

Our results also hold in the more general situation where k is replaced by a p -modular system (K, R, k) for a prime p , where k is algebraically closed. Here R is a complete discrete valuation ring with maximal ideal $\mathfrak{p}$, quotient field K , and residue class field $k = R/\mathfrak{p}$ of characteristic p . The basic theory needed includes the Krull-Schmidt-Azumaya Theorem [7, Proposition 56.4] and existence of projective covers of modules which follows from [6, Theorem 6.23] and [7, Propositions 56.2 and 56.4]. In this situation, A is a fully G -graded R -algebra (free and finitely generated as an R -module), and the twisted group algebra $\mathcal{E}/J_G(\mathcal{E})$ of the Clifford correspondence is a k -algebra as $J_G(\mathcal{E}) \supseteq \mathfrak{p}\mathcal{E}$. For simplicity however, we state our results in the case of a single base field k . All algebras and modules will be finite dimensional over k , and tensor products will be over k unless otherwise indicated.

The author thanks J. L. Alperin for suggesting this work and P. R. Boisen, E. C. Dade, and H. Ellers for helpful conversations.

2. REDUCTION TO THE INVARIANT CASE

In this section we first give a correspondence between blocks covering a block B of A_1 and blocks of the endomorphism algebra $\text{End}_{A^{op} \otimes A}(B_{\Delta_B} \uparrow^{A^{op} \otimes A})$. We then give a reduction to the invariant case. Let G_B be the subgroup of G fixing an arbitrary block B of A_1 , that is,

$$G_B := \{g \in G \mid A_{g^{-1}} B A_g = B\}.$$

Let $A_B := \sum_{g \in G_B} A_g$ and $\Delta_B := \sum_{g \in G_B} (A^{op})_g \otimes A_g$, where $(A^{op})_g = (A_{g^{-1}})^{op}$. Then Δ_B is a fully $\delta(G_B)$ -graded algebra where $\delta(G_B) = \{(g, g) \mid g \in G_B\} \subseteq G \times G$. Note that B is naturally a Δ_B -module, denoted B_{Δ_B} . We observe that if $\tilde{B}$ is a block of A covering B , so that B is a direct summand of $\tilde{B} \downarrow_{A_1^{op} \otimes A_1}$, then $\tilde{B}$ covers all G -conjugates of B : The restricted module $\tilde{B} \downarrow_{A_1^{op} \otimes A_1}$ is fixed by the G -action, as $\tilde{B}$ is. So for any $g \in G$, $B^g = A_{g^{-1}} B A_g$ is also a direct summand of $\tilde{B} \downarrow_{A_1^{op} \otimes A_1}$. In addition $\tilde{B}$ covers no other blocks of A_1 , as we see in the following lemma. We refer the reader to [3], [10], or [18] for facts about conjugate modules and induced modules for a fully group-graded ring. The following lemma is analogous to [1, Lemma 4.3] or [7, Lemma 61.3 (v)].

Lemma 2.1. *Let B be a block of A_1 .*

- (i) *The blocks of A covering B correspond one-to-one with the indecomposable summands of the $A^{op} \otimes A$ -module $B_{\Delta_B} \uparrow^{A^{op} \otimes A}$.*
- (ii) *The blocks of A covering B correspond one-to-one with the blocks of the algebra $\text{End}_{A^{op} \otimes A}(B_{\Delta_B} \uparrow^{A^{op} \otimes A})$.*
- (iii) *If the block $\tilde{B}$ of A covers B , then $\tilde{B}$ covers all G -conjugates of B and no other blocks of A_1 .*

Proof. (i) Write $A_1 \cong B_1 \oplus \cdots \oplus B_k \oplus M$ as $A_1^{op} \otimes A_1$ -modules, where $B = B_1, \dots, B_k$ are the distinct conjugates of B , and M is the sum of the remaining blocks of A_1 . The induced $A^{op} \otimes A$ -module $(A_1)_\Delta \uparrow^{A^{op} \otimes A}$ is isomorphic to A [3, Lemma 3.3]; this isomorphism is given simply by sending an element $\sum_i a_i \otimes (b_i \otimes c_i)$ of $(A_1)_\Delta \uparrow^{A^{op} \otimes A}$ to $\sum_i b_i a_i c_i$. Similarly, $B_{\Delta_B} \uparrow^{A^{op} \otimes A} \cong A B A$ as $A^{op} \otimes A$ -modules. As $B_1 \oplus \cdots \oplus B_k$ and M are naturally Δ -submodules of $(A_1)_\Delta$, we have

$$A \cong (B_1 \oplus \cdots \oplus B_k)_\Delta \uparrow^{A^{op} \otimes A} \oplus M_\Delta \uparrow^{A^{op} \otimes A}$$

as $A^{op} \otimes A$ -modules. The Δ -module $(B_1 \oplus \cdots \oplus B_k)_\Delta$ is isomorphic to $B_{\Delta_B} \uparrow^\Delta$ by an argument similar to that used in [3, Lemma 3.3]. Therefore $B_{\Delta_B} \uparrow^{A^{op} \otimes A} \cong B_{\Delta_B} \uparrow^\Delta \uparrow^{A^{op} \otimes A} \cong (B_1 \oplus \cdots \oplus B_k)_\Delta \uparrow^{A^{op} \otimes A}$, and so $A \cong B_{\Delta_B} \uparrow^{A^{op} \otimes A} \oplus M_\Delta \uparrow^{A^{op} \otimes A}$.

Let $\tilde{B}$ be a block of A contained in $B_{\Delta_B} \uparrow^{A^{op} \otimes A}$. By Mackey's Theorem for group-graded rings [18, Theorem 8.4],

$$B_{\Delta_B} \uparrow^{A^{op} \otimes A} \downarrow_{A_1^{op} \otimes A_1} \cong \sum_{(s,t) \in \delta(G_B) \setminus G \times G} B \otimes_{A_1^{op} \otimes A_1} ((A^{op})_s \otimes A_t).$$

A similar argument to [3, Lemma 3.3] shows that the $A_1^{op} \otimes A_1$ -module $A_{s^{-1}} B A_t$ is isomorphic to the conjugate module $B \otimes_{A_1^{op} \otimes A_1} ((A^{op})_s \otimes A_t)$. As B is indecomposable and A is fully graded, these conjugate modules $A_{s^{-1}} B A_t$ are indecomposable as well. Since $\tilde{B}$ divides $B_{\Delta_B} \uparrow^{A^{op} \otimes A}$, the Krull-Schmidt Theorem now implies that some $A_{s^{-1}} B A_t$ divides $\tilde{B} \downarrow_{A_1^{op} \otimes A_1}$. Therefore B is a summand of $A_s(\tilde{B} \downarrow_{A_1^{op} \otimes A_1}) A_{t^{-1}} \cong \tilde{B} \downarrow_{A_1^{op} \otimes A_1}$; that is, $\tilde{B}$ covers B .

Now assume $\tilde{B}$ is a block of A covering B , but does not divide $B_{\Delta_B} \uparrow^{A^{op} \otimes A}$. Then $\tilde{B}$ divides $M_{\Delta} \uparrow^{A^{op} \otimes A}$, and so B divides $M_{\Delta} \uparrow^{A^{op} \otimes A} \downarrow_{A_1^{op} \otimes A_1}$. As before, we apply Mackey's Theorem to $M_{\Delta} \uparrow^{A^{op} \otimes A} \downarrow_{A_1^{op} \otimes A_1}$, and conclude that B divides $A_{s^{-1}} M A_t$ for some $s, t \in G$. This implies that the conjugate $A_1^{op} \otimes A_1$ -module $B^{s^{-1}} = A_s B A_{s^{-1}}$ divides $A_s A_{s^{-1}} M A_t A_{s^{-1}} = M A_{ts^{-1}}$. Letting e be the primitive central idempotent of A_1 corresponding to the block $B^{s^{-1}}$, we derive a contradiction, as $e B^{s^{-1}} \neq 0$ and $e M = 0$. Therefore the blocks of A covering B are exactly the indecomposable summands of $B_{\Delta_B} \uparrow^{A^{op} \otimes A}$.

(ii) As a summand of A , $B_{\Delta_B} \uparrow^{A^{op} \otimes A}$ is an algebra, and

$$\text{End}_{A^{op} \otimes A}(B_{\Delta_B} \uparrow^{A^{op} \otimes A}) \cong Z(B_{\Delta_B} \uparrow^{A^{op} \otimes A}),$$

the center of $B_{\Delta_B} \uparrow^{A^{op} \otimes A}$. Thus blocks of $B_{\Delta_B} \uparrow^{A^{op} \otimes A}$ correspond one-to-one with blocks of $\text{End}_{A^{op} \otimes A}(B_{\Delta_B} \uparrow^{A^{op} \otimes A})$.

(iii) This follows immediately from the proof of (i) and the observation preceding the lemma. $\square$

We next show that the blocks of A covering B correspond one-to-one with the blocks of A_B covering B , a reduction allowing us to consider only the G -invariant case from now on. This is a version of the Fong-Reynolds Theorem [17, V.2.5].

Lemma 2.2. *Let B be a block of A_1 . The blocks of A covering B correspond one-to-one with the blocks of A_B covering B . Further, this correspondence is given by induction of blocks of A_B , as $A_B^{op} \otimes A_B$ -modules, to $A^{op} \otimes A$ -modules.*

Proof. We prove the first statement by showing that there is an isomorphism of algebras

$$\text{End}_{A_B^{op} \otimes A_B}(B_{\Delta_B} \uparrow^{A_B^{op} \otimes A_B}) \cong \text{End}_{A^{op} \otimes A}(B_{\Delta_B} \uparrow^{A^{op} \otimes A}),$$

and applying Lemma 2.1 (ii). By the Nakayama relations [2, Proposition 2.8.3] we have two natural isomorphisms

$$\text{End}_{A_B^{op} \otimes A_B}(B_{\Delta_B} \uparrow^{A_B^{op} \otimes A_B}) \cong \text{Hom}_{\Delta_B}(B_{\Delta_B}, B_{\Delta_B} \uparrow^{A_B^{op} \otimes A_B} \downarrow_{\Delta_B})$$

and

$$\text{End}_{A^{op} \otimes A}(B_{\Delta_B} \uparrow^{A^{op} \otimes A}) \cong \text{Hom}_{\Delta_B}(B_{\Delta_B}, B_{\Delta_B} \uparrow^{A^{op} \otimes A} \downarrow_{\Delta_B}).$$

We may consider $B_{\Delta_B} \uparrow^{A_B^{op} \otimes A_B} \downarrow_{\Delta_B}$ as a direct summand of $B_{\Delta_B} \uparrow^{A^{op} \otimes A} \downarrow_{\Delta_B}$ by Mackey's Theorem [18, Theorem 8.4]. In order to achieve the desired isomorphism, we need only show that all Δ_B -homomorphisms from B_{Δ_B} to $B_{\Delta_B} \uparrow^{A^{op} \otimes A} \downarrow_{\Delta_B}$ in fact have image contained in $B_{\Delta_B} \uparrow^{A_B^{op} \otimes A_B} \downarrow_{\Delta_B}$. Consider a summand $A_s B A_t = A_s B A_{s^{-1}} A_{st} = B^{s^{-1}} A_{st}$ of $B_{\Delta_B} \uparrow^{A^{op} \otimes A} \downarrow_{\Delta_B}$ as an $A_1^{op} \otimes A_1$ -module. If $s \notin G_B$, then the primitive central idempotent e of A_1 associated with B yields the identity map on B but 0 on $B^{s^{-1}} \neq B$. In this case there are no $A_1^{op} \otimes A_1$ -homomorphisms, and so no Δ_B -homomorphisms from B_{Δ_B} to $A_s B A_t$. A similar argument works if $s \in G_B$ and $t \notin G_B$. Therefore we have proved the first statement of the lemma.

By Lemma 2.1 (i), blocks of A_B covering B are the indecomposable $A_B^{op} \otimes A_B$ -direct summands of $B_{\Delta_B} \uparrow^{A_B^{op} \otimes A_B}$, while blocks of A covering B are the indecomposable $A^{op} \otimes A$ -direct summands of

$$B_{\Delta_B} \uparrow^{A^{op} \otimes A} \cong (B_{\Delta_B} \uparrow^{A_B^{op} \otimes A_B}) \uparrow^{A^{op} \otimes A}.$$

Suppose that $B_{\Delta_B} \uparrow^{A_B^{op} \otimes A_B} \cong \tilde{B}_1 \oplus \cdots \oplus \tilde{B}_n$ is a decomposition into indecomposable $A_B^{op} \otimes A_B$ -modules. Then

$$B_{\Delta_B} \uparrow^{A^{op} \otimes A} \cong \tilde{B}_1 \uparrow^{A^{op} \otimes A} \oplus \cdots \oplus \tilde{B}_n \uparrow^{A^{op} \otimes A}$$

as $A^{op} \otimes A$ -modules. If some $\tilde{B}_i \uparrow^{A^{op} \otimes A}$ were not indecomposable, there would be a contradiction to the first statement of the lemma. Therefore each $\tilde{B}_i \uparrow^{A^{op} \otimes A}$ is a block of A . $\square$

3. TWO ENDOMORPHISM ALGEBRAS

In this section we assume B is a G -invariant block of A_1 , so that $G_B = G$, $A_B = A$, and $\Delta_B = \Delta = \sum_{g \in G} (A^{op})_g \otimes A_g$.

We first discuss a Clifford correspondence for *indecomposable* modules which is implicit in Dade's work [14]. This provides a starting point for the Clifford correspondence for blocks. We show that an endomorphism algebra arising in this indecomposable module situation (as applied to a block) is in fact the fixed point subalgebra of the endomorphism algebra of Lemma 2.1 (ii) under an appropriate G -action. It seems necessary to look at both endomorphism algebras to achieve the Clifford correspondence for blocks from this point of view. Moreover this development clarifies the connection to the classical Clifford correspondence, and explains why G -conjugacy classes arise in the Clifford correspondence for blocks.

Let V be an indecomposable right A_1 -module, and $\mathcal{E} = \text{End}_A(V \uparrow^A)$, where $V \uparrow^A = V \otimes_{A_1} A$ is the induced A -module. As $V \uparrow^A$ is a graded module [20], $\mathcal{E}$ is a G -graded algebra with identity component $\mathcal{E}_1 \cong \text{End}_{A_1}(V)$. One way to see this is to consider the isomorphisms of vector spaces

$$\mathcal{E} \cong \text{Hom}_{A_1}(V, V \uparrow^A \downarrow_{A_1}) \cong \sum_{g \in G} \text{Hom}_{A_1}(V, V^g)$$

that follow from [2, Proposition 2.8.3] and Mackey's Theorem [18, Theorem 8.4]. Here V^g is the A_1 -module $V \otimes_{A_1} A_g$, and $\mathcal{E}_g \cong \text{Hom}_{A_1}(V, V^g)$ as a vector space. If V is G -invariant, so that $V^g \cong V$ for all $g \in G$, then $\mathcal{E}$ is *fully* G -graded, as may be seen from the above decomposition of $\mathcal{E}$ or [20, I.5.2]. If V is also irreducible, then Schur's Lemma implies that $\mathcal{E}$ is a twisted group algebra. This is the twisted group algebra of the classical Clifford correspondence. In case V is indecomposable but not irreducible, or in case k is replaced by a complete discrete valuation ring R , the fact that $\mathcal{E}_1 \cong \text{End}_{A_1}(V)$ is local [6, Proposition 6.10] replaces Schur's Lemma as follows: Let $J_G(\mathcal{E})$ denote the *graded Jacobson radical* of $\mathcal{E}$, the intersection of all maximal graded right ideals. By [5, Theorem 4.4] $J_G(\mathcal{E})$ is contained in the ordinary Jacobson radical $J(\mathcal{E})$. In case V is G -invariant, [14, Proposition 2.19], [6, Proposition 5.22], and [8, Lemma 14.2] imply that $\mathcal{E}/J_G(\mathcal{E})$ is a twisted group algebra for a subgroup of G . For the sake of completeness, we will give these arguments in greater detail later for the special case where V is a block.

Whether or not V is G -invariant, there is a one-to-one correspondence between indecomposable A -direct summands of $V \uparrow^A$ and indecomposable right $\mathcal{E}/J_G(\mathcal{E})$ -direct summands of $\mathcal{E}/J_G(\mathcal{E})$: Indecomposable summands of $V \uparrow^A$ correspond one-to-one with indecomposable right summands of $\mathcal{E}$ by sending a summand of $V \uparrow^A$ to the corresponding projection endomorphism, an idempotent of $\mathcal{E}$ [6, Proposition 6.3]. As $J_G(\mathcal{E}) \subseteq J(\mathcal{E})$, indecomposable right $\mathcal{E}$ -direct summands of $\mathcal{E}$ correspond with indecomposable right $\mathcal{E}/J_G(\mathcal{E})$ -direct summands of $\mathcal{E}/J_G(\mathcal{E})$ [6, Corollaries 6.22 and 6.25].

In the block situation, we replace A by the fully G -graded algebra $A_1^{op} \otimes A$, A_1 by $A_1^{op} \otimes A_1$, and V by the block B of A_1 . (We remark that replacing A by $A^{op} \otimes A_1$ would work equally well.) The relevant G -graded endomorphism algebra is then

$$\mathcal{E} := \text{End}_{A_1^{op} \otimes A}(B \uparrow^{A_1^{op} \otimes A}),$$

and the Clifford correspondence for indecomposable modules described above applies here. It will need to be modified to yield the Clifford correspondence for blocks given in Theorem 4.5. We caution that, although B is G -invariant as a block, it may not be $1 \times G$ -invariant as an $A_1^{op} \otimes A_1$ -module. Thus it does not follow from the above arguments that $\mathcal{E}/J_G(\mathcal{E})$ is a twisted group algebra; in Section 4 we show that this fact follows from a characterization of the graded Jacobson radical given in [14]. In addition, we are interested in $A^{op} \otimes A$ -modules rather than $A_1^{op} \otimes A$ -modules. This is where a G -action on $\mathcal{E}$ arises. The following proposition is Theorem 1.3 of [19]; see also [11, Theorem 2.1].

Proposition 3.1 (Miyashita action). *Let G be a finite group and S a fully G -graded algebra. Let M and N be S -modules, $g \in G$, and $\phi \in \text{Hom}_{S_1}(M, N)$. Then there is a unique element $\phi^g \in \text{Hom}_{S_1}(M, N)$ such that $\phi^g(ms_g) = \phi(m)s_g$ for all $m \in M$ and $s_g \in S_g$. If $M = N$, then G acts as algebra automorphisms of $\text{End}_{S_1}(M)$.*

We give the definition of ϕ^g , which will be needed in the next section. As S is fully G -graded, we have $S_{g^{-1}}S_g = S_1$ for all $g \in G$. As $1 \in S_1$ [20], there are elements $\alpha_i \in S_{g^{-1}}$, $\beta_i \in S_g$ such that $\sum_{i=1}^n \alpha_i \beta_i = 1$. Then, for all $m \in M$,

$$(3.1) \quad \phi^g(m) := \sum_{i=1}^n \phi(m\alpha_i)\beta_i.$$

Now let $S = A^{op} \otimes A$ be the fully G -graded algebra with $S_g = (A^{op})_g \otimes A$. The proposition gives a G -action on $\text{End}_{A_1^{op} \otimes A}(B_\Delta \uparrow^{A^{op} \otimes A})$, where $\Delta = \sum_{g \in G} (A^{op})_g \otimes A_g$, and B_Δ is the Δ -module B . This provides the connection between $\mathcal{E} = \text{End}_{A_1^{op} \otimes A}(B \uparrow^{A_1^{op} \otimes A})$ and the algebra $\text{End}_{A^{op} \otimes A}(B_\Delta \uparrow^{A^{op} \otimes A})$ of Lemma 2.1 (ii), as we see next. We note that $B_\Delta \uparrow^{A^{op} \otimes A} \downarrow_{A_1^{op} \otimes A} \cong B \uparrow^{A_1^{op} \otimes A}$, where the latter module is induced from $A_1^{op} \otimes A_1$. This follows from an application of Mackey's Theorem [18, Theorem 8.4], as there is only one $\delta(G), 1 \times G$ -double coset in $G \times G$, and $B_\Delta \downarrow_{A_1^{op} \otimes A_1} = B$. Therefore

$$\text{End}_{A_1^{op} \otimes A}(B_\Delta \uparrow^{A^{op} \otimes A}) \cong \text{End}_{A_1^{op} \otimes A}(B \uparrow^{A_1^{op} \otimes A}) = \mathcal{E}.$$

Thus the above proposition yields an action of G as automorphisms of $\mathcal{E}$. The next lemma describing the fixed point subalgebra $\mathcal{E}^G$ follows immediately.

Lemma 3.2. $\mathcal{E}^G \cong \text{End}_{A^{op} \otimes A}(B_\Delta \uparrow^{A^{op} \otimes A})$.

Note that the second endomorphism algebra in the lemma is the algebra of Lemma 2.1 (ii). We mention that if we had taken $S_g = A^{op} \otimes A_g$ instead, we would have gotten a different G -graded endomorphism algebra, but upon taking G -fixed points, we would have obtained the same $A^{op} \otimes A$ -endomorphism algebra.

We now give the relationship between our endomorphism algebras and the centralizer algebras of [9]. Let $C_A(A_1)$ denote the centralizer in A of A_1 . Let e be the primitive central idempotent of A_1 corresponding to the block B . Then as graded algebras, $\mathcal{E}$ is isomorphic to $eC_A(A_1)$: We identify $B \uparrow^{A_1^{op} \otimes A}$ with

$\sum_{g \in G} BA_g$ as in the proof of Lemma 2.1 (i). Given $\phi \in \mathcal{E}$, ϕ is determined by $\phi(e)$, as $\phi(b) = \phi(eb) = \phi(e)b$ for any $b \in B \uparrow^{A_1^{op} \otimes A}$. Further, $\phi(e) \in eC_A(A_1)$ as $e\phi(e) = \phi(e^2) = \phi(e)$ and $\phi \in \mathcal{E}$ implies that $\phi(e)$ commutes with elements of A_1 . Conversely, any element of $eC_A(A_1)$ defines an element of $\mathcal{E}$ in this way. The G -grading on $eC_A(A_1)$ inherited from A corresponds to that of $\mathcal{E}$, and the G -action on $\mathcal{E}$ provided by Proposition 3.1 gives rise to a G -action on $eC_A(A_1)$, which is just the usual Miyashita action on this algebra.

Lemma 3.3. *The subalgebras $\mathcal{E}^G$ and $\mathcal{E}_1$ of $\mathcal{E}$ are both contained in the center of $\mathcal{E}$.*

Proof. Let $\phi \in \mathcal{E}$ and $\psi \in \mathcal{E}_g$ for some $g \in G$, and suppose $\psi(e) = r_g \in eC_A(A_1)$, so that $\psi(b) = r_gb$ for all $b \in B \uparrow^{A_1^{op} \otimes A}$. Applying Proposition 3.1, $\phi^{g^{-1}} \circ \psi(e) = \phi^{g^{-1}}(r_ge) = r_g\phi(e) = \psi \circ \phi(e)$, so that $\phi^{g^{-1}} \circ \psi = \psi \circ \phi$. If $\phi \in \mathcal{E}^G$, then ϕ is in the center of $\mathcal{E}$. If $\psi \in \mathcal{E}_1$, then $\phi \circ \psi = \psi \circ \phi$ for all $\phi \in \mathcal{E}$, so that ψ is in the center of $\mathcal{E}$. $\square$

The next lemma shows that blocks of $\mathcal{E}^G$ correspond one-to-one to G -conjugacy classes of blocks of $\mathcal{E}$, as $\mathcal{E}^G$ is central in $\mathcal{E}$ by the previous lemma. Combined with Lemma 2.1 (ii) for $G = G_B$, this shows that blocks of A covering B correspond one-to-one with G -conjugacy classes of blocks of $\mathcal{E}$.

Lemma 3.4. *Let S be a finite dimensional k -algebra with an action of G as automorphisms, and suppose that S^G is central in S . Then the blocks of S^G correspond one-to-one with the G -conjugacy classes of blocks of S .*

Proof. A G -conjugacy class of blocks of S corresponds to a G -conjugacy class $e_1, \dots, e_k$ of primitive central idempotents of S . Their sum $e' = e_1 + \dots + e_k$ is a central idempotent of S^G . If e' were not primitive in S^G , then it would decompose into a sum of primitive central idempotents in S^G , each of which is a central idempotent of S as S^G is central in S . Thus each decomposes into a sum of primitive central idempotents of S , and by uniqueness these must be the $e_1, \dots, e_k$. This contradicts the assumption that $e_1, \dots, e_k$ is a single G -conjugacy class. Conversely, a block of S^G corresponds to a primitive central idempotent of S^G , which is a central idempotent of S , and decomposes into primitive central idempotents of S . The G -action must permute these primitive central idempotents transitively. $\square$

4. THE CLIFFORD CORRESPONDENCE

We continue under the assumption that B is a G -invariant block of A_1 . In this section we show that blocks of $\mathcal{E} = \text{End}_{A_1^{op} \otimes A}(B \uparrow^{A_1^{op} \otimes A})$ correspond one-to-one with blocks of $\mathcal{E}/J_G(\mathcal{E})$, a version of [9, Lemma 3.1 and Theorem 3.5]. Together with the results of Sections 2 and 3, this yields the Clifford correspondence in Theorem 4.5. In order to achieve this correspondence, it seems necessary to introduce a *fully* group-graded subalgebra $\mathcal{E}[B]$ of $\mathcal{E}$, as is done for centralizer algebras in [9].

Let $G[B]$ be the subgroup of G defined by $g \in G[B]$ if and only if $\mathcal{E}_g$ contains a unit, and

$$\mathcal{E}[B] := \mathcal{E}_{G[B]} = \sum_{g \in G[B]} \mathcal{E}_g.$$

Then $\mathcal{E}[B]$ is a *fully* $G[B]$ -graded algebra, as for each $g \in G[B]$, $\mathcal{E}_g = u_g \mathcal{E}_1$ for a unit $u_g \in \mathcal{E}_g$. There are examples for which $G[B] \neq G$ [16, Lemma 3.11].

The graded Jacobson radical $J_G(\mathcal{E})$ of any group-graded algebra $\mathcal{E}$ is a graded two-sided ideal of $\mathcal{E}$ [20, Lemma I.7.4]. We give the characterization of $J_G(\mathcal{E})$ in [14], providing details for the sake of completeness. The components of $J_G(\mathcal{E})$ are given by

$$J_G(\mathcal{E})_g = \{\phi \in \mathcal{E}_g \mid \phi\mathcal{E}_{g^{-1}} \subseteq J(\mathcal{E}_1)\}.$$

This follows immediately from the one-to-one correspondence between all maximal G -graded right ideals M of $\mathcal{E}$ and all maximal right ideals N of $\mathcal{E}_1$ given by sending M to $N = M_1$, and N to M where $M_g = \{\phi \in \mathcal{E}_g \mid \phi\mathcal{E}_{g^{-1}} \subseteq N\}$. This correspondence is straightforward to verify. As B is an indecomposable $A_1^{op} \otimes A_1$ -module, $\mathcal{E}_1 \cong \text{End}_{A_1^{op} \otimes A_1}(B)$ is local, and so $\mathcal{E}_g - J_G(\mathcal{E})_g$ is the set of units in $\mathcal{E}_g$ [14, Lemma 2.18]. It follows that

$$J_G(\mathcal{E})_g = \begin{cases} J(\mathcal{E}_1)\mathcal{E}_g & \text{if } g \in G[B], \\ \mathcal{E}_g & \text{if } g \notin G[B], \end{cases}$$

as in [14, Proposition 2.19]. These arguments also apply to $\mathcal{E}[B]$, so in that case $J_{G[B]}(\mathcal{E}[B]) = J(\mathcal{E}_1)\mathcal{E}[B]$, and $\mathcal{E}[B]/J_{G[B]}(\mathcal{E}[B])$ is a twisted group algebra for $G[B]$ by [8, Lemma 14.2]. The above characterizations of $J_G(\mathcal{E})$ and $J_{G[B]}(\mathcal{E}[B])$ also immediately imply the following lemma.

Lemma 4.1. *There is an isomorphism of twisted group algebras*

$$\mathcal{E}/J_G(\mathcal{E}) \cong \mathcal{E}[B]/J_{G[B]}(\mathcal{E}[B]).$$

The next lemma, a generalization of [9, Lemma 3.1], may be applied to our situation with $H = G[B]$ and $S = \mathcal{E}[B]$ as $\mathcal{E}_1$ is central in $\mathcal{E}[B]$ by Lemma 3.3, and $J_{G[B]}(\mathcal{E}[B]) = J(\mathcal{E}_1)\mathcal{E}[B]$. This shows the necessity of dealing with $\mathcal{E}[B]$ rather than $\mathcal{E}$.

Lemma 4.2. *Let H be a finite group, and S a fully H -graded algebra in which S_1 is central and $J_H(S) = J(S_1)S$. Then the blocks of S correspond one-to-one with blocks of $S/J_H(S)$.*

Proof. Let $S/J_H(S) = b_1 \oplus \cdots \oplus b_k$ be a decomposition into blocks. Consider each b_i as a right S -module via the quotient map from S to $S/J_H(S)$. Let $P(b_i)$ be the projective cover of b_i as a right S -module. As $J_H(S) \subseteq J(S)$ [5, Theorem 4.4], S itself is the projective cover of $S/J_H(S)$ as a right S -module [6, Corollary 6.22]. As projective covers preserve direct sums, we have $S \cong P(b_1) \oplus \cdots \oplus P(b_k)$ as a right S -module. We identify the $P(b_i)$ with right ideals of S via this isomorphism. We will show that the $P(b_i)$ are also left ideals, and that they are indecomposable $S^{op} \otimes S$ -modules; that is, the $P(b_i)$ are blocks of S .

We let $P'(b_i)$ be the projective cover of b_i as a left S -module, so that $S \cong P'(b_1) \oplus \cdots \oplus P'(b_k)$ as a left S -module. Consider the sum m_i of all blocks b_j with $j \neq i$. We have $S \cong P(b_i) \oplus P(m_i)$ as a right S -module, and $S \cong P'(b_i) \oplus P'(m_i)$ as a left S -module. Restrict these S -modules to S_1 -modules. As S_1 is central in S , we do not distinguish between left and right S_1 -modules. We have $P(b_i) = P(b_i)S \cong P(b_i)P'(b_i) \oplus P(b_i)P'(m_i)$ and $P(m_i) \cong P(m_i)P'(b_i) \oplus P(m_i)P'(m_i)$ as S_1 -modules. Therefore, as S_1 -modules,

$$S \cong P(b_i)P'(b_i) \oplus P(b_i)P'(m_i) \oplus P(m_i)P'(b_i) \oplus P(m_i)P'(m_i).$$

The right S_1 -module $P(b_i)P'(m_i)$ becomes 0 on passing to the quotient $S/J_H(S)$. As $J_H(S) = J(S_1)S$, it follows that $J(S_1)P(b_i)P'(m_i) = P(b_i)P'(m_i)$. Therefore

by Nakayama's Lemma, $P(b_i)P'(m_i) = 0$, and similarly $P(m_i)P'(b_i) = 0$. Thus

$$S \cong P(b_i)P'(b_i) \oplus P(m_i)P'(m_i)$$

as S_1 -modules, where $P(b_i) = P(b_i)P'(b_i) = P'(b_i)$, and $P(m_i) = P(m_i)P'(m_i) = P'(m_i)$. It follows that each $P(b_i)$ is both a left and a right ideal of S .

Suppose $P(b_i) = B_1 \oplus B_2$ as $S^{op} \otimes S$ -modules. Under the quotient map from S to $S/J_H(S)$, the image of one of B_1, B_2 must be 0. Considering B_1 and B_2 as S_1 -modules, another application of Nakayama's Lemma implies that one of B_1, B_2 is 0. $\square$

Next we see that G -conjugacy classes of blocks of $\mathcal{E}/J_G(\mathcal{E})$ correspond one-to-one with G -conjugacy classes of blocks of $\mathcal{E}$, resulting in the Clifford correspondence. For this, we need a lemma.

Lemma 4.3. *The subgroup $G[B]$ is normal in G , and $(\mathcal{E}_h)^g = \mathcal{E}_{hg}$ for all $g, h \in G$.*

Proof. Let $g, h \in G$, and $\phi \in \mathcal{E}_h$. By (3.1), $\phi^g(b) = \sum_i \beta_i \phi(\alpha_i b)$ for all $b \in B \uparrow^{A_1^{op} \otimes A}$, where $\alpha_i \in A_g$, $\beta_i \in A_{g^{-1}}$, and $\sum_i \beta_i \alpha_i = 1$. Let e be the primitive central idempotent of A_1 corresponding to the block B , so $\phi(e) \in A_h$ as discussed in the text preceding Lemma 3.3. As e is a G -invariant element of $C_A(A_1)$, e is central in A_1 so $\phi^g(e) = \sum_i \beta_i \phi(\alpha_i e) = \sum_i \beta_i \phi(e) \alpha_i$, since ϕ is an $A_1^{op} \otimes A$ -map. But $\sum_i \beta_i \phi(e) \alpha_i \in A_{g^{-1}} A_h A_g = A_{hg}$. Therefore $\phi^g \in \mathcal{E}_{hg}$, and $(\mathcal{E}_h)^g \subseteq \mathcal{E}_{hg}$. Conversely, let $\psi \in \mathcal{E}_{hg}$, and let $\phi = \psi^{g^{-1}} \in (\mathcal{E}_{hg})^{g^{-1}} \subseteq \mathcal{E}_h$ by the above argument. Then $\psi = \phi^g \in (\mathcal{E}_h)^g$, and so $\mathcal{E}_{hg} = (\mathcal{E}_h)^g$.

To see that $G[B]$ is normal in G , let $h \in G[B]$, $g \in G$, and u_h a unit in $\mathcal{E}_h$. Then $(u_h)^g$ is a unit in $\mathcal{E}_{hg}$. $\square$

By the lemma, the G -action on $\mathcal{E}$ yields a G -action on $\mathcal{E}[B]$. Further, G fixes the ideal $J_{G[B]}(\mathcal{E}[B]) = J(\mathcal{E}_1)\mathcal{E}[B]$, as G fixes $\mathcal{E}_1$ by the lemma and so permutes the maximal ideals of $\mathcal{E}_1$. Therefore the G -action on $\mathcal{E}[B]$ induces a G -action on $\mathcal{E}[B]/J_{G[B]}(\mathcal{E}[B])$.

Lemma 4.4. *The G -conjugacy classes of blocks of $\mathcal{E}$ correspond one-to-one with G -conjugacy classes of blocks of $\mathcal{E}/J_G(\mathcal{E})$.*

Proof. By Lemmas 4.1 and 4.2, we need only show that G -conjugacy classes of blocks of $\mathcal{E}[B]$ correspond one-to-one with G -conjugacy classes of blocks of $\mathcal{E}$. Given a G -conjugacy class of blocks of $\mathcal{E}[B]$, there is a corresponding G -invariant central idempotent e of $\mathcal{E}[B]$, which is also then a G -invariant central idempotent of $\mathcal{E}$, as $\mathcal{E}^G$ is central in $\mathcal{E}$. Consider the image $\bar{e}$ of e in $\mathcal{E}/J_G(\mathcal{E}) \cong \mathcal{E}[B]/J_{G[B]}(\mathcal{E}[B])$. If e may be decomposed in $\mathcal{E}$ as the sum of two nontrivial G -invariant central idempotents, this decomposition gives such a decomposition of $\bar{e}$, contradicting Lemma 4.2.

Conversely, a G -conjugacy class of blocks of $\mathcal{E}$ corresponds to a G -invariant central idempotent e of $\mathcal{E}$. This corresponds to an idempotent $\bar{e}$ of $\mathcal{E}/J_G(\mathcal{E}) \cong \mathcal{E}[B]/J_{G[B]}(\mathcal{E}[B])$, which by Lemma 4.2 corresponds to an idempotent e' of $\mathcal{E}[B]$. As $J_{G[B]}(\mathcal{E}[B]) \subseteq J_G(\mathcal{E})$, e and e' are central idempotents of $\mathcal{E}$ having the same image modulo $J_G(\mathcal{E})$, and so $e = e'$. $\square$

The Clifford correspondence now follows immediately.

Theorem 4.5. *Let B be a G -invariant block of A_1 . The blocks of A covering B correspond one-to-one with G -conjugacy classes of blocks of the twisted group algebra $\mathcal{E}/J_G(\mathcal{E})$.*

Proof. By Lemma 2.1 (ii) with $G = G_B$, Lemma 3.2, and Lemma 3.4, the blocks of A covering B correspond one-to-one with G -conjugacy classes of blocks of $\mathcal{E}$. By Lemma 4.4, G -conjugacy classes of blocks of $\mathcal{E}$ correspond one-to-one with G -conjugacy classes of blocks of $\mathcal{E}/J_G(\mathcal{E})$. $\square$

We close by returning to the general case where B is a block of A_1 that is *not* necessarily G -invariant, and G_B is the set of all elements g of G such that $A_{g^{-1}}BA_g = B$. By Lemma 2.2 and Theorem 4.5 with G replaced by G_B , we have

Corollary 4.6. *Let B be a block of A_1 and $\mathcal{E} = \text{End}_{A_1^{op} \otimes A_B}(B \uparrow^{A_1^{op} \otimes A_B})$. The blocks of A covering B correspond one-to-one with G_B -conjugacy classes of blocks of the twisted group algebra $\mathcal{E}/J_{G_B}(\mathcal{E})$.*

REFERENCES

- [1] J. L. ALPERIN AND D. W. BURRY, Block theory with modules, *J. Algebra* **65** (1980), pp. 225–233. MR **81k**:20018
- [2] D. J. BENSON, *Representations and Cohomology I: Basic representation theory of finite groups and associative algebras*, Cambridge University Press, 1991. MR **92m**:20005
- [3] P. R. BOISEN, The representation theory of fully group-graded algebras, *J. Algebra* **151** (1992), pp. 160–179. MR **93i**:20005
- [4] A. H. CLIFFORD, Representations induced in an invariant subgroup, *Ann. of Math. (2)* **38** (1937), pp. 533–550.
- [5] M. COHEN AND S. MONTGOMERY, Group-graded rings, smash products, and group actions, *Trans. Amer. Math. Soc.* **282** (1984), pp. 237–257. MR **85i**:16002
- [6] C. W. CURTIS AND I. REINER, *Methods of Representation Theory with Applications to Finite Groups and Orders, Volume I*, Wiley, 1981. MR **82i**:20001
- [7] ———, *Methods of Representation Theory with Applications to Finite Groups and Orders, Volume II*, Wiley, 1987. MR **88f**:20002
- [8] E. C. DADE, Compounding Clifford’s theory, *Ann. of Math.* **91** (1970), pp. 236–290. MR **41**:6992
- [9] ———, Block extensions, *Illinois J. Math.* **17** (1973), pp. 198–272. MR **48**:6226
- [10] ———, Group-graded rings and modules, *Math. Z.* **174** (1980), pp. 241–262. MR **82c**:16028
- [11] ———, The equivalence of various generalizations of group rings and modules, *Math. Z.* **181** (1982), pp. 335–344. MR **84a**:16018
- [12] ———, Counting characters in blocks, I, *Inv. Math.* **109** (1992), pp. 187–210. MR **93g**:20021
- [13] ———, Counting characters in blocks, II, *J. reine angew. Math.* **448** (1994), pp. 97–190. MR **95a**:20007
- [14] ———, Clifford theory for blocks. Preprint, 1995.
- [15] ———, Counting characters in blocks, 2.9, in *Representation Theory of Finite Groups*, Proceedings of a Special Research Quarter at the Ohio State University, de Gruyter, 1997. MR **99b**:20016
- [16] H. ELLERS, Cliques of irreducible representations, quotient groups, and Brauer’s theorems on blocks, *Can. J. Math.* **47** (5) (1995), pp. 929–945. MR **96g**:20013
- [17] W. FEIT, *The Representation Theory of Finite Groups*, North-Holland, 1982. MR **83g**:20001
- [18] J. HAEFNER, Graded equivalence theory with applications, *J. Algebra* **172** (1995), pp. 385–424. MR **96f**:16052
- [19] Y. MIYASHITA, On Galois extensions and crossed products, *J. Fac. Sci. Hokkaido Univ. Ser. I* **21** (1970), pp. 97–121. MR **42**:6046
- [20] C. NÄSTÄSCU AND F. VAN OYSTAEYEN, *Graded Ring Theory*, North-Holland, 1982. MR **84i**:16002

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF TORONTO, TORONTO, ONTARIO, CANADA M5S 3G3

E-mail address: `sjw@math.toronto.edu`

Current address: Department of Mathematics, University of Wisconsin, Madison, Wisconsin 53706

E-mail address: `sjw@math.wisc.edu`