ON AUTOMATIC CONTINUITY OF HOMOMORPHISMS

A. BEDDAA, S. J. BHATT, AND M. OUDADESS

(Communicated by Dale Alspach)

ABSTRACT. Introducing a weaker notion of regularity in a topological algebra, we examine and improve an automatic continuity theorem given by the second author. Examples and applications are given.

All topological algebras considered are commutative and Hausdorff, having a unit element. A topological algebra A is a Q -algebra if the set $G(A)$ of invertible elements is open. Let B be a subalgebra of an algebra A . Then B is *inverse closed* in A if $G(B) = B \cap G(A)$. A is *strongly semisimple* if for every $x \in A$, $x \neq 0$, there exists a nonzero continuous multiplicative linear functional χ such that $\chi(x) \neq 0$. A is *advertibly complete* if a Cauchy net x_α in A converges in A whenever for some y in A , $x_\alpha + y - x_\alpha y$ converges to 0. A Q -algebra is advertibly complete [Ma, p. 45]. A *uniform seminorm* on an algebra A is a seminorm p such that $p(x^2) = p(x)^2$ for all x in A . Such a p is submultiplicative [BK]. A *uniform topological algebra* A is a topological algebra whose topology is defined by a family of uniform seminorms. Such an A is semisimple. The abbreviation *lmca* will stand for locally m -convex algebra.

In [B], the following is given.

Theorem ([B, Theorem 2.2]). *Let A be a spectrally bounded, regular, complete, uniform topological algebra. If B is an lmca and $\phi : A \rightarrow B$ is a one-to-one homomorphism such that $(\text{Im } \phi)^-$ (the closure of $\text{Im } \phi$) is a semisimple Q -algebra, then $\phi^{-1}/\text{Im } \phi$ is continuous.*

In the proof, the author considers the map $\phi^* : \sigma(C) \rightarrow \sigma(A)$, with $\phi^*(f) = f \circ \phi$, $\sigma(C)$ and $\sigma(A)$ denoting respectively the spaces of nonzero continuous multiplicative functionals on $C = (\text{Im } \phi)^-$ and A . In Math. Reviews, the reviewer R. J. Loy [L] asserted that the continuity of ϕ has been implicitly used in [B]. Indeed, ϕ^* is not always well defined when ϕ is not continuous as the following example shows.

Example 1. Let Ω denote the first uncountable ordinal and $[0, \Omega)$ the set of all ordinals smaller than Ω . Consider the algebra $C[0, \Omega)$ of complex continuous functions on $[0, \Omega)$ with compact open topology τ . Every $f \in C[0, \Omega)$ is bounded. It is a regular uniform lmca. The identity map $\phi : (C[0, \Omega), \tau) \rightarrow (C[0, \Omega), \|\cdot\|_\infty)$, $\phi(f) = f$, satisfies the hypotheses of the above statement. It is well known that $(C[0, \Omega), \tau)$ has discontinuous multiplicative linear functionals (see, for example, [Mi], [Z]); let χ

Received by the editors September 10, 1997 and, in revised form, May 18, 1998.

1991 *Mathematics Subject Classification.* Primary 46H40; Secondary 46H05.

Key words and phrases. Uniform topological algebra, locally m -convex algebra, weakly regular algebra, advertibly complete algebra, Q -algebra, automatic continuity of homomorphism.

be such a functional. It is in $\sigma(C[0, \Omega], \|\cdot\|_\infty)$, but $\phi^*(\chi) = \chi$ is not in $\sigma([0, \Omega], \tau)$. One may give another example in a more general situation. Let X be a non-compact, locally compact pseudocompact space. Take $A = C(X)$ the algebra of all continuous functions on X with compact open topology, B the algebra $C(X)$ endowed with the sup norm $\|\cdot\|_\infty$ and $\phi : A \rightarrow B$ be $\phi(f) = f$. Since X is pseudocompact, non-compact, it is not realcompact [S, p. 44]. Let y be an element of the real compactification of X with y not in X . The evaluation δ_y at y is in $\sigma(B)$ but not in $\sigma(A)$. Hence ϕ^* is not defined on δ_y .

On the other hand, ϕ^* can be well defined even when ϕ is not continuous; in this case, the proof given in [B] works. Here is an example of such a situation.

Example 2. Let X be a compact Hausdorff space. Consider the algebra $C(X)$ of continuous complex functions on X . Take $A = (C(X), \tau_d)$, τ_d being the topology of uniform convergence on all countable compact subsets of X , B the algebra $(C(X), \|\cdot\|_\infty)$ and ϕ the identity map from A to B . It is of course discontinuous, only if X is uncountable. But in this case, $\sigma(A)$ and $\sigma(B)$ are both homeomorphic to X . So ϕ^* is well defined.

The following theorem repairs the above result. On one hand, it provides a positive result in a context more general than above; on the other hand, it shows that if one assumes the continuity of ϕ in the above result, then the algebra A is necessarily a Banach algebra.

Theorem 1. *Let A be weakly regular, advertibly complete, uniform topological algebra, let B be an lmca, and let $\phi : A \rightarrow B$ be a one-to-one homomorphism such that $(\text{Im } \phi)^-$ is a semisimple Q -algebra.*

- (1) *If A is functionally continuous, then $\phi^{-1}/\text{Im } \phi$ is continuous.*
- (2) *If ϕ is continuous, then the topology of A is normable.*

Following [Mi, p. 51], A is *functionally continuous* (FC) if every multiplicative functional on A is continuous. Note that $(C(X), \tau_d)$ in Example 2 is FC, but not Q . A major unsolved problem in topological algebras is the Michael problem: Is every multiplicative linear functional on a Frechet lmc algebra continuous? This has led to several sufficient conditions for A to be FC. This makes FC a reasonable assumption.

Let A be a commutative topological algebra. A is *weakly regular* if given a closed set $F \subset \sigma(A)$, $F \neq \sigma(A)$, there exists $x \neq 0$ in A such that $f(x) = 0$ for all $f \in F$. In the context of Banach algebras, weak regularity arises naturally in the study of uniqueness of the uniform norm [BD]; and it is weaker than regularity. This is exhibited in an example due to Barnes [Me, Example 1]. Let $D = \{z \in \mathbb{C} : |z| < 1\}$, $X = \bar{D} \times [0, 1]$. Let $A = \{f \in C(X) : f \text{ is holomorphic on } D \times \{0\}\}$, a uniform Banach algebra. Then A is weakly regular, but not regular. Since regularity in a uniform algebra is a stringent property, the validity of Theorem 1 under weak regularity is interesting.

Proof of Theorem 1. (1) Let $C = (\text{Im } \phi)^-$. Since A is FC, $\phi^* : \sigma(C) \rightarrow \sigma(A)$, $\phi^*(f) = f \circ \phi$ is well defined. It is also continuous with respective Gelfand topologies. Since the algebra C is a locally convex Q -algebra, $\sigma(C)$ is compact [Ma, p. 187]. Then $\phi^*(\sigma(C))$ is a compact subset of $\sigma(A)$. We have $\phi^*(\sigma(C)) = \sigma(A)$. Indeed, if $\phi^*(\sigma(C)) \neq \sigma(A)$, there exists $x \neq 0$ in A such that $\chi(\phi(x)) = 0$ for all χ in $\sigma(C)$.

Since C is commutative semisimple and lmc, $\phi(x) = 0$; and then $x = 0$ for ϕ is one-to-one. Thus $\phi^*(\sigma(C)) = \sigma(A)$. Now let $P = (p_\alpha)$ be a family of uniform seminorms defining the topology τ of A . Since A is advertibly complete, the spectrum $\text{Sp}_A(x) = \{\chi(x) : \chi \in \sigma(A)\}$ and the spectral radius $\rho_A(x) = \sup_\alpha \{\lim_{n \rightarrow \infty} (p_\alpha(x^n))^{1/n}\}$ for all x in A [Ma, p. 104, p. 99]. Since $\sigma(A)$ is compact, $\text{Sp}_A(x)$ is bounded. Since $p_\alpha(x^2) = p_\alpha(x)^2$ for all x and α , $\rho_A(x) = \sup_\alpha p_\alpha(x)$ for all $x \in A$. Also, $\sigma(A) = \phi^*(\sigma(C))$ gives $\text{Sp}_A(x) = \{f(\phi(x)) : f \in \sigma(C)\} \subset \text{Sp}_C(\phi(x))$. Further, as C is a Q -algebra, $s(C) = \{x \in C : \rho_C(x) \leq 1\}$ is a neighbourhood of 0 by [Mi, Prop. 13.5, p. 58]; and there exists a convex balanced open set W such that $0 \in W \subseteq s(C)$. The Minkowski functional q of W in C is a continuous seminorm satisfying $\rho_C(y) \leq q(y)$ for all $y \in C$. Hence for each α , for each $x \in A$, $p_\alpha(x) \leq \rho_A(x) \leq \rho_C(\phi(x)) \leq q(\phi(x))$. This proves that $\phi^{-1}/\text{Im } \phi$ is continuous.

(2) Suppose ϕ is continuous. Then $\phi^* : \sigma(C) \rightarrow \sigma(A)$ is well defined even if A is not FC. Then $\phi^{-1}/\text{Im } \phi$ is continuous as above making ϕ a topological isomorphism. Thus $\phi(A)$ is advertibly complete; and hence inverse closed in its completion. Whence it is inverse closed in the Q -algebra C , for the completion of $\phi(A)$ is contained in C . Therefore $\phi(A)$, and so A , is a Q -algebra. Hence the topology on A given by the algebra norm ρ_A is finer than τ . Now since A is a Q -algebra, $s(A) = \{x \in A : \rho_A(x) \leq 1\}$ is a neighbourhood of 0 on (A, τ) . Thus ρ determines τ . $\square$

Remark. Once ϕ^* is defined, the full strength of weak regularity has not been used. In fact, one has only to find a nonzero element vanishing on a given compact set.

We now give a result in the absence of FC. We consider the space $\sigma^*(A)$ consisting of all nonzero multiplicative functionals on A endowed with the weak topology $\sigma(A^*, A)$. We then introduce the following notion of weak σ^* -compact-regular weakened in the sense of the previous remark.

Definition. A commutative topological algebra A is called *weakly σ^* -compact-regular* if for a compact subset K of $\sigma^*(A)$, $K \neq \sigma^*(A)$, there exists a nonzero $x \in A$ such that $\chi(x) = 0$ for all $\chi \in K$.

Theorem 2. *Let A be a weakly σ^* -compact-regular advertibly complete uniform algebra, B a locally convex algebra and $\phi : A \rightarrow B$ a one-to-one homomorphism such that $C = (\text{Im } \phi)^-$ is a strongly semisimple Q -algebra. Then $\phi^{-1}/\text{Im } \phi$ is continuous. If ϕ is continuous, then the topology of A is normable.*

For the proof, consider the map $\phi^{**} : \sigma(C) \rightarrow \sigma^*(A)$, $\phi^{**}(\chi) = \chi \circ \phi$. If $\chi \circ \phi$ is identically zero, then by the continuity of χ , one obtains that χ is also identically zero. This contradicts $\chi \in \sigma(C)$. Thus ϕ^{**} is defined; and then it is continuous. Now one obtains $\phi^{**}(\sigma(C)) = \sigma^*(A)$; and the proof can be completed as in Theorem 1.

We conjecture that the semisimplicity of $(\text{Im } \phi)^-$ in Theorem 1 (and strong semisimplicity in Theorem 2) can be omitted. The following supports this.

Theorem 3. *Let A be a uniform lmca, B a locally convex algebra, and $\phi : A \rightarrow B$ a one-to-one homomorphism such that $(\text{Im } \phi)^-$ is a Q -algebra. Assume that at least one of the following holds.*

- (a) A is advertibly complete and $\text{Im } \phi$ is FC with continuous product.
- (b) A is FC, Ptak (as a l.c. space), regular, having locally equicontinuous spectrum $\sigma(A)$ (in particular, A is FC, Frechet, regular, having locally compact spectrum

$\sigma(A)$), and B is *lmca*. Then $\phi^{-1}/\text{Im } \phi$ is continuous. If ϕ is continuous, then the topology of A is normable.

Proof. (1) Assume (a). Then $\sigma^*((\text{Im } \phi)^-) = \sigma((\text{Im } \phi)^-)$ (since a Q -algebra is FC) $= \sigma(\text{Im } \phi)$ (by the joint continuity of multiplication in $\text{Im } \phi = \sigma^*(\text{Im } \phi)$ and $\phi^*(\sigma^*(\text{Im } \phi)) = \sigma^*(A)$ as ϕ is one-to-one. Then, for all $x \in A$, $\text{Sp}_A(x) = \{\chi(x) : \chi \in \sigma(A)\} = \{\chi(x) : \chi \in \sigma^*(A)\} = \{f(\phi(x)) : f \in \sigma^*(\text{Im } \phi)\} = \{f(\phi(x)) : f \in \sigma^*(C)\}$. Hence for some continuous seminorm q , $\rho_A(x) = \rho_C(x) \leq q(\phi(x))$ ($x \in A$).

(2) Assume (b). By [Ma, Coro. 1.3, p. 184], local equicontinuity of $\sigma(A)$ implies continuity of the Gelfand map $x \rightarrow \hat{x}$ and local compactness of $\sigma(A)$. We show that $\sigma(A) = \phi^*(\sigma(C))$. Note that $\phi^*(\sigma(C)) \subset \sigma(A)$. Suppose $\chi \in \sigma(A) \setminus \phi^*(\sigma(C))$. By the local compactness, there exists a compact set $K \subseteq \sigma(A)$ and disjoint open sets U, V in $\sigma(A)$ such that $\chi \in K \subset U$, $\phi^*(\sigma(C)) \subset V$. As A is Ptak, regular and having continuous Gelfand map, [Ma, Coro. 4.4, p. 344] implies that there exist $x, y \in A$ such that $g(x) = 1$ ($g \in \phi^*(\sigma(C))$), $g(x) = 0$ ($g \in \sigma(A) \setminus V$); $g(y) = 1$ ($g \in K$), $g(y) = 0$ ($g \in \sigma(A) \setminus U$). Then $g(x)g(y) = 0$ for all $g \in \sigma(A)$. By the semisimplicity of A , $xy = 0 = \phi(x)\phi(y)$. On the other hand, for all $f \in \sigma(C)$, $f(\phi(x)) = 1$. Thus $0 \notin \{f(\phi(x)) : f \in \sigma(C)\} = \text{Sp}_C(\phi(x))$, C being *lmc* and a Q -algebra. Thus $\phi(x)$ is invertible in C . Hence $\phi(y) = \phi(x)^{-1}\phi(x)\phi(y) = 0$, so that $y = 0$, a contradiction. It follows that $\phi^*(\sigma(C)) = \sigma(A)$. Now the proof can be completed as in Theorem 1. Note that if A is Frechet, then every compact subset of $\sigma(A)$ is equicontinuous [Mi, Prop. 4.2, p. 17]. Hence by [Ma, Th. 1.1, p. 182], the Gelfand map is continuous. Further if $\sigma(A)$ is locally compact, then it is locally equicontinuous [Ma, Cor. 1.3, p. 184]. $\square$

Remarks. (1) If B is *lmca* and $C = (\text{Im } \phi)^-$ is a semisimple Q -algebra, then C is strongly semisimple.

(2) Actually in the above theorems, $\phi^{-1} : \phi(A) \rightarrow (A, \|\cdot\|)$ is continuous, where $\|\cdot\|$ is the uniform norm given by $\|x\| = \sup\{p(x) : p \text{ is a continuous uniform seminorm}\}$. The existence of this norm implies that A is spectrally bounded.

(3) Theorem 2 also applies to Example 1. Indeed, the algebra $(C[0, \Omega], \tau)$ is a complete uniform algebra. It is weakly σ^* -compact-regular, for $\sigma^*(C[0, \Omega])$ is homeomorphic to the Stone-Cech compactification $\beta[0, \Omega]$ of $[0, \Omega]$ and $C[0, \Omega]$ is isomorphic to the algebra $C(\beta[0, \Omega])$ [GJ].

(4) The hypothesis $(\text{Im } \phi)^-$ is a Q -algebra cannot be omitted. Let $A = C_b(\mathbb{R})$ be the algebra of all continuous bounded functions on the real line. Endowed with the sup norm, it is a uniform Banach algebra. By the same arguments as in (3), one shows that A is weakly σ^* -compact-regular. Consider $B = C(\mathbb{R})$ to be the algebra of all continuous functions with the compact open topology. Consider $\phi : A \rightarrow B$, $\phi(f) = f$. Then $(\text{Im } \phi)^- = C(\mathbb{R})$. It is well known that it is not a Q -algebra. Clearly ϕ^{-1} is discontinuous.

(5) The referee has asked: (In above theorems) does the automatic continuity of ϕ^{-1} on $\text{Im } \phi$ necessitate $(\text{Im } \phi)^-$ a Q -algebra? The following answers this.

Proposition 4. *Let A be a complete Q -lmca, B an *lmca*, and $\phi : A \rightarrow B$ a one-to-one homomorphism such that $\phi^{-1}/\text{Im } \phi$ is continuous. Then $(\text{Im } \phi)^-$ is a Q -algebra.*

Proof. We may assume that B is complete, hence $C = (\text{Im } \phi)^-$ is complete. Then $\phi^{-1}/\text{Im } \phi$ extends as a continuous homomorphism $\psi : C \rightarrow A$. By assumption, there exists a continuous seminorm p on A such that for all x in A , $r_C(\phi(x)) \leq r_{\text{Im } \phi}(\phi(x)) \leq r_A(x) \leq p(x) \leq p(\psi(\phi(x)))$. Since ψ is continuous, there exists a

continuous seminorm q on C such that $p(\psi(y)) \leq q(y)$ ($y \in C$). Hence in the above, $r_C(\phi(x)) \leq p(\psi(\phi(x))) \leq q(\phi(x))$ for all x in A . Now let $y \in C$, $y = \lim \phi(x_\alpha)$ for some net (x_α) in A . For any continuous multiplicative functional f on C , $|f(y)| \leq |f(y - \phi(x_\alpha))| + |f(\phi(x_\alpha))| \leq |f(y - \phi(x_\alpha))| + q(\phi(x_\alpha)) \rightarrow q(y)$. Hence $r_C(y) = \sup |f(y)| \leq q(y)$ ($y \in C$) showing that C is a Q -algebra.

If A is not a Q -algebra, then this does not hold. For the open unit disc U in the complex plane, let $A = H(U)$ be the uniform Frechet algebra consisting of holomorphic functions on U with the compact-open topology, $B = C(U)$ with the compact-open topology, and $\phi : A \rightarrow B$ be $\phi(f) = f$. Clearly ϕ is a homeomorphism and $(\text{Im } \phi)^- = B$ fails to be a Q -algebra. $\square$

APPLICATIONS

(1) Proposition. *Let A be an advertibly complete lmca. Let $\|\cdot\|$ be any continuous norm on A . Then A cannot be simultaneously weakly regular and uniform unless the topology of A is normable.*

Proof. Let τ denote the lmc topology on A . Let $P = (p_\alpha)$ be a family of submultiplicative seminorms on A defining τ . Then $P_0 = P \cup \{\|\cdot\|\}$ also determines τ . Suppose (A, τ) is uniform. Then τ is defined by a family $S = (q_i)$ of uniform seminorms. By closing S with maxima of finite subfamilies and applying the continuity of $\|\cdot\|$, there exists a q in S which is a norm. Let A_q be the uniform Banach algebra obtained by completing (A, q) . Let $\phi : A \rightarrow A_q$ be $\phi(x) = x$. Now if A is weakly regular, then Theorem 1 applied to ϕ implies that τ is normable.

It follows that an advertibly complete non-normed weakly regular uniform algebra cannot support a continuous norm. Let X be a compact Hausdorff space. By a well known result of Kaplansky, if $|\cdot|$ is any norm on $C(X)$ making it a normed algebra, then the supnorm $\|\cdot\| \leq |\cdot|$. The following has a bearing with this. A norm on an algebra A is *semisimple* if the completion of $(A, |\cdot|)$ is semisimple [BD]. $\square$

(2) Corollary. *Let $\|\cdot\|$ be a uniform norm on an algebra A such that $(A, \|\cdot\|)$ is a weakly regular Q -normed algebra. Let $|\cdot|$ be any submultiplicative norm on A .*

- (i) *If $|\cdot|$ is semisimple, then $\|\cdot\| \leq |\cdot|$. Further if $|\cdot|$ is continuous, then $|\cdot|$ is equivalent to $\|\cdot\|$.*
- (ii) *Let $(A, \|\cdot\|)$ be complete and regular. Then $\|\cdot\| \leq |\cdot|$ for any submultiplicative norm $|\cdot|$.*

Indeed let $\phi : (A, \|\cdot\|) \rightarrow (\tilde{A}, |\cdot|)$ ($\tilde{A}$ = completion of $(A, |\cdot|)$), $\phi(x) = x$. Theorem 1 implies that there exists $k > 0$ such that $\|\cdot\| \leq k|\cdot|$. Since $(A, \|\cdot\|)$ is Q , $\rho_A(x) = \inf \|x^n\|^{1/n} = \lim \|x^n\|^{1/n} = \|x\| \leq \lim |x^n|^{1/n} \leq |x|$ for all $x \in A$. (ii) follows by Theorem 3(b).

(3) Let $A = \mathbb{C} \times C_c^\infty(\mathbb{R})$ (resp. $B = \mathbb{C} \times C_c(\mathbb{R})$) be the algebra of all complex C^∞ -functions (resp. continuous functions) on $\mathbb{R}$ which are constant outside some compact set (depending on the function). We endow A (resp. B) with the inductive limit topology τ_D (resp. τ_K). The algebra (A, τ_D) is a complete regular lmca, (B, τ_K) is a lmc Q -algebra [Ma, p. 128] and A is dense in B [K, p. 148]. Consider $\phi : A \rightarrow B$, $\phi(f) = f$. Since the topology τ_D is finer than τ_K on A , ϕ is continuous. It is classical that A is not normable. Hence by Theorem 1, (A, τ_D) cannot be uniform.

Let A be the algebra $\mathbb{C} \times C_c^\infty(\mathbb{R})$ endowed with the compact open topology τ . It is a weakly σ^* -compact-regular uniform lmca. Since it is inverse closed in $C(\mathbb{R})$, it

is advertibly complete. Take (B, τ_K) as above. Let $\phi : (A, \tau) \rightarrow (B, \tau_K)$, $\phi(f) = f$. By Theorem 1, ϕ^{-1} is continuous.

(4) Let $U \subset \mathbb{C}^d$ be open. Let $H(U)$ be the uniform Frechet algebra of all holomorphic functions on U with the compact open topology. Let $H^\infty(U) = \{f \in H(U) : f \text{ is bounded}\}$, a uniform Banach algebra. Let $X \subset \mathbb{C}^d$ be compact. Let $H(X)$ be the algebra of holomorphic germs on X . Choose a decreasing sequence (U_n) of open neighbourhoods of X such that $\bar{U}_{n+1} \subset U_n$ and $\bar{U}_{n+1}$ is compact. In view of the continuous embeddings

$$\cdots \rightarrow H^\infty(U_n) \rightarrow H(U_n) \rightarrow H^\infty(U_{n+1}) \rightarrow H(U_{n+1}) \rightarrow \cdots,$$

$H(X)$ can be realized as inductive limits $H(X) = \varinjlim H(U_n) = \varinjlim H^\infty(U_n)$, its topology τ being the finest locally convex topology making all $\phi_n : H(U_n) \rightarrow H(X)$, $\phi_n(f) = f/X$ and similarly making all $\psi_n : H^\infty(U_n) \rightarrow H(X)$, $\psi_n(f) = f/X$ continuous.

None of $H^\infty(U)$ and $H(U)$ is weakly regular. Note that $(H(X), \tau)$ is a complete semisimple Q -algebra [Ma, p. 134]. If $H(U_n)$ is weakly regular, then by Theorem 1, it becomes a Banach algebra and ϕ_n becomes a homeomorphism. If $H^\infty(U_n)$ is weakly regular, then ψ_n becomes a homeomorphism. Either of these forces $H(X)$ to be a uniform Banach algebra. Being a complete, non-normed Q -algebra, $H(X)$ is not a uniform algebra [BD].

(5) Let $D = \{z \in \mathbb{C} : |z| < 1\}$, $Y = D \times [0, 1]$, $Z = \bar{D} \times [0, 1]$. Let $A = \{f \in C(Y) : f \text{ is holomorphic on } D \times \{0\}\}$, $B = \{f \in C(Z) : f \text{ is holomorphic on } D \times \{0\}\}$. Let $0 < r < 1$. Let $\|f\|_r = \sup\{|f(x)| : x \in \bar{D}_r \times [0, 1]\}$ ($f \in A$); $|f|_r = \sup\{|f(x)| : x \in \bar{D} \times [0, r]\}$ ($f \in B$). Each of A and B with the topology defined respectively by $\{\|\cdot\|_r : 0 < r < 1\}$ and $\{|\cdot|_r : 0 < r < 1\}$ is a uniform Frechet algebra. Any $f \in C(Y)$ (resp. $f \in C(Z)$) vanishing on $D \times \{0\}$ is in A (resp. in B). This implies that both A and B are weakly regular, not regular. Thus each of A and B fails to support a continuous norm.

ACKNOWLEDGEMENT

A query by the referee resulted in Proposition 4. The referee has also made several suggestions regarding the exposition. We sincerely thank the referee for these.

REFERENCES

- [BK] S. J. Bhatt and D. J. Karia, *Uniqueness of the uniform norm with an application to topological algebras*, Proc. Amer. Math. Soc. **116** (2) (1992), 499–503. MR **92m**:46068
- [B] S. J. Bhatt, *Automatic continuity of homomorphisms in topological algebras*, Proc. Amer. Math. Soc. **119** (1) (1993), 135–139. MR **93k**:46042
- [BD] S. J. Bhatt and H. V. Dedania, *Banach algebras with unique uniform norm*, Proc. Amer. Math. Soc. **124** (2) (1996), 579–584. MR **96d**:46067
- [GJ] L. Gillman and M. Jerison, *Rings of continuous functions*, Springer-Verlag, New York, Heidelberg, Berlin, 1976. MR **53**:11352
- [K] Vo-Khac Khoan, *Distributions Analyse de Fourier Operateurs aux dérivées partielles*, vol. T. 1, Vuibert, Paris, 1972.
- [L] R. J. Loy, *Math. Reviews*, 93k:46042, p. 6116, Amer. Math. Soc., 1993.
- [Ma] A. Mallios, *Topological Algebras*, Selected Topics, North-Holland, 1986. MR **87m**:46099
- [Me] M. J. Meyer, *Spectral extension property and extension of multiplicative linear functionals*, Proc. Amer. Math. Soc. **112** (1991), 855–861. MR **91j**:46059
- [Mi] E. Michael, *Locally multiplicatively-convex algebras*, Mem. Amer. Math. Soc., vol. 11, Providence, RI, 1952. MR **14**:482a

- [S] J. Schmets, *Espaces de fonctions continues*, Lecture Notes in Mathematics 519, Springer-Verlag, 1976. MR **54**:11041
- [Z] W. Zelazko, *Banach algebras*, Elsevier Publishing Co. Amsterdam, London, New York, 1973. MR **56**:6389

(A. Beddaa and M. Oudadess) ECOLE NORMALE SUPERIEURE, B.P. 5118 TAKADDOUN, RABAT, MOROC

(S. J. Bhatt) DEPARTMENT OF MATHEMATICS, SARDAR PATEL UNIVERSITY, VALLABH VIDYANAGAR 388120, GUJARAT, INDIA