

## A CHARACTERIZATION OF MÖBIUS TRANSFORMATIONS

ROLAND HÖFER

(Communicated by Christopher Croke)

**ABSTRACT.** Let  $n \geq 2$  be an integer and let  $\mathcal{D}$  be a domain of  $\mathbb{R}^n$ . Let  $f : \mathcal{D} \rightarrow \mathbb{R}^n$  be an injective mapping which takes hyperspheres whose interior is contained in  $\mathcal{D}$  to hyperspheres in  $\mathbb{R}^n$ . Then  $f$  is the restriction of a Möbius transformation.

### 1. INTRODUCTION

Let  $n \geq 2$  be an integer. A theorem of A.D. Alexandrov [1] states that any bijective transformation of  $\mathbb{R}^{n+1}$  which preserves the Lorentz distance 0 between pairs of points in both directions is the product of a Lorentz transformation and a dilatation. The following Theorem 1.3 is due to A.D. Alexandrov [2], J.A. Lester [7], and I. Popovici and D.C. Rădulescu [9] and generalizes Alexandrov's theorem.

**Definition 1.1.** Let  $n \in \mathbb{N}$ ,  $n \geq 2$ . For  $x, y \in \mathbb{R}^n$  let  $x \cdot y$  denote the standard euclidean product between  $x$  and  $y$ . The Lorentz product, resp. Lorentz distance, between  $x, y \in \mathbb{R}^{n+1}$  is defined by

$$\begin{aligned} x \diamond y &:= x_1 y_1 + \dots + x_n y_n - x_{n+1} y_{n+1}, \\ d(x, y) &:= (y - x) \diamond (y - x). \end{aligned}$$

**Definition 1.2** (cf. [6]). Let  $n \in \mathbb{N}$ ,  $n \geq 2$ .

a) Let  $\mathcal{D} \subset \mathbb{R}^n$ . A mapping  $f : \mathcal{D} \rightarrow \mathbb{R}^n$  is the restriction of a *Möbius transformation* if  $\mathbb{R}\sigma_1(f(x)) = \mathbb{R}(\sigma_1(x)A_1)$  is satisfied for all  $x \in \mathcal{D}$ , where

$$\sigma_1(z) := \left( \frac{1 - z \cdot z}{2}, z, \frac{1 + z \cdot z}{2} \right)$$

for all  $z \in \mathbb{R}^n$ , and where  $A_1$  is an  $(n+2) \times (n+2)$ -Lorentz matrix,  $A_1 M_1 A_1^T = M_1 := \text{diag}(1, \dots, 1, -1)$ .

b) Let  $\mathcal{D} \subset \mathbb{R}^{n+1}$ . A mapping  $f : \mathcal{D} \rightarrow \mathbb{R}^{n+1}$  is the restriction of a *Lie transformation* if  $\mathbb{R}\sigma_2(f(x)) = \mathbb{R}(\sigma_2(x)A_2)$  for all  $x \in \mathcal{D}$ , where

$$\sigma_2(z) := \left( \frac{1 - z \diamond z}{2}, z, \frac{1 + z \diamond z}{2} \right)$$

for all  $z \in \mathbb{R}^{n+1}$ , and where  $A_2$  is an  $(n+3) \times (n+3)$ -matrix with  $A_2 M_2 A_2^T = M_2 := \text{diag}(1, \dots, 1, -1, -1)$ .

---

Received by the editors June 4, 1998.

1991 *Mathematics Subject Classification.* Primary 51B10; Secondary 51M04, 51M09.

*Key words and phrases.* Möbius transformation, Lie transformation, mappings preserving hyperspheres, Alexandrov's theorem for domains.

**Theorem 1.3.** *Let  $\mathcal{D}$  be a domain (i.e. an open, connected subset) of  $\mathbb{R}^{n+1}$ ,  $n \geq 2$ . Let  $f : \mathcal{D} \rightarrow \mathbb{R}^{n+1}$  be a mapping such that*

$$d(x, y) = 0 \quad \Leftrightarrow \quad d(f(x), f(y)) = 0$$

*for all  $x, y \in \mathcal{D}$ . Then  $f$  is the restriction of a Lie transformation.*

Alexandrov's theorem and Theorem 1.3 are important results in a modern field of geometrical research which is called *characterizations of geometrical mappings under mild hypotheses* [3], [4], [8]. In particular no regularity assumptions such as differentiability or even continuity are needed in these kinds of characterizations. In the same sense, C. Carathéodory proved [5] that any injective mapping of a domain  $\mathcal{D}$  of  $\mathbb{R}^2$  to  $\mathbb{R}^2$  is the restriction of a Möbius transformation if the following condition is satisfied:

*The image of any circle contained with its interior in  $\mathcal{D}$ , is itself a circle.*

## 2. RESULTS

There is a close connection between Carathéodory's theorem and Theorem 1.3 ( $n = 2$ ). In fact we will generalize Carathéodory's theorem to arbitrary dimensions with the help of Theorem 1.3.

**Theorem 2.1.** *Let  $n \geq 2$  be an integer and let  $\mathcal{D}$  be a domain of  $\mathbb{R}^n$ . Let  $f : \mathcal{D} \rightarrow \mathbb{R}^n$  be an injective mapping such that  $f(H)$  is a hypersphere, whenever  $H \subset \mathcal{D}$  is a hypersphere and the interior of  $H$  is contained in  $\mathcal{D}$ . Then  $f$  is the restriction of a Möbius transformation.*

**Definition 2.2.** A *similarity* of  $\mathbb{R}^n$ ,  $n \geq 2$ , is a mapping  $f : \mathbb{R}^n \rightarrow \mathbb{R}^n$ ,  $f(x) = kxQ + t$  where  $k > 0$ ,  $t \in \mathbb{R}^n$ , and  $Q$  is an orthogonal  $n \times n$ -matrix,  $QQ^T = E$ .

It is well known that a mapping  $f : \mathbb{R}^n \rightarrow \mathbb{R}^n$  which is induced by a Möbius transformation is a similarity. Hence, Theorem 2.1 implies the following corollary.

**Corollary 2.3.** *Let  $n \geq 2$ . Let  $f : \mathbb{R}^n \rightarrow \mathbb{R}^n$  be an injective mapping such that images of euclidean hyperspheres are euclidean hyperspheres. Then  $f$  is a similarity.*

Now let  $\mathcal{D}$  be the set  $I^n := \{x \in \mathbb{R}^n \mid x \cdot x < 1\}$  of points in Poincaré's sphere model of  $n$ -dimensional hyperbolic geometry,  $n \geq 2$ . A hyperbolic hypersphere in  $I^n$  is a euclidean hypersphere which is contained in  $I^n$ . If  $f : I^n \rightarrow I^n$  is induced by a Möbius transformation and if  $f$  is surjective, then  $f$  is a hyperbolic motion.

**Corollary 2.4.** *Let  $n \geq 2$ . Let  $f : I^n \rightarrow I^n$  be a bijection of  $n$ -dimensional hyperbolic space which maps hyperbolic hyperspheres onto hyperbolic hyperspheres. Then  $f$  is a hyperbolic motion.*

## 3. PROOF OF THEOREM 2.1

We show that, whenever  $H$  is a hypersphere contained in  $\mathcal{D}$  such that the interior  $I$  of  $H$  is also contained in  $\mathcal{D}$ , then  $f|_I$  is the restriction of a Möbius transformation. This implies Theorem 2.1 since

a) Any Möbius transformation is uniquely determined by its restriction to any non-empty open subset of  $\mathbb{R}^n$ .

b) For any two points  $x, y \in \mathcal{D}$ , there is a finite sequence  $I_1, \dots, I_k \subset \mathcal{D}$  of interiors of hyperspheres with  $x \in I_1$ ,  $y \in I_k$ ,  $I_j \cap I_{j+1} \neq \emptyset$  for all  $j \in \{1, \dots, k-1\}$ .

Let  $H$  be a hypersphere contained in  $\mathcal{D}$  such that the interior  $I$  of  $H$  is also contained in  $\mathcal{D}$ .

1. Let  $I'$  denote the interior of the hypersphere  $H' := f(H)$ . Then either  $f(I) \subset I'$  or  $f(I) \subset \mathbb{R}^n \setminus (H' \cup I')$ .

*Proof.* Let  $x, y \in I$ . Then there is a hypersphere  $H_1 \subset I$  which contains  $x$  and  $y$ . Since  $f$  is injective and  $f(H_1)$  is a hypersphere, either  $f(H_1) \subset I'$  or  $f(H_1) \subset \mathbb{R}^n \setminus (H' \cup I')$ . Thus  $f(x), f(y)$  are on the same side of  $f(H_1)$ .

2. Let  $\mu : \mathbb{R}^n \setminus I' \rightarrow \mathbb{R}^n$  denote the restriction of a Möbius transformation which satisfies  $\mu(H') = H'$  and  $\mu(\mathbb{R}^n \setminus (I' \cup H')) \subset I'$ . Let  $g : H \cup I \rightarrow \mathbb{R}^n$  be defined by  $g := f|_{H \cup I}$  if  $f(I) \subset I'$ , and  $g := \mu \circ f|_{H \cup I}$  if  $f(I) \subset \mathbb{R}^n \setminus (H' \cup I')$ . Then  $g(I) \subset I'$ ,  $g(H) = H'$ , and  $g$  is an injective mapping which takes hyperspheres in  $H \cup I$  to hyperspheres in  $H' \cup I'$ .

3. Let  $H_1 \subset I$  be a hypersphere with interior  $I_1$ . Then  $g(I_1)$  is contained in the interior  $I'_1$  of  $H'_1 := g(H_1)$ , and  $g(I \setminus (H_1 \cup I_1))$  is contained in the exterior of  $H'_1$ .

*Proof.* Let  $z \in I \setminus (H_1 \cup I_1)$ . There is a hypersphere  $H_2 \subset H \cup I$  with  $z \in H_2$ ,  $\#(H \cap H_2) = 1$  and  $H_1 \cap H_2 = \emptyset$ . Then  $H'_2 := g(H_2) \subset H' \cup I'$ ,  $g(z) \in H'_2$ ,  $\#(H' \cap H'_2) = 1$  and  $H'_1 \cap H'_2 = \emptyset$ . Hence  $g(z) \notin I'_1$ , and  $g(I \setminus (H_1 \cup I_1)) \subset I' \setminus (H'_1 \cup I'_1)$ . From the proof of 1. we know that  $g(I_1)$  is either contained in the interior or in the exterior of  $H'_1$ . We take a hypersphere  $H_3 \subset I$ ,  $\#(H_1 \cap H_3) > 1$ . Then  $H_3 \cap I_1 \neq \emptyset$  and  $\#(H'_1 \cap g(H_3)) > 1$ . Hence  $g(H_3 \cap I_1) \cap I'_1 \neq \emptyset$  and  $g(I_1) \subset I'_1$ .

**Definition 3.1.** Two hyperspheres  $H_1, H_2 \subset \mathbb{R}^n$  are in *interior (exterior) contact* if  $\#(H_1 \cap H_2) = 1$  and  $H_i$  is contained in the interior (exterior) of  $H_j$  where  $(i, j) = (1, 2)$  or  $(i, j) = (2, 1)$ .

4. Two hyperspheres  $H_1, H_2 \subset I$  are in interior (exterior) contact iff  $g(H_1), g(H_2)$  are in interior (exterior) contact.

*Proof.* Since  $g$  is injective,  $\#(H_1 \cap H_2) = 1$  iff  $\#(g(H_1) \cap g(H_2)) = 1$ . The assertion now follows from 3.

**Definition 3.2.** For any hypersphere  $H_1$  let  $\gamma(H_1) \in \mathbb{R}^n$ ,  $\rho(H_1) > 0$  denote the euclidean center and radius of  $H_1$ . Let  $\lambda(H_1) := (\gamma(H_1), \rho(H_1)) \in \mathbb{R}^n \times \mathbb{R}_{>0}$ .

5. Two distinct hyperspheres  $H_1, H_2$  of  $\mathbb{R}^n$  are in interior contact iff the Lorentz distance between  $\lambda(H_1)$  and  $\lambda(H_2)$  is zero.

6. The set  $\mathcal{C} := \{\lambda(H_1) \mid H_1 \subset I \text{ is a hypersphere}\}$  is a domain of  $\mathbb{R}^{n+1}$ .

*Proof.*  $\mathcal{C} = \{x \in \mathbb{R}^n \times ]0, \rho(H)[ \mid d(x, \lambda(H)) < 0\}$  is open and connected.

7. The mapping  $\varphi := \lambda \circ g \circ \lambda^{-1} : \mathcal{C} \rightarrow \mathcal{C}' := \{\lambda(g(H_1)) \mid H_1 \subset I \text{ is a hypersphere}\}$  satisfies  $d(x, y) = 0$  iff  $d(\varphi(x), \varphi(y)) = 0$  for all  $x, y \in \mathcal{C}$ .

*Proof.* From 5. and 4., for all distinct hyperspheres  $H_1, H_2 \subset I$ ,

$$\begin{aligned} d(\lambda(H_1), \lambda(H_2)) = 0 &\Leftrightarrow H_1 \text{ and } H_2 \text{ are in interior contact} \\ &\Leftrightarrow g(H_1) \text{ and } g(H_2) \text{ are in interior contact} \\ &\Leftrightarrow d(\lambda(g(H_1)), \lambda(g(H_2))) = 0. \end{aligned}$$

8. From 7. and Theorem 1.3,  $\varphi$  is the restriction of a Lie transformation, i.e. there is an  $(n+3) \times (n+3)$ -matrix  $A_2 =: (a_{ij})_{i,j=1,\dots,n+3}$  as in Definition 1.2 b), such that  $\mathbb{R}\sigma_2(y) = \mathbb{R}(\sigma_2(x)A_2)$  for all  $x \in \mathcal{C}$ ,  $y = \varphi(x)$ .

**Definition 3.3.** A *light line* of  $\mathbb{R}^{n+1}$  is a line  $u + \mathbb{R}v$ ,  $u, v \in \mathbb{R}^{n+1}$ ,  $v \neq 0$ , where  $d(v, v) = 0$ .

9.  $f|_I$  is the restriction of a Möbius transformation.

*Proof.* Let  $x \in I$ . Let  $l_1, l_2$  be two distinct light lines which contain  $(x, 0)$ . Then  $\{(x, 0)\} = l_1 \cap l_2 \subset \partial\mathcal{C}$ . The images  $\varphi(l_i \cap \mathcal{C}) \neq \emptyset$  are contained in uniquely determined light lines  $l'_i$ ,  $i = 1, 2$ . Since  $\varphi$  is continuous,  $\{(g(x), 0)\} = l'_1 \cap l'_2$  is contained in  $\partial\mathcal{C}'$ . Hence for all  $x \in I$ , we have  $\mathbb{R}\sigma_2((g(x), 0)) = \mathbb{R}(\sigma_2((x, 0))A_2)$  which implies

$$(3.1) \quad \mathbb{R}\sigma_1(g(x)) = \mathbb{R}(\sigma_1(x)A_1),$$

$$(3.2) \quad \sigma_1(x) \cdot (a_{1,n+2}, \dots, a_{n+1,n+2}, a_{n+3,n+2}) = 0$$

where  $A_1$  is the  $(n+2) \times (n+2)$ -matrix obtained from  $A_2$  by deleting the  $(n+2)$ th row and  $(n+2)$ th column. Equation (3.2) is a quadratic equation in  $x = (x_1, \dots, x_n)$  which holds for any  $x \in I$ , and we obtain  $a_{1,n+2} = \dots = a_{n+1,n+2} = a_{n+3,n+2} = 0$ . Together with  $A_2 M_2 A_2^T = M_2$  we have  $A_1 M_1 A_1^T = M_1$ , where  $M_1$  is chosen as in Definition 1.2 a). Equation (3.1) implies that  $g$  is the restriction of a Möbius transformation. Hence also  $f|_I$  is the restriction of a Möbius transformation.  $\square$

*Remark 3.4.* It is possible to prove Theorem 2.1 by Carathéodory's theorem. If  $n = 3$  and  $f : \mathcal{D} \rightarrow \mathbb{R}^3$  is injective and has the sphere preserving property, then we can apply Carathéodory's theorem to any hypersphere  $H \subset \mathcal{D}$  whose interior is contained in  $\mathcal{D}$ , after removing a point  $p \in H$  and  $f(p) \in f(H)$ , to show that  $f$  is a Möbius transformation between  $H$  and its image  $f(H)$ . This Möbius transformation is the restriction of the same Möbius transformation for all hyperspheres. Induction proves the result for all  $n \geq 2$ .

I would like to thank the referee for helpful comments, especially Remark 3.4.

## REFERENCES

1. A. D. Alexandrov. Seminar report. *Uspekhi Mat. Nauk*, 37(3):187, 1950.
2. A. D. Alexandrov. On the axioms of relativity theory. *Vestnik Leningrad Univ. Math.*, 19:5–28, 1976.
3. W. Benz. Characterizations of geometrical mappings under mild hypotheses: Über ein modernes Forschungsgebiet der Geometrie. *Hamb. Beitr. Wiss.gesch.*, 15:393–409, 1994.
4. W. Benz. *Real Geometries*. BI Wissenschaftsverlag, Mannheim, Leipzig, Wien, Zürich, 1994. MR **95k**:51019
5. C. Carathéodory. The most general transformations of plane regions which transform circles into circles. *Bull. Am. Math. Soc.*, 43:573–579, 1937.
6. T. E. Cecil. *Lie Sphere Geometry*. Springer-Verlag, New York Berlin Heidelberg, 1992. MR **94m**:53076
7. J. A. Lester. A physical characterization of conformal transformations of Minkowski spacetime. *Ann. Discrete Math.*, 18:567–574, 1983. MR **84g**:83004b

8. J. A. Lester. Distance preserving transformations. In F. Buekenhout, editor, *Handbook of Incidence Geometry*, pages 921–944, Amsterdam, 1995. Elsevier. MR **96j**:51019
9. I. Popovici and D. C. Rădulescu. Characterizing the conformality in a Minkowski space. *Ann. Inst. H. Poincaré. Phys. Théor.*, 35:131–148, 1981. MR **83b**:53011

MATHEMATISCHES SEMINAR, UNIVERSITÄT HAMBURG, BUNDESSTR. 55, 20146 HAMBURG, GERMANY

*E-mail address:* `hoefer@math.uni-hamburg.de`