NONVANISHING OF SYMMETRIC SQUARE L -FUNCTIONS OF CUSP FORMS INSIDE THE CRITICAL STRIP

WINFRIED KOHNEN AND JYOTI SENGUPTA

(Communicated by Dennis A. Hejhal)

ABSTRACT. We shall give a certain nonvanishing result for the symmetric square L -function of an elliptic cuspidal Hecke eigenform w.r.t. the full modular group inside the critical strip.

1. INTRODUCTION

Let f be a normalized cuspidal Hecke eigenform of integral weight k on the full modular group $SL_2(\mathbf{Z})$ and denote by $D_f^*(s)$ ($s \in \mathbf{C}$) the symmetric square L -function of f completed with its archimedean Γ -factors. As is well-known [7], [8], $D_f^*(s)$ has a holomorphic continuation to $\mathbf{C}$ and is invariant under $s \mapsto 2k - 1 - s$. Note that, by [3], $D_f^*(s)$ (up to a variable shift) also is the standard zeta function of a cuspidal automorphic representation of $GL(3)$, and so by [4] zeros of $D_f^*(s)$ can occur only inside the critical strip $k - 1 < \operatorname{Re}(s) < k$. According to the generalized Riemann hypothesis, the zeros of $D_f^*(s)$ should all lie on the critical line $\operatorname{Re}(s) = k - \frac{1}{2}$.

The last statement of course is far from being settled. On the other hand, it turns out to be comparatively easy to prove nonvanishing results for $D_f^*(s)$ on the average. For example, in [6] Xian-Jin Li used an approximate functional equation for an average sum of the $D_f^*(s)$ to show that for any given s with $k - 1 < \operatorname{Re}(s) < k$, $s \neq k - \frac{1}{2}$, $\zeta(s - k + 1) \neq 0$, there are infinitely many different f such that $D_f^*(s)$ is not zero.

In the present note, using a different approach we will prove that given any s with $k - 1 < \operatorname{Re}(s) < k$, $\operatorname{Re}(s) \neq k - \frac{1}{2}$, then for all k large enough there exists a Hecke eigenform f of weight k such that $D_f^*(s) \neq 0$. For the proof we use a “kernel function” for $D_f^*(s)$ as given by Zagier in [8] and then proceed in a similar way as in [5], where a corresponding result for Hecke L -functions was proved.

2. NOTATION

For $s \in \mathbf{C}$ we usually write $s = \sigma + it$ with $\sigma, t \in \mathbf{R}$.

Received by the editors July 31, 1998.
 2000 *Mathematics Subject Classification*. Primary 11Fxx.

3. STATEMENT OF RESULT

Let k be an even integer ≥ 12 and let S_k be the space of cusp forms of weight k w.r.t. the full modular group $\Gamma_1 = SL_2(\mathbf{Z})$, equipped with the usual Petersson scalar product $\langle \cdot, \cdot \rangle$. For $f(z) = \sum_{n \geq 1} a(n)e^{2\pi i n z}$ ($z \in \mathcal{H} =$ upper half plane) a normalized Hecke eigenform in S_k (recall that normalized means $a(1) = 1$), we denote by

$$D_f(s) = \prod_p (1 - \alpha_p^2 p^{-s})^{-1} (1 - \alpha_p \beta_p p^{-s})^{-1} (1 - \beta_p^2 p^{-s})^{-1} \quad (\sigma > k)$$

the symmetric square L -function of f , where the product is taken over all rational primes p and α_p, β_p are defined by

$$\alpha_p + \beta_p = a(p), \quad \alpha_p \beta_p = p^{k-1}.$$

By [7], [8], $D_f(s)$ has a holomorphic continuation to $\mathbf{C}$, and the function

$$D_f^*(s) = 2^{-s} \pi^{-3s/2} \Gamma(s) \Gamma\left(\frac{s-k+2}{2}\right) D_f(s)$$

satisfies the functional equation

$$(1) \quad D_f^*(2k-1-s) = D_f^*(s).$$

Let $\{f_{k,1}, \dots, f_{k,g_k}\}$ ($g_k = \dim S_k$) be the basis of normalized Hecke eigenforms of S_k .

Theorem. *Let $t_0 \in \mathbf{R}$ and $0 < \epsilon < \frac{1}{2}$. Then there exists a positive constant $C(t_0, \epsilon)$ depending only on t_0 and ϵ such that for $k > C(t_0, \epsilon)$ the function*

$$\sum_{\nu=1}^{g_k} \frac{1}{\langle f_{k,\nu}, f_{k,\nu} \rangle} D_{f_{k,\nu}}^*(s)$$

does not vanish at any point $s = \sigma + it_0$, $k-1 < \sigma < k - \frac{1}{2} - \epsilon$, $k - \frac{1}{2} + \epsilon < \sigma < k$.

Corollary. *Let $s \in \mathbf{C}$ be fixed with $k-1 < \sigma < k$, $\sigma \neq k - \frac{1}{2}$. Then for all k large enough there exists a normalized Hecke eigenform f in S_k such that $D_f^*(s) \neq 0$.*

4. PROOF

The proof proceeds along similar lines as in [5]. We consider the cusp forms dual w.r.t. the Petersson scalar product to the values $D_f^*(s)$ ($2-k < \sigma < k-1$) where f is any normalized Hecke eigenform in S_k . These have been constructed by Zagier [8] and will be denoted Φ_s as in [8] in what follows. To state the relevant properties of Φ_s , we need to introduce several notations.

Let Δ be a discriminant, i.e. $\Delta \in \mathbf{Z}$ and $\Delta \equiv 0, 1 \pmod{4}$. Put

$$L(s, \Delta) = \begin{cases} \zeta(2s-1), & \text{if } \Delta = 0, \\ L_D(s) \sum_{d|f, d>0} \mu(d) \left(\frac{D}{d}\right) d^{-s} \sigma_{1-2s}\left(\frac{f}{d}\right), & \text{if } \Delta \neq 0, \end{cases}$$

where if $\Delta \neq 0$ we have written $\Delta = Df^2$ with $f \in \mathbf{N}$ and D the discriminant of $\mathbf{Q}(\sqrt{\Delta})$, $\left(\frac{D}{\cdot}\right)$ is the Kronecker symbol, $L_D(s)$ the associated L -function defined by analytic continuation of the series $\sum_{n \geq 1} \left(\frac{D}{n}\right) n^{-s}$ ($\sigma > 1$), μ is the Möbius function and $\sigma_\nu(m) = \sum_{d|m, d>0} d^\nu$ ($m \in \mathbf{N}$, $\nu \in \mathbf{C}$).

Furthermore, for t an integer with $\Delta < t^2$ and $s \in \mathbf{C}$ with $\frac{1}{2} < \sigma < k$ we define

$$\begin{aligned} I_k(\Delta, t; s) &= \int_0^\infty \int_{-\infty}^\infty \frac{y^{k+s-2}}{(x^2 + y^2 + ity - \frac{1}{4}\Delta)^k} dx dy \\ (2) \quad &= \frac{\Gamma(k - \frac{1}{2})\Gamma(\frac{1}{2})}{\Gamma(k)} \int_0^\infty \frac{y^{k+s-2}}{(y^2 + ity - \frac{1}{4}\Delta)^{k-\frac{1}{2}}} dy \end{aligned}$$

where the second integral converges absolutely for $1 - k < \sigma < k$ if $\Delta \neq 0$ [8, Proposition 4]. We are now in a position to state Zagier's theorem.

Theorem ([8]). *Let $k \geq 4$ be an even integer. For $m \in \mathbf{N}$, $s \in \mathbf{C}$ set*

$$\begin{aligned} c_m(s) &= m^{k-1} \sum_{t \in \mathbf{Z}} (I_k(t^2 - 4m, t; s) + I_k(t^2 - 4m, -t; s)) L(s, t^2 - 4m) \\ &+ \begin{cases} \frac{(-1)^{\frac{k}{2}} \Gamma(s+k-1) \zeta(2s)}{2^{2s+k-3} \pi^{s-1} \Gamma(k)} u^{k-s-1}, & \text{if } m = u^2, u > 0, \\ 0, & \text{if } m \text{ is not a perfect square.} \end{cases} \end{aligned}$$

Then the following assertions hold:

- i) The series converges absolutely and uniformly for $2 - k < \sigma < k - 1$.
- ii) The function

$$\Phi_s(z) = \sum_{m \geq 1} c_m(s) e^{2\pi i m z} \quad (z \in \mathcal{H}, 2 - k < \sigma < k - 1)$$

is in S_k .

- iii) Let f be a normalized Hecke eigenform in S_k . Then the Petersson scalar product of Φ_s and f is given by

$$\langle \Phi_s, f \rangle = c_k \frac{\pi^{\frac{1}{2}(s+k-1)}}{2^{s+k-1} \Gamma(\frac{1+s}{2})} D_f^*(s+k-1)$$

where

$$c_k = \frac{(-1)^{\frac{k}{2}} \pi}{2^{k-3} (k-1)}.$$

From the theorem, taking $m = 1$ we deduce

$$\begin{aligned} (3) \quad c_1(s) &= c_k \frac{\pi^{\frac{1}{2}(s+k-1)}}{2^{s+k-1} \Gamma(\frac{1+s}{2})} \sum_{\nu=1}^{g_k} \frac{1}{\langle f_{k,\nu}, f_{k,\nu} \rangle} D_{f_{k,\nu}}^*(s+k-1) \\ &\quad (2 - k < \sigma < k - 1). \end{aligned}$$

In view of the functional equation (1), it is sufficient to prove the theorem in the range $k - \frac{1}{2} + \epsilon < \sigma < k$. Suppose that the right-hand side of (3) vanishes at $s = \frac{1}{2} + \delta + it_0$ where $\epsilon < \delta < \frac{1}{2}$. Then from the definition of $c_1(s)$ we obtain

$$\begin{aligned} &\sum_{t \in \mathbf{Z}} (I_k(t^2 - 4, t; \frac{1}{2} + \delta + it_0) + I_k(t^2 - 4, -t; \frac{1}{2} + \delta + it_0)) L(\frac{1}{2} + \delta + it_0, t^2 - 4) \\ &+ (-1)^{\frac{k}{2}} \frac{\Gamma(k - \frac{1}{2} + \delta + it_0) \zeta(1 + 2\delta + 2it_0)}{2^{k-2+2\delta+2it_0} \pi^{-\frac{1}{2}+\delta+it_0} \Gamma(k)} = 0, \end{aligned}$$

or

$$(4) \quad \frac{(-1)^{\frac{k}{2}-1} 2^k \Gamma(k)}{\Gamma(k - \frac{1}{2} + \delta + it_0)} \sum_{t \in \mathbf{Z}} (I_k(t^2 - 4, t; \frac{1}{2} + \delta + it_0) + I_k(t^2 - 4, -t; \frac{1}{2} + \delta + it_0)) \\ \cdot L(\frac{1}{2} + \delta + it_0, t^2 - 4) = \frac{\zeta(1 + 2\delta + 2it_0)}{2^{2\delta-2+2it_0} \pi^{-\frac{1}{2}+\delta+it_0}}.$$

Clearly the right-hand side of (4) does not depend on k and is never zero for $\epsilon \leq \delta \leq \frac{1}{2}$. Therefore in absolute value it is bounded from below by a positive absolute constant depending only on ϵ .

We will show that the left-hand side of (4) goes to zero uniformly for $\epsilon < \delta < \frac{1}{2}$ as $k \rightarrow \infty$, thereby arriving at a contradiction.

We first look at the terms $I_k(t^2 - 4, t; \frac{1}{2} + \delta + it_0) + I_k(t^2 - 4, -t; \frac{1}{2} + \delta + it_0)$ in (4).

If $t = 0$, we obtain from (2)

$$2I_k(-4, 0; \frac{1}{2} + \delta + it_0) = 2 \frac{\Gamma(k - \frac{1}{2}) \Gamma(\frac{1}{2})}{\Gamma(k)} \int_0^\infty \frac{y^{k-\frac{3}{2}+\delta+it_0}}{(y^2 + 1)^{k-\frac{1}{2}}} dy \\ = \frac{\Gamma(k - \frac{1}{2}) \Gamma(\frac{1}{2})}{\Gamma(k)} B(\frac{k - \frac{1}{2} + \delta + it_0}{2}, \frac{k - \frac{1}{2} - \delta - it_0}{2})$$

where $B(z, w)$ is the Beta function and in the last line we have used [1, 6.2.1]. Since $B(z, w) = \frac{\Gamma(z)\Gamma(w)}{\Gamma(z+w)}$ we deduce

$$(5) \quad 2I_k(-4, 0; \frac{1}{2} + \delta + it_0) = \sqrt{\pi} \frac{\Gamma(\frac{k-\frac{1}{2}+\delta+it_0}{2}) \Gamma(\frac{k-\frac{1}{2}-\delta-it_0}{2})}{\Gamma(k)}.$$

Next suppose that $t \neq 0, \pm 2$. We will then use the fact that $I_k(t^2 - 4, t; s)$ can be expressed in terms of standard Legendre functions $P_\nu^\mu(z)$ [8, Proof of Proposition 4]. More precisely, one has

$$I_k(t^2 - 4, t; s) = \left(\frac{|t^2 - 4|}{4}\right)^{\frac{s-k}{2}} \frac{\Gamma(k - \frac{1}{2}) \sqrt{\pi}}{\Gamma(k)} \cdot \begin{cases} I_{k,s}(\frac{it}{\sqrt{|t^2-4|}}), & \text{if } t^2 < 4, \\ e^{\frac{\pi i}{2}(s-k)\text{sign}(t)} I_{k,s}(\frac{|t|}{\sqrt{t^2-4}}), & \text{if } t^2 > 4, \end{cases}$$

where

$$I_{k,s}(z) = \frac{2^{1-k} \sqrt{\pi}}{\Gamma(k - \frac{1}{2})} \Gamma(k - 1 + s) \Gamma(k - s) (z^2 - 1)^{-\frac{k-1}{2}} P_{-s}^{1-k}(z)$$

$(1 - k < \sigma < k, z \in \mathbf{C} \setminus (-\infty, 1])$. Also for $|z - 1| < 2$ the identity

$$P_{-s}^{1-k}(z) = \frac{1}{\Gamma(k)} \left(\frac{z+1}{z-1}\right)^{\frac{1-k}{2}} F(s, 1-s, k; \frac{1-z}{2})$$

holds, where $F(a, b, c; z)$ ($|z| < 1$) denotes the Gauss hypergeometric series [1, 8.1.2].

Therefore for $t = \pm 1$ we easily find that

$$(6) \quad 2(I_k(-3, 1; \frac{1}{2} + \delta + it_0) + I_k(-3, -1; \frac{1}{2} + \delta + it_0)) \\ \ll_{t_0} \frac{|\Gamma(k - \frac{1}{2} + \delta + it_0) \Gamma(k - \frac{1}{2} - \delta - it_0)|}{2^k \Gamma(k)^2},$$

and for $\pm t \geq 3$ we get

$$(7) \quad 2(I_k(t^2 - 4, t; \frac{1}{2} + \delta + it_0) + I_k(t^2 - 4, -t; \frac{1}{2} + \delta + it_0)) \ll_{t_0} (t^2 - 4)^{-\frac{1}{4} + \frac{\delta}{2}} \\ \cdot \left(\frac{|t| - \sqrt{t^2 - 4}}{|t| + \sqrt{t^2 - 4}} \right)^{\frac{k-1}{2}} \frac{|\Gamma(k - \frac{1}{2} + \delta + it_0)\Gamma(k - \frac{1}{2} - \delta - it_0)|}{2^k \Gamma(k)^2}$$

where the constants implied in $\ll_{t_0}$ depend only on t_0 .

Finally, for $t = \pm 2$ we have by [8, Proposition 4]

$$(8) \quad 2(I_k(0, 2; \frac{1}{2} + \delta + it_0) + I_k(0, -2; \frac{1}{2} + \delta + it_0)) \\ = 2(e^{\frac{\pi i}{2}(\frac{1}{2} - k + \delta + it_0)} + e^{-\frac{\pi i}{2}(\frac{1}{2} - k + \delta + it_0)})\sqrt{\pi} \\ \cdot \frac{\Gamma(\delta + it_0)\Gamma(k - \frac{1}{2} - \delta - it_0)}{\Gamma(k)} 2^{-k + \frac{1}{2} + \delta + it_0} \\ \ll_{t_0, \epsilon} \frac{|\Gamma(k - \frac{1}{2} - \delta - it_0)|}{2^k \Gamma(k)}.$$

We now look at the quantities $L(\frac{1}{2} + \delta + it_0, t^2 - 4)$ on the left-hand side of (4).

For $|t| = 2$, we have by definition

$$L(\frac{1}{2} + \delta + it_0, 0) = \zeta(2\delta + 2it_0)$$

which is a continuous function in the range $\epsilon \leq \delta \leq \frac{1}{2}$, provided $t_0 \neq 0$. If $t_0 = 0$, the same applies to the function

$$2i^k \cos\left(\frac{\pi}{2}\left(\frac{1}{2} + \delta + it_0\right)\right) \cdot \zeta(2\delta + 2it_0)$$

where

$$2i^k \cos\left(\frac{\pi}{2}\left(\frac{1}{2} + \delta + it_0\right)\right) = e^{\frac{\pi i}{2}(\frac{1}{2} - k + \delta + it_0)} + e^{-\frac{\pi i}{2}(\frac{1}{2} - k + \delta + it_0)}$$

is the second factor on the right-hand side of (8).

On the other hand, if $|t| \neq 2$, then for all $\delta \geq 0$ and all $\epsilon' > 0$ one has

$$(9) \quad L(\frac{1}{2} + \delta + it_0, t^2 - 4) \ll_{t_0, \epsilon'} |t^2 - 4|^{\frac{1}{2} + \epsilon'}.$$

In fact, if $t^2 - 4$ is a fundamental discriminant, this follows from [2, Chapter 12, Example 22 (b)], and the general case then easily follows from the definitions.

Denote the left-hand side of (4) by L_{k, δ, t_0} . Then from (5)–(9) (fixing any small $\epsilon' > 0$ in (9)) and the separate discussion in the case $|t| = 2$, $t_0 = 0$ above we deduce that

$$(10) \quad |L_{k, \delta, t_0}| \ll_{t_0, \epsilon} \left| \frac{2^k \Gamma(\frac{k - \frac{1}{2} + \delta + it_0}{2}) \Gamma(\frac{k - \frac{1}{2} - \delta - it_0}{2})}{\Gamma(k - \frac{1}{2} + \delta + it_0)} \right| + \frac{|\Gamma(k - \frac{1}{2} - \delta - it_0)|}{\Gamma(k)} \\ + \left| \frac{\Gamma(k - \frac{1}{2} - \delta - it_0)}{\Gamma(k - \frac{1}{2} + \delta + it_0)} \right| + \frac{|\Gamma(k - \frac{1}{2} - \delta - it_0)|}{\Gamma(k)} \sum_{t \geq 3} (t^2 - 4)^{\frac{1}{4} + \frac{\delta}{2} + \epsilon'} \left(\frac{t - \sqrt{t^2 - 4}}{t + \sqrt{t^2 - 4}} \right)^{\frac{k-1}{2}}.$$

Elementary considerations show that the sum over $t \geq 3$ converges and is bounded by an absolute constant independent of k .

Note that by Legendre's duplication formula for the Γ -function, the first term on the right-hand side of (10) can also be written in the form

$$2^{\frac{3}{2}-\delta} \sqrt{\pi} \left| \frac{\Gamma(\frac{k}{2} - \frac{1}{4} - \frac{\delta}{2} - \frac{it_0}{2})}{\Gamma(\frac{k}{2} + \frac{1}{4} + \frac{\delta}{2} + \frac{it_0}{2})} \right|.$$

Using the fact that

$$\lim_{x \rightarrow \infty} x^{b-a} \frac{\Gamma(x+a)}{\Gamma(x+b)} = 1 \quad (a, b \in \mathbf{C} \setminus \mathbf{R}; x \rightarrow \infty)$$

[1, 6.1.46, 6.1.47], we now see indeed that

$$(11) \quad L_{k,\delta,t_0} \rightarrow 0 \quad (k \rightarrow \infty).$$

Moreover, using more precisely the explicit asymptotics of $x^{b-a} \frac{\Gamma(x+a)}{\Gamma(x+b)}$ for $x \rightarrow \infty$ given in [1, 6.1.47], one sees that the convergence in (11) is uniform in δ , since $\delta > \epsilon > 0$ is bounded away from zero.

This proves the theorem.

REFERENCES

- [1] N. Abramowitz and I. Stegun: "Handbook of Mathematical Functions", Dover, New York, 1965
- [2] T. Apostol: "Introduction to Analytic Number Theory", Springer, Berlin-Heidelberg-New York, 1976 MR **55**:7892
- [3] S. Gelbart and H. Jacquet: A relation between automorphic representations of GL_2 and GL_3 , Ann. Sci. E.N.S. IV ser., 11, 471-542 (1978) MR **81e**:10025
- [4] H. Jacquet and J. A. Shalika: A nonvanishing theorem for zeta functions of GL_n , Invent. Math. 38, 1-16 (1976/77) MR **55**:5583
- [5] W. Kohnen: Nonvanishing of Hecke L -functions associated to cusp forms inside the critical strip, J. of Number Theory vol. 67, no. 2, 182-189 (1997) MR **98j**:11037
- [6] X.-J. Li: On the poles of Rankin-Selberg convolutions of modular forms, Trans. Amer. Math. Soc. 348, no. 3, 1213-1234 (1996) MR **96h**:11038
- [7] G. Shimura: On the holomorphy of certain Dirichlet series, Proc. London Math. Soc. 31, 75-98 (1975) MR **52**:3064
- [8] D. Zagier: Modular forms whose Fourier coefficients involve zeta-functions of quadratic fields, in "Modular Functions of One Variable VI" (eds. J.-P. Serre and D. Zagier), pp. 105-169, LNM no. 627, Springer, Berlin-Heidelberg-New York, 1976. MR **58**:5525

UNIVERSITÄT HEIDELBERG, MATHEMATISCHES INSTITUT, IM NEUENHEIMER, FELD 288, D-69120 HEIDELBERG, GERMANY

E-mail address: winfried@mathi.uni-heidelberg.de

SCHOOL OF MATHEMATICS, TATA INSTITUTE FOR FUNDAMENTAL RESEARCH, HOMI BHABHA ROAD, BOMBAY 400 005, INDIA

E-mail address: sengupta@math.tifr.res.in