UNIQUENESS OF THE LEAST-ENERGY SOLUTION FOR A SEMILINEAR NEUMANN PROBLEM

MASSIMO GROSSI

(Communicated by Lesley M. Sibner)

ABSTRACT. We prove that the least-energy solution of the problem

$$\begin{cases} -d\Delta u + u = u^p & \text{in } B, \\ u > 0 & \text{in } B, \\ \frac{\partial u}{\partial \nu} = 0 & \text{on } \partial B, \end{cases}$$

where B is a ball, $d > 0$ and $1 < p < \frac{N+2}{N-2}$ if $N \geq 3$, $p > 1$ if $N = 2$, is unique (up to rotation) if d is small enough.

INTRODUCTION

In this paper we consider the following problem:

$$(0.1) \quad \begin{cases} -d\Delta u + u = u^p & \text{in } \Omega, \\ u > 0 & \text{in } \Omega, \\ \frac{\partial u}{\partial \nu} = 0 & \text{on } \partial\Omega, \end{cases}$$

where Ω is a bounded smooth domain in $\mathbb{R}^N$, ν is the unit outer normal to $\partial\Omega$, $d > 0$ and $1 < p < \frac{N+2}{N-2}$ if $N \geq 3$, $p > 1$ if $N = 2$.

To our knowledge the first existence result to (0.1) is due to Lin, Ni and Takagi (see [LNT]). In this paper the authors consider the functional

$$(0.2) \quad J_d(u) = \frac{1}{2} \int_{\Omega} (d|\nabla u|^2 + u^2) - \frac{1}{p+1} \int_{\Omega} (u^+)^{p+1}$$

and

$$(0.3) \quad c_d = \inf_{h \in \Gamma} \max_{t \in [0,1]} J_d(h(t))$$

where $\Gamma = \{f \in C([0,1], H^1(\Omega)) \mid f(0) = 0, f(1) = e\}$ and $e \neq 0$ is a non-negative function of $H^1(\Omega)$ such that $J_d(e) = 0$. From the mountain-pass theorem (see [AR]) we have that c_d is a positive critical value of J_d . Moreover, it turns out that c_d is the least positive critical value for J_d (see [LNT] or [NT]).

Therefore, we define a critical point u_d of J_d satisfying $J_d(u_d) = c_d$ a *least-energy solution* of (0.1).

Received by the editors July 9, 1998.

1991 *Mathematics Subject Classification*. Primary 35J70.

Key words and phrases. Uniqueness results, semilinear elliptic equations, Neumann problem.

This research was supported by M.U.R.S.T. (Project “Metodi Variazionali ed Equazioni Differenziali Non Lineari”).

Clearly (0.1) admits the trivial solutions $u \equiv 1$ and $u \equiv 0$. In [LNT] it was proved that $c_d = O(d^{N/2})$ as $d \rightarrow 0$ and from this $0 < J_d(u_d) < J_d(1)$, i.e. u_d is a nonconstant solution provided d is small enough.

Another important result concerns the shape of the least-energy solution u_d . In [NT] it was shown that u_d has only one local maximum in $\overline{\Omega}$, and it is achieved at exactly one point which lies on the boundary of Ω , provided d is small enough.

In this paper we will prove the uniqueness of the least-energy solution if Ω is a ball. Of course, in this case, if $u(x)$ is a solution of (0.1), then also $u(Tx)$ solves (0.1) for any T belonging to the orthogonal group $O(N)$; therefore by uniqueness we mean that any two solutions can be obtained from each other using the action of $O(N)$.

Theorem 1. *Let us consider the problem (0.1) when Ω is a ball. Then there exists a $d_0 > 0$ such that for any $d < d_0$ the least-energy solution of (0.1) is unique.*

We want to point out that uniqueness results for (0.1) cannot be obtained if we do not restrict the class of the solutions of (0.1). Indeed, in [DY] Dancer and Yan prove the existence of k -peak solutions to (0.1) (solutions with more than one local maximum point). Clearly the energy of these solutions is strictly greater than c_d . We mention here the papers of [BDS], [Gu], [W1], and [W2] for the existence of multipeak solutions in other domains.

The paper is organized as follows: in Section 1 we recall some important results which we use in the proof of Theorem 1. In Section 2 we prove Theorem 1.

1. KNOWN RESULTS

In this section we recall some known facts which will be useful in the proof of Theorem 1. Here Ω is an arbitrary bounded smooth domain of $\mathbb{R}^N$ and p is a real number such that $1 < p < \frac{N+2}{N-2}$ if $N \geq 3$ and $p > 1$ if $N = 2$. We start by recalling some results on the existence and the shape of the least-energy solution u_d .

Theorem 1.1. *Let $p > 1$ if $N = 1$ or 2 and let $1 < p < \frac{N+2}{N-2}$ if $N \geq 3$. Then there exists a $d_0 > 0$ such that, for any $d < d_0$, (0.1) has a positive nonconstant solution u_d satisfying*

$$(1.1) \quad J_d(u_d) = c_d,$$

$$(1.2) \quad \int_{\Omega} u_d^q \leq C_1(q) d^{N/2} \quad \text{for any } q \geq 1,$$

and

$$(1.3) \quad \sup_{\Omega} u_d(x) \leq C_2$$

with $C_1(q), C_2$ independent of d .

Proof. See [LNT]. □

Theorem 1.2. *Let u_d be a least-energy solution to (1.3). Then u_d has at most one local maximum in Ω and it is achieved exactly at one point which must lie on the boundary, provided d is sufficiently small.*

Proof. See [NT], Theorem 2.1. □

Theorem 1.3. *There is exactly one solution (up to translation) of the problem*

$$(1.4) \quad \begin{cases} -\Delta u + u = u^p & \text{in } \mathbb{R}^N, \\ u > 0 & \text{in } \mathbb{R}^N, \\ u(x) \rightarrow 0 & \text{as } |x| \rightarrow \infty. \end{cases}$$

Proof. See [K]. □

Next let us introduce a diffeomorphism which straightens a boundary portion of Ω as follows. We denote by B_R the ball of $\mathbb{R}^N$ centered at the origin and radius R . Through translation and rotation of the coordinate system we may assume that P is the origin and the inner normal to $\partial\Omega$ at P is pointing in the direction of the positive x_N -axis. Then there exists a smooth function $\psi(x')$, $x' = (x_1, \dots, x_{N-1})$, defined for $|x| < \delta_0$ such that $\psi(0) = 0$, $\nabla\psi(0) = 0$ and

$$\partial\Omega \cap \mathcal{N} = \{x = (x', x_N) \mid x_N = \psi(x')\}$$

and

$$\Omega \cap \mathcal{N} = \{x = (x', x_N) \mid x_N > \psi(x')\}.$$

For $|y| < \delta'$ let us define $x = \Phi(y)$, with $\Phi(y) = (\Phi_1(y), \dots, \Phi_N(y))$, by

$$(1.5) \quad \Phi_j(y) = \begin{cases} y_j - y_N \frac{\partial\psi}{\partial x_j}(y') & \text{for } j = 1, \dots, N-1, \\ y_N + \psi(y') & \text{if } j = N. \end{cases}$$

Since $\nabla\psi(0) = 0$ the differential map $D\Phi$ of Φ satisfies $D\Phi(0) = I$, the identity map. Thus there exists the inverse map $y = \Phi^{-1}(x)$ between $C \subset \mathbb{R}^N$ and $B(0, \delta)$ with $\delta < \delta'$.

Theorem 1.4. *Let u_d be a least-energy solution to (0.1) and let us denote by P_d the point where the maximum of u_d is achieved. Then for any $\epsilon > 0$ there exist a positive constant d_0 and a subdomain $\Omega_d^{(i)} \subset \Omega$ such that for $d \in]0, d_0[$ the following statements hold:*

$$(1.6) \quad P_d \in \partial\Omega_d \quad \text{and} \quad \text{diam}(\Omega_d^{(i)}) \leq C\sqrt{d},$$

$$(1.7) \quad u_d(\Phi(\sqrt{d}y)) \rightarrow U(y) \text{ in } C_{\text{loc}}^2(\mathbb{R}_+^N) \text{ as } d \rightarrow 0,$$

$$(1.8) \quad |u_d(x)| \leq C_1 \epsilon e^{-\frac{\mu_1 \delta(x)}{\sqrt{d}}} \quad \text{for } x \in \Omega \setminus \Omega_d^{(i)},$$

where $\delta(x) = \min\{\text{dist}(x, \partial\Omega_d), \eta_0\}$, C, C_1, μ_1, η_0 are positive constants depending only on Ω and $U(x)$ is the unique solution of (1.4) with maximum achieved at $P_0 = \lim_{d \rightarrow 0} P_d$.

Proof. See [NT], (i) and (iii) of Theorem 2.3 and (4.29). □

Theorem 1.5. *Let us consider the problem*

$$(1.9) \quad \begin{cases} -\Delta z + z = pU(x)^{p-1}z & \text{in } \mathbb{R}^N, \\ \|z\|_{L^\infty(\mathbb{R}^N)} \leq C, \end{cases}$$

where $U(x)$ is the unique solution of (1.4) symmetric with respect to the origin. Then the only nontrivial solutions of (1.9) are given by

$$(1.10) \quad z = \sum_{i=1}^N \alpha_i \frac{\partial U}{\partial x_i}, \quad \alpha_i \in \mathbb{R}.$$

Proof. This result is contained in the proof of Proposition 1 of [D]. □

2. THE UNIQUENESS RESULT

In this section we denote by $x = (x', x_N) \in \mathbb{R}^{N-1} \times \mathbb{R}$ a generic point of $\mathbb{R}^N$. Let us assume that $\Omega = B = \{x \in \mathbb{R}^N \mid |x'|^2 + (x_N - 1)^2 < 1\}$. From Theorem 1.2 the maximum of a least-energy solution u of (0.1) is achieved on the boundary of B if d is small enough. So, since the problem (0.1) is invariant with respect to the orthogonal group of $\mathbb{R}^N$, by a suitable rotation we can assume that u attains its maximum at the origin. Hence $\nabla u(0) = 0$ for any least-energy solution of (0.1).

We suppose by contradiction that there exist a sequence $d_n \searrow 0$ and two distinct least-energy solutions $u_{1,n}$ and $u_{2,n}$ which solve (0.1) with $d = d_n$. So we can suppose that $u_{1,n}$ and $u_{2,n}$ achieve their maximum at the origin for any $n \in \mathbb{N}$. Since $u_{1,n} - u_{2,n} \not\equiv 0$ we can consider

$$(2.1) \quad z_n(x) = \frac{u_{1,n}(x) - u_{2,n}(x)}{\|u_{1,n} - u_{2,n}\|_{L^\infty}}.$$

By assumption on $u_{1,n}$ and $u_{2,n}$ we have

$$(2.2) \quad \nabla z_n(0) = 0.$$

Note that z_n satisfies

$$(2.3) \quad \begin{cases} -d_n \Delta z_n + z_n = c_n(x) z_n & \text{in } B, \\ \frac{\partial z_n}{\partial \nu} = 0 & \text{on } \partial B, \\ \|z_n\|_{L^\infty(B)} = 1, \end{cases}$$

where

$$(2.4) \quad c_n(x) = p \int_0^1 (t u_{1,n}(x) + (1-t) u_{2,n}(x))^{p-1} dt.$$

Let us set

$$(2.5) \quad \tilde{z}_n(x) = z_n(\sqrt{d_n}x), \quad \tilde{z}_n : B/\sqrt{d_n} \mapsto \mathbb{R}.$$

We remark that for any compact set $K \subset \mathbb{R}_+^N$, where $\mathbb{R}_+^N = \{x \in \mathbb{R}^N \mid x_N > 0\}$, we have that $K \subset B/\sqrt{d_n}$ for n large enough. In the following proposition we study the asymptotic behavior of $\tilde{z}_n$.

Proposition 2.1. *We have that*

$$(2.6) \quad \tilde{z}_n \rightarrow z \quad \text{in } C_{\text{loc}}^1(\mathbb{R}_+^N),$$

where z is a bounded solution of

$$-\Delta z + z = pU^{p-1}z \quad \text{in } \mathbb{R}^N$$

and U is the unique solution of (1.4) symmetric with respect to the origin.

Proof. Let us consider the diffeomorphism Φ between $B(0, \delta)$ and $C \subset \mathbb{R}^N$ which straightens the boundary of B (see (1.5)). Now, as in [NT], let us set

$$(2.7) \quad v_n(y) = z_n(\Phi(y)), \quad y \in \overline{B_{2k}^+},$$

where $B_{2k}^+ = \{y \in B_{2k} \mid y_N > 0\}$.

Step 1. In this step we prove that

$$(2.8) \quad v_n(\sqrt{d_n}y) \rightarrow z(y) \quad \text{in } C_{\text{loc}}^2(\mathbb{R}_+^N).$$

First of all we extend v_n to $B(0, 2k)$ by reflection,

$$\tilde{v}_n(y) = \begin{cases} v_n(y) & \text{if } y \in \overline{B_{2k}^+}, \\ v_n(y', -y_N) & \text{if } y \in \overline{B_{2k}^-}, \end{cases}$$

with $\overline{B_{2k}^-} = \{y \in B_{2k} \mid y_N < 0\}$. Moreover we define a scaled function $w_n(z)$ by

$$(2.9) \quad w_n(y) = \tilde{v}_n(\sqrt{d_n}y) \quad \text{for } y \in \overline{B_{\frac{k}{\sqrt{d_n}}}}.$$

It is easily seen that

$$w_n \in C^2(\overline{B_{\frac{k}{\sqrt{d_n}}}} \setminus \{y_N = 0\}) \cap C^1(\overline{B_{\frac{k}{\sqrt{d_n}}}})$$

since $\frac{\partial v_n}{\partial y_N}$ on $y_N = 0$ and that w_n satisfies the elliptic equation

$$(2.10) \quad -L_n w_n + w_n = \tilde{c}_n(y) w_n$$

with

$$(2.11) \quad L_n w_n = \sum_{i,j=1}^N a_{ij}^n(y) \frac{\partial^2 w_n}{\partial z_i \partial z_j} + \sqrt{d_n} \sum_{j=1}^N b_j^n(y) \frac{\partial w_n}{\partial z_j}$$

in $B_{\frac{k}{\sqrt{d_n}}} \setminus \{y_N = 0\}$, where a_{ij}^n , b_j^n and $\tilde{c}_n(y)$ are defined as follows. First, put

$$\Psi = \Phi^{-1},$$

$$\begin{aligned} a_{ij}(y) &:= \sum_{l=1}^N \frac{\partial \Psi_i}{\partial x_l}(\Phi(y)) \frac{\partial \Phi_j}{\partial x_l}(\Phi(y)), \quad 1 \leq i, j \leq N, \\ b_j(y) &:= (\Delta \Psi_j)(\Phi(y)), \quad j = 1, \dots, N, \\ \hat{c}_n(y) &= c_n(\Phi(y)). \end{aligned}$$

Then set

$$\begin{aligned} a_{ij}^n(y) &:= \begin{cases} a_{ij}(\sqrt{d_n}y) & \text{if } y_N \geq 0, \\ (-1)^{\delta_N^i + \delta_N^j} a_{ij}(\sqrt{d_n}y', -\sqrt{d_n}y_N) & \text{if } y_N < 0, \end{cases} \\ b_j^n(y) &:= \begin{cases} b_j(\sqrt{d_n}y) & \text{if } y_N \geq 0, \\ (-1)^{\delta_N^j} b_j(\sqrt{d_n}y', -\sqrt{d_n}y_N) & \text{if } y_N < 0, \end{cases} \\ \tilde{c}_n(y) &:= \begin{cases} \hat{c}_n(\sqrt{d_n}y) & \text{if } y_N \geq 0, \\ (-1)^{\delta_N^j} \hat{c}_n(\sqrt{d_n}y', -\sqrt{d_n}y_N) & \text{if } y_N < 0, \end{cases} \end{aligned}$$

where δ_j^i is the Kroneker symbol.

Note that by (1.3) the functions w_n and c_n are uniformly bounded with respect to n . Then, by using the standard L^r -estimate and the interior Schauder estimate as in [NT], pp. 834-836, we obtain that

$$(2.12) \quad w_n \rightarrow z \quad \text{in } C_{\text{loc}}^2(\mathbb{R}^N)$$

and

$$(2.13) \quad -L_n w_n \rightarrow -\Delta z.$$

Hence, since by (1.7) and (2.4) we get $\tilde{c}_n \rightarrow pU^{p-1}$, z is a solution of

$$(2.14) \quad -\Delta z + z = pU^{p-1}z \quad \text{in } \mathbb{R}^N.$$

By the definition of w_n and $\tilde{v}_n$ (2.8) follows.

Step 2. In this step we prove the claim of the proposition.

The function $\tilde{z}_n$ satisfies the equation

$$(2.15) \quad \begin{cases} -\Delta \tilde{z}_n + \tilde{z}_n = c_n(\sqrt{d_n})\tilde{z}_n & \text{in } B/\sqrt{d_n}, \\ \tilde{z}_n > 0 & \text{in } B/\sqrt{d_n}, \\ \frac{\partial \tilde{z}_n}{\partial \nu} = 0 & \text{on } \partial B/\sqrt{d_n}. \end{cases}$$

From (1.3) we get $\|c_n \tilde{z}_n\|_\infty \leq C$ and then by standard L^p -estimates we deduce, for a subsequence, $\|\tilde{z}_n\|_{C^1(K)} \leq C$ for any compact set $K \subset \mathbb{R}^N$.

By Step 1 we know that $\|z_{d_n}(\Phi(\sqrt{d_n}y)) - U(y)\|_{C_{\text{loc}}^0(\mathbb{R}_+^N)} \rightarrow 0$. So if we show that

$$(2.16) \quad \|z_{d_n}(\Phi(\sqrt{d_n}y)) - \tilde{u}_n(y)\|_{C_{\text{loc}}^0(\mathbb{R}_+^N)} \rightarrow 0,$$

the claim follows.

Let us fix a compact set $K \subset \mathbb{R}^N$. By the definition of Φ (see (1.5)) we get

$$(2.17) \quad |\Phi(\sqrt{d_n}y) - \sqrt{d_n}y|^2 = |d_n y_N^2 |\nabla_{y'} \psi(\sqrt{d_n}y)|^2 + \psi^2(\sqrt{d_n}y) \rightarrow \psi^2(0) = 0$$

uniformly in K .

Then, since $|\nabla u_{d_n}| \leq C$ in K , we obtain that

$$(2.18) \quad \|z_{d_n}(\Phi(\sqrt{d_n}y)) - \tilde{z}_n(y)\|_{C^0(K)} \leq C \|\Phi(\sqrt{d_n}y) - \sqrt{d_n}y\|_{C^0(K)} \rightarrow 0,$$

and this finishes the proof. □

Proof of Theorem 1. Let us again consider the functions z_n and $\tilde{z}_n$ defined by (2.1) and (2.5). First of all we remark that since $u_{1,n}$ and $u_{2,n}$ solve (0.1) we have

$$\int_{\Omega} u_{1,n} u_{2,n} (u_{1,n}^{p-1} - u_{2,n}^{p-1}) = 0$$

and then $\tilde{z}_n$ does change sign.

Since we have that $\|\tilde{z}_n\|_{L^\infty(B/\sqrt{d_n})} = 1$ we can assume that there exists a sequence of points $\tilde{x}_n \in B/\sqrt{d_n}$ such that $\tilde{z}_n(\tilde{x}_n) = \max_{x \in B/\sqrt{d_n}} \tilde{z}_n(x) = 1$. Moreover we have that

$$(2.19) \quad \nabla \tilde{z}_n(\tilde{x}_n) = 0 \quad \text{and} \quad \Delta \tilde{z}_n(\tilde{x}_n) \leq 0$$

(this is obvious if $\tilde{x}_n \in B/\sqrt{d_n}$ and it follows by $\frac{\partial \tilde{z}_n}{\partial \nu}(\tilde{x}_n) = 0$ if $\tilde{x}_n \in \frac{\partial B}{\sqrt{d_n}}$).

From Proposition 2.1 the following alternative occurs: either

$$(2.20) \quad \tilde{z}_n \rightarrow 0$$

or

$$(2.21) \quad \tilde{z}_n \rightarrow \sum_{i=1}^N \alpha_i \frac{\partial U}{\partial x_i} \quad \text{for some } (\alpha_1, \dots, \alpha_N) \in \mathbb{R}^N \setminus \{0\}.$$

We will prove that in any case we reach a contradiction.

Case 1. $|\tilde{z}_n| \rightarrow z \equiv 0$.

Here we have the following alternative:

- i) $|\tilde{x}_n| \leq C$,
- ii) $|\tilde{x}_n| \rightarrow \infty$.

Case i). $|\tilde{x}_n| \leq C$.

Let $x_0 = \lim_n \tilde{x}_n$. If $x_0 \notin \partial\mathbb{R}_+^N$, the claim follows easily. If $x_0 \in \partial\mathbb{R}_+^N$, we again consider the diffeomorphism Φ between $B(0, \delta')$ and $B(0, \delta)$ which straightens the boundary. Since $\tilde{x}_n \in B/\sqrt{d_n}$ we have that $\tilde{x}_n = \frac{x_n}{\sqrt{d_n}}$ with $x_n \in B$ and $x_n \rightarrow 0$. Then there exists $P_n \in B_\delta^+$ such that

$$(2.22) \quad \Phi(P_n) = x_n.$$

Set $y_n = \frac{P_n}{\sqrt{d_n}}$. Then, since $\Phi(0) = 0$ and Φ is a diffeomorphism in B_δ^+ ,

$$(2.23) \quad |\sqrt{d_n}y_n| = |P_n| = |\Phi^{-1}(x_n)| = |\Phi^{-1}(\sqrt{d_n}\tilde{x}_n)| \leq C|\sqrt{d_n}\tilde{x}_n|,$$

so $|y_n| \leq C$. Finally, by (2.7)-(2.9)

$$(2.24) \quad 1 = \tilde{z}_n(\tilde{x}_n) = z_n(x_n) = z_n(\Phi(P_n)) = z_n(\Phi(\sqrt{d_n}y_n)) \rightarrow 0$$

and this gives a contradiction.

Case ii). $|\tilde{x}_n| \rightarrow \infty$.

Let us prove that $c_n(\sqrt{d_n}\tilde{x}_n) \rightarrow 0$. We have

$$c_n(\sqrt{d_n}\tilde{x}_n) = p \int_0^1 (tu_{1,n}(\sqrt{d_n}\tilde{x}_n) + (1-t)u_{2,n}(\sqrt{d_n}\tilde{x}_n))^{p-1} dt.$$

Again since $\tilde{x}_n \in B/\sqrt{d_n}$ there exists $x_n \in B$ such that $x_n = \sqrt{d_n}\tilde{x}_n$ and

$$(2.25) \quad \frac{|x_n|}{\sqrt{d_n}} \rightarrow \infty.$$

Then (1.6) and (1.7) of Theorem 1.4 applies: since by (2.25) $x_n \notin B_{d_n}^{(i)}$ we obtain, for any $\epsilon > 0$,

$$(2.26) \quad u_{1,n}(x_n) < \epsilon \text{ and } u_{2,n}(x_n) < \epsilon \text{ for } n \text{ large.}$$

Hence $c_n(\sqrt{d_n}\tilde{x}_n) = c_n(x_n) \rightarrow 0$. Finally from (2.19) we obtain

$$(2.27) \quad 0 \leq -\Delta \tilde{z}_n(\tilde{x}_n) = c_n(x_n) - 1 \rightarrow -1,$$

and this gives a contradiction.

Case 2. $|\tilde{z}_n| \rightarrow z = \sum_{i=1}^N \alpha_i \frac{\partial U}{\partial x_i}$ for some $(\alpha_1, \dots, \alpha_N) \in \mathbb{R}^N \setminus \{0\}$.

First of all let us prove that

$$(2.28) \quad \frac{\partial^2 U}{\partial x_j^2}(0) < 0 \quad \text{and} \quad \frac{\partial^2 U}{\partial x_i \partial x_j}(0) = 0 \text{ if } i \neq j, \quad i, j \in \{1, \dots, N\}.$$

This is a well-known fact (see [NT] for example), but we repeat the proof for the reader's convenience. Since the function $U(\rho)$ is radially decreasing for $\rho > 0$ (see [GNN]), we have that $\frac{\partial U}{\partial x_j} = U'(\rho) \frac{x_j}{\rho} > 0$ in $T_j = \{x \in \mathbb{R}^N \mid x_j < 0\}$. Hence $\frac{\partial U}{\partial x_j}|_{\partial T_j} \equiv 0$ and then $\frac{\partial^2 U}{\partial x_i \partial x_j}(0) = 0$ for any $i \neq j$. Moreover $f = \frac{\partial U}{\partial x_j}$ solves

$$\begin{cases} -\Delta f + f = pU(x)^{p-1}f & \text{in } T_j, \\ f > 0 & \text{in } T_j, \\ f = \frac{\partial U}{\partial x_j} = 0 & \text{on } \partial T_j. \end{cases}$$

So, by Hopf's Lemma, $\frac{\partial f}{\partial x_j} = \frac{\partial^2 U}{\partial x_j^2}(0) < 0$ on ∂T_j , and then (2.28) is proved.

Now let us recall that $\nabla \tilde{z}_n(0) = 0$. Since $\Phi(0) = 0$ we have

$$(2.29) \quad \nabla(z_n \circ \Phi)(0) = 0.$$

Then since $(\alpha_1, \dots, \alpha_N) \neq (0, \dots, 0)$ there exists $\alpha_j \neq 0$. Finally, from (2.8), (2.28) and (2.29) we get

$$(2.30) \quad 0 = \frac{\partial(z_n \circ \Phi)}{\partial x_j}(0) \rightarrow \sum_{i=1}^N \alpha_i \frac{\partial^2 U}{\partial x_i \partial x_j}(0) = \alpha_j \frac{\partial^2 U}{\partial x_j^2} \neq 0,$$

a contradiction. The proof of Theorem 1 is finished.

REFERENCES

- [AR] A. Ambrosetti and P. H. Rabinowitz, "Dual variational methods in critical point theory and applications", J. Funct. Anal. **14**, 349-381, (1973). MR **51**:6412
- [BDS] P. Bates, E.N. Dancer and J. Shi, "Multi-spike stationary solutions on the Cahn-Hilliard equation in higher dimension and instability", preprint.
- [D] E.N. Dancer, "On the uniqueness of the positive solution of a singularly perturbed problem", Rocky Mountain Journal of Mathematics **25** (1995), 957-975. MR **96j**:35021
- [DY] E.N. Dancer and S. Yan, "Multipeak solutions for a singularly perturbed Neumann problem" (to appear).
- [GNN] B. Gidas, W.N. Ni and L. Nirenberg, "Symmetry of positive solutions of nonlinear elliptic equation in $\mathbb{R}^N$ ", Advances in Math. Studies **7 A**, 369-402, (1981). MR **84a**:35083
- [GT] D. Gilbarg and N. Trudinger, "Elliptic partial differential equations of second order", Springer Verlag (1983). MR **86c**:35035
- [Gu] C. Gui, "Multipeak solutions for a semilinear Neumann problem", Duke Math. J. **84**, 739-769, (1996). MR **97i**:35052
- [K] M.K. Kwong, "Uniqueness of positive solutions of $\Delta u - u + u^p = 0$ in $\mathbb{R}^N$ ", Arch. Rat. Mech. Anal., **105**, 243-266, (1989). MR **90d**:35015
- [LN] C.S. Lin and W.M. Ni, "On the diffusion coefficient of a semilinear Neumann problem, Calculus of Variations and Partial Differential Equations", S. Hildebrandt, D. Kinderlehrer and M. Miranda, eds., Lecture Notes in Math. 1340, Springer-Verlag, Berlin-Heidelberg-New York-Tokyo, 160-174, (1988). MR **90d**:35101
- [LNT] C.S. Lin, W.M. Ni and I. Takagi, "Large amplitude stationary solutions to a chemotaxis system", J. Diff. Eqns. **72**, 1-27, (1988). MR **89e**:35075
- [NT] W.M. Ni and I. Takagi, "On the shape of Least Energy Solutions to a Semilinear Neumann Problem", Comm. Pure Math. Appl., Vol **XLIV**, 819-851, (1991). MR **92i**:35052
- [W1] Z.Q. Wang, "On the existence of multiple single-peaked solution for a semilinear Neumann problem", Arch. Rat. Mech. Anal., **120**, 375-399, (1992). MR **93k**:35109
- [W2] Z.Q. Wang, "Nonradial solutions of nonlinear Neumann problems in radially symmetric domains", Topology in Nonlinear analysis (Warsaw, 1994), 85-96, Banach Center Publications, **35**, Polish Acad. Sci., Warsaw, (1996). MR **98e**:35075

DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DI ROMA "LA SAPIENZA", P.LE A. MORO 2, 00185, ROMA, ITALY

E-mail address: grossi@mat.uniroma1.it