NORMALIZERS OF THE CONGRUENCE SUBGROUPS OF THE HECKE GROUP G_5 II

MONG-LUNG LANG AND SER-PEOW TAN

(Communicated by Ronald M. Solomon)

ABSTRACT. Let $\lambda = 2\cos(\pi/5)$. Let (τ) be an ideal of $\mathbb{Z}[\lambda]$ and let (τ_0) be the maximal ideal of $\mathbb{Z}[\lambda]$ such that $(\tau_0^2) \subseteq (\tau)$. Then $N(G_0(\tau)) \leq G_0(\tau_0)$. In particular, if τ is square free, then $G_0(\tau)$ is self-normalized in $PSL_2(\mathbb{R})$.

1. INTRODUCTION

In this paper, we continue our study into the extent to which properties of the modular group hold for the Hecke groups; see [CLLT], [LLT1], [LLT2], [LT1], [LT2] for some previous results. We are in particular interested in the Hecke group G_5 which we denote by G and its congruence subgroups $G_0(\tau)$. Our main results are:

- (i) **Theorem 8.** Let $\lambda = 2\cos(\pi/5) = (1 + \sqrt{5})/2$. Let (τ) be an ideal of $\mathbb{Z}[\lambda]$ and let (τ_0) be the maximal ideal of $\mathbb{Z}[\lambda]$ such that $(\tau_0^2) \subseteq (\tau)$. Then $N(G_0(\tau)) \leq G_0(\tau_0)$.
- (ii) **Main Theorem.** Let (τ) be an ideal of $\mathbb{Z}[\lambda]$. Suppose that τ is square free. Then $G_0(\tau)$ is self-normalized in $PSL_2(\mathbb{R})$.

This contrasts with the cases of the congruence subgroups of the modular group Γ which admit Atkin-Lehner involutions, so have strictly larger normalizers ([AL], [C], [LN]).

We recall the following definitions, notation and results. For $q \geq 4$, the Hecke groups G_q are the (discrete) subgroups $\langle w, u_q \rangle$ of $PSL_2(\mathbb{Z}[\lambda_q])$, where $\lambda_q = 2\cos(\pi/q)$ and

$$w = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}, \quad u_q = \begin{pmatrix} 1 & \lambda_q \\ 0 & 1 \end{pmatrix}.$$

When $q = 3$, we recover the modular group Γ so the above can be thought of as a natural generalization of Γ . Alternatively, we can interpret the generalization as G_q being maximal discrete subgroups whose entries are in some extension of $\mathbb{Z}$. Finally, we have the geometric interpretation: Γ is a $(2, 3, \infty)$ triangle group and the Hecke group G_q is a $(2, q, \infty)$ triangle group.

Received by the editors September 25, 1998.

1991 *Mathematics Subject Classification.* Primary 11F06.

Key words and phrases. Congruence subgroups, Hecke groups.

Let $\mathcal{A}$ be an ideal of $\mathbb{Z}[\lambda_q]$. We define

$$G_0(\mathcal{A}) = \left\{ \sigma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in G_q \mid c \in \mathcal{A} \right\}.$$

Again, this is a natural generalization of the congruence subgroups $\Gamma_0(n)$ of Γ . It works because the elements of G_q sit naturally in the ring $\mathbb{Z}[\lambda_q]$.

Recall that G_q is commensurable with $PSL_2(\mathbb{Z})$ if and only if $q = 4$ or 6 . The elements of such groups are completely known; see [P], for example.

Suppose G_q is not commensurable with $PSL_2(\mathbb{Z})$. By the results of Leutbecher, [L1], [L2], $\mathbb{Q}(\lambda) \cup \{\infty\}$ is the set of cusps of G_q if and only if $q = 5$. Also, 5 is the only q other than 4, 6 for which $\mathbb{Q}(\lambda)$ is a quadratic field. For all other q 's, the degree is > 2 . As a consequence, $q = 5$ is the next most workable and interesting q . Some of the classical results on the modular group can be generalized to $G = G_5$ ([CLLT], [LLT2]). From now on $q = 5$, so $\lambda_q = (1 + \sqrt{5})/2$.

The main facts used in the proof of our main theorem are:

- (a) $\mathbb{Z}[\lambda]$ is a principal ideal domain.
- (b) The set of cusps of G is $\mathbb{Q}(\lambda) \cup \{\infty\}$ ([L1], [L2]). Furthermore, if $x \in \mathbb{Q}(\lambda)$ is a cusp, x has a unique reduced form $x = \frac{a}{b}$ ([R]). By definition, this means that $a, b \in \mathbb{Z}[\lambda]$ with $b > 0$ and there exists $c, d \in \mathbb{Z}[\lambda]$ such that $\begin{pmatrix} a & c \\ b & d \end{pmatrix} \in G$.

Clearly, $(a, b) = 1$ so that if $x = \frac{a'}{b'}$ with $(a', b') = 1$, then $a = \mu a'$, $b = \mu b'$ where μ is a unit in $\mathbb{Z}[\lambda]$.

- (c) (Proposition 6 of [LLT1]) Suppose x_i, x_j are members in $\mathbb{Q}(\lambda) \cup \{\infty\}$ with reduced form a_i/b_i and a_j/b_j respectively, and suppose that $x_i < x_j$. Then the following statements are equivalent:
 - (i) $\begin{pmatrix} a_j & a_i \\ b_j & b_i \end{pmatrix} \in G$;
 - (ii) (x_i, x_j) is an even line, that is, it is the image of the complete hyperbolic geodesic with ends at 0 and ∞ under the action of some $A \in G$;
 - (iii) $a_j b_i - a_i b_j = 1$.
- (d) The even lines give a tessellation of $\mathbb{H}$ (the upper half plane) into ideal pentagons, that is, hyperbolic 5-gons whose vertex angles are all zero ([K], [LLT1]).
- (e) $\begin{pmatrix} 1 & b \\ 0 & 1 \end{pmatrix} \in G$ if and only if $b = m\lambda$, $m \in \mathbb{Z}$ ([R]). Similarly, $\begin{pmatrix} 1 & 0 \\ c & 1 \end{pmatrix} \in G$ if and only if $c = n\lambda$, $n \in \mathbb{Z}$. (This result is well known, a proof of this result can be found in Corollary 5 of [LLT1].)

The rest of the paper is organized as follows. In Sections 2 and 3, we study the reduced forms of $\mathbb{Q}(\lambda)$. Facts about reduced forms that are useful in the determination of $N(G_0(\tau))$ are given in Corollary 3 and Lemma 4. The next two sections are devoted to the reduction of the supergroups of $N(G_0(\tau))$ (a group S is a supergroup of $N(G_0(\tau))$ if $N(G_0(\tau))$ is a subgroup of S). In Section 6, we complete the proof of our main theorem.

Throughout the paper, $\lambda = 2\cos(\pi/5)$, (τ) is an ideal of $\mathbb{Z}[\lambda]$, and $N(G_0(\tau))$ is the normalizer of $G_0(\tau)$ in $PSL_2(\mathbb{R})$.

2. REDUCED FORMS

Recall first that a/b is a reduced form if

- (i) $a, b \in \mathbb{Z}[\lambda]$ with $b > 0$,
- (ii) there exist $c, d \in \mathbb{Z}[\lambda]$ such that $\begin{pmatrix} a & c \\ b & d \end{pmatrix} \in G$.

Let $x/y = (x_1 + x_2\lambda)/(y_1 + y_2\lambda) > 0$ be a reduced form. Since $\lambda^2 = \lambda + 1$, we may assume that x_1, x_2, y_1 , and y_2 are nonnegative rational integers ([LLT1]). The following lemmas must be well known among the experts. For the reader's convenience, the proof is included.

Lemma 1. *Let $x/y > 0$ ($y > 0$) be a reduced form. Then $y \in \mathbb{N}$ if and only if $y = 1$.*

Proof. Let a/b and c/d ($0 \leq a/b < c/d$) be reduced forms such that $(a/b, c/d)$ is an even line. The vertices of the ideal pentagon lying below $(a/b, c/d)$ are given by

$$\{a/b, (\lambda a + c)/(\lambda b + d), \lambda(a + c)/\lambda(b + d), (a + \lambda c)/(b + \lambda d), c/d\}.$$

Since x/y is a reduced form, x/y is a vertex of an ideal pentagon I . Suppose that I lies below $(a/b, c/d)$. As $x/y \notin \{0, \infty\}$, we may assume that

$$x/y \in \{(\lambda a + c)/(\lambda b + d), \lambda(a + c)/\lambda(b + d), (a + \lambda c)/(b + \lambda d)\}.$$

Let $a = a_1 + a_2\lambda$, $b = b_1 + b_2\lambda$, $c = c_1 + c_2\lambda$, and $d = d_1 + d_2\lambda$. Note that we may assume that $a_1, a_2, b_1, b_2, c_1, c_2, d_1$, and d_2 are nonnegative rational integers.

Suppose that $y \in \mathbb{N}$.

Case 1. $x/y = (\lambda a + c)/(\lambda b + d)$. It follows that $b_1 + b_2 + d_2 = 0$ and $b_2 + d_1 = y$. Hence $b_1 = b_2 = d_2 = 0$. As a consequence, $b = 0$ and $d = y$. This is a contradiction ($0 \leq a/b < c/d$).

Case 2. $x/y = (a + \lambda c)/(b + \lambda d)$. It follows that $b_2 + d_1 + d_2 = 0$ and $b_1 + d_2 = y$. Hence $b_2 = d_1 = d_2 = 0$. As a consequence, $b = y$, $d = 0$, and $(a/b, c/d) = (a/y, 1/0)$. By Proposition 6 (iii) of [LLT1], $y = 1$.

Case 3. $x/y = \lambda(a + c)/\lambda(b + d)$. It follows that $b_1 + b_2 + d_1 + d_2 = 0$ and $b_2 + d_2 = y$ is an integer. It follows that $b_1 = b_2 = d_1 = d_2 = 0$. This is a contradiction.

This completes the proof of the lemma. $\square$

Lemma 2. *Let m be a positive rational integer. Then the only reduced forms between 0 and λ with denominator $m\lambda$ are $1/m\lambda$ and $\lambda - 1/m\lambda = (m\lambda^2 - 1)/m\lambda = (m\lambda + (m - 1))/m\lambda$.*

Proof. Let $0 < x/m\lambda < \lambda$ be a reduced form. Similar to Lemma 1, we may assume that

$$x/y \in \{(\lambda a + c)/(\lambda b + d), \lambda(a + c)/\lambda(b + d), (a + \lambda c)/(b + \lambda d)\}$$

for some even line $(a/b, c/d)$. Let $a = a_1 + a_2\lambda$, $b = b_1 + b_2\lambda$, $c = c_1 + c_2\lambda$, and $d = d_1 + d_2\lambda$. Note that we may assume that $a_1, a_2, b_1, b_2, c_1, c_2, d_1$, and d_2 are nonnegative rational integers.

Suppose that $x/m\lambda = (\lambda a + c)/(\lambda b + d)$. It follows that $b_1 + b_2 + d_2 = m$ and $b_2 + d_1 = 0$. Hence $b_2 = d_1 = 0$. This implies that $b = b_1 \in \mathbb{N}$. Since a/b is a reduced form, $b = b_1 = 1$ (Lemma 1). By Corollary 5 of [LLT1], $a = k\lambda$ for some

$k \in \mathbb{Z}$. Since $0 \leq a/b < \lambda$, $a = 0$. It follows that $d = (m-1)\lambda$. By Proposition 6 (iii) of [LLT1], $c = 1$. Hence $x/m\lambda = (\lambda a + c)/(\lambda b + d) = 1/m\lambda$.

Suppose that $x/m\lambda = (a + \lambda c)/(\lambda b + d)$. It follows that $b_2 + d_1 + d_2 = m$ and $b_1 + d_2 = 0$. Hence $b_1 = d_2 = 0$. This implies that $d = d_1 \in \mathbb{N} \cup \{0\}$. Since c/d is a reduced form, $d = d_1 = 1$ (Lemma 1). Applying Corollary 5 of [LLT1], $c = 0$ or $k\lambda$, where $k \in \mathbb{Z}$. Since $0 \leq a/b < c/d \leq \lambda$, $c = \lambda$. Since $c/d = \lambda/1$, $b = b_2 = (m-1)\lambda$. By Proposition 6 (iii) of [LLT1], $a = -1 + (m-1)\lambda^2$. Hence $x/m\lambda = (\lambda a + c)/(\lambda b + d) = (m\lambda^2 - 1)/m\lambda$.

Suppose that $x/m\lambda = \lambda(a+c)/\lambda(b+d)$. It follows that $b_2 = d_2 = 0$. This implies that $b = b_1 \in \mathbb{N}$, $d = d_1 \in \mathbb{N} \cup \{0\}$, and $b_1 + d_1 = m$. By Lemma 1, $b = b_1 = 1$ and $d = d_1 = 0$. It follows that $m = 1$. Since $1/\lambda$ and $\lambda/\lambda = \lambda - 1/\lambda$ are the only reduced forms between 0 and λ with denominator λ , the lemma holds.

In summary, the only reduced forms between 0 and λ with denominator $m\lambda$ are $1/m\lambda$ and $(m\lambda^2 - 1)/m\lambda$. $\square$

Applying Lemma 2, we have the following.

Corollary 3. *Let m be a positive rational integer. Then $x/m\lambda$ is a reduced form if and only if $x = km\lambda^2 \pm 1$, where $k \in \mathbb{Z}$. Further, if $x/m\lambda$ and $y/m\lambda$ are reduced forms such that $x/m\lambda - y/m\lambda = 2/m\lambda$, then $x/m\lambda = k\lambda + 1/m\lambda$ and $y/m\lambda = k\lambda - 1/m\lambda$.*

Proof. By Lemma 2, the reduced forms with denominator $m\lambda$ are given by $k\lambda \pm (1/m\lambda)$. This completes the proof of the corollary. $\square$

3. PSEUDO-EUCLIDEAN ALGORITHM AND REDUCED FORMS

Let $a, b \in \mathbb{Z}[\lambda] \setminus \{0\}$. There exists a unique rational integer q such that

- (i) $a = (q\lambda)b + r$,
- (ii) $|r| = |a - (q\lambda)b| \leq |a - (x\lambda)b|$ for all $x \in \mathbb{Z}$.

We call such a division algorithm *pseudo-Euclidean* (see [R] for more details). In terms of matrices, the above can be written as

$$\begin{pmatrix} 1 & -q\lambda \\ 0 & 1 \end{pmatrix} \begin{pmatrix} a \\ b \end{pmatrix} = \begin{pmatrix} r \\ b \end{pmatrix}.$$

Note that $\begin{pmatrix} 1 & -q\lambda \\ 0 & 1 \end{pmatrix} \in G$. Let $n \in \mathbb{N}$. Applying the pseudo-Euclidean algorithm repeatedly, one has,

$$n = (q_1\lambda)1 + r_1,$$

$$1 = (q_2\lambda)r_1 + r_2,$$

$$r_1 = (q_3\lambda)r_2 + r_3,$$

.....

.....

.....

$$r_k = (q_{k+1}\lambda)r_{k+1} + r_{k+2},$$

$$r_{k+1} = (q_{k+2}\lambda)r_{k+2} + 0.$$

The finiteness of the algorithm is governed by the fact that the set of cusps of G is $\mathbb{Q}(\lambda) \cup \{\infty\}$. Note that in terms of matrices, the above can be written as

$$(3.1) \quad \begin{pmatrix} n \\ 1 \end{pmatrix} = A \begin{pmatrix} r_{k+2} \\ 0 \end{pmatrix},$$

where $A \in G$. It is clear that

$$\gcd(n, 1) = \gcd(1, r_1) = \cdots = \gcd(r_{k+1}, r_{k+2}) = r_{k+2}$$

is a unit. As

- (i) $|r_{k+2}| < 1$,
- (ii) λ is a primitive unit,

there exists $e(n) \in \mathbb{N}$ such that

$$|r_{k+2}| = \lambda^{-e(n)}.$$

Multiplying (3.1) by $\lambda^{e(n)}$, one has

$$\begin{pmatrix} n\lambda^{e(n)} \\ \lambda^{e(n)} \end{pmatrix} = A \begin{pmatrix} \pm 1 \\ 0 \end{pmatrix}.$$

Since $A \in G$ and $\pm 1/0$ is a reduced form, we conclude that $n\lambda^{e(n)}/\lambda^{e(n)}$ is the reduced form of n . Consequently, $-\lambda^{e(n)}/n\lambda^{e(n)}$ is the reduced form of $-1/n$. Note that since the reduced form is unique, $e(n)$ is the unique rational integer such that

$$-\lambda^{e(n)}/n\lambda^{e(n)} \text{ and } n\lambda^{e(n)}/\lambda^{e(n)}$$

are reduced forms.

Lemma 4. *Let $m \geq 1$ and $x > 1$ be rational integers. Then there exists $f \in \mathbb{N}$ such that $e(mx^{f+1}) > e(mx^f)$.*

Proof. Let $s, t \in \mathbb{N} \cup \{0\}$. Applying the pseudo-Euclidean algorithm, one has

$$mx^s = (q_s\lambda)1 + r_s$$

and

$$mx^t = (q_t\lambda)1 + r_t.$$

Suppose that $r_s = r_t$. It follows that

$$mx^s - mx^t = (q_s - q_t)\lambda.$$

As λ is irrational, the above implies that $mx^s = mx^t$. Consequently, $r_s \neq r_t$ if and only if $s \neq t$. Therefore, mx^s is not congruent to mx^t modulo λ if and only if $s \neq t$. This implies that

$$\Omega = \{mx^s\lambda^{e(mx^s)}/\lambda^{e(mx^s)} \pmod{\lambda} : s \in \mathbb{N} \cup \{0\}\}$$

is infinite. Let

$$mx^s\lambda^{e(mx^s)}/\lambda^{e(mx^s)} = k_s\lambda + mx^s\lambda^{e(mx^s)}/\lambda^{e(mx^s)} \pmod{\lambda},$$

where $k_s \in \mathbb{Z}$. It follows that

$$(3.2) \quad \begin{pmatrix} 1 & -k_s\lambda \\ 0 & 1 \end{pmatrix} \begin{pmatrix} mx^s\lambda^{e(mx^s)} \\ \lambda^{e(mx^s)} \end{pmatrix} = \begin{pmatrix} mx^s\lambda^{e(mx^s)} - \lambda^{e(mx^s)}k_s\lambda \\ \lambda^{e(mx^s)} \end{pmatrix}.$$

As $mx^s \lambda^{e(mx^s)} / \lambda^{e(mx^s)}$ is in reduced form, (3.2) implies that

$$(mx^s \lambda^{e(mx^s)} - \lambda^{e(mx^s)} k_s \lambda) / \lambda^{e(mx^s)} = mx^s \lambda^{e(mx^s)} / \lambda^{e(mx^s)} \pmod{\lambda}$$

is also in reduced form. Hence every member in Ω is in reduced form. Since

- (i) Ω is infinite,
- (ii) there exist only finitely many cusps u/v (in reduced form) in $[0, \lambda]$ such that $v \leq \lambda^{e(m)}$,

there exists some s such that

$$\lambda^{e(mx^s)} > \lambda^{e(m)}.$$

Hence there exists some $f \in \mathbb{N}$ such that $e(mx^{f+1}) > e(mx^f)$. $\square$

4. FIRST REDUCTION

Throughout this section, (τ) is a nontrivial ideal of $\mathbb{Z}[\lambda]$ and m is the smallest positive rational integer in (τ) . Let $PGL_2^+(\mathbb{Q}(\lambda)) = \{A \in PGL_2(\mathbb{Q}(\lambda)) : \det A > 0\}$. A matrix M in $PGL_2^+(\mathbb{Q}(\lambda))$ is said to be primitive if

- (i) all the entries of M are members of $\mathbb{Z}[\lambda]$,
- (ii) the greatest common divisor of the entries of M is 1.

Denote by $N(G_0(\tau))$ the normalizer of $G_0(\tau)$ in $PSL_2(\mathbb{R})$. The purpose of this section is to show that every element in $N(G_0(\tau))$ is of the form $\pi A'$, where $\pi \in \mathbb{Q}(\lambda)^{1/2} = \{z \in \mathbb{R} : z^2 \in \mathbb{Q}(\lambda)\}$ and $A' \in PGL_2^+(\mathbb{Q}(\lambda))$ is primitive.

For any

$$A = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in N(G_0(\tau)),$$

we have

$$(4.1) \quad \begin{pmatrix} a & b \\ c & d \end{pmatrix} \begin{pmatrix} 1 & \lambda \\ 0 & 1 \end{pmatrix} \begin{pmatrix} a & b \\ c & d \end{pmatrix}^{-1} = \begin{pmatrix} 1 - ac\lambda & a^2\lambda \\ -c^2\lambda & 1 + ac\lambda \end{pmatrix} \in G_0(\tau),$$

$$(4.2) \quad \begin{pmatrix} a & b \\ c & d \end{pmatrix}^{-1} \begin{pmatrix} 1 & \lambda \\ 0 & 1 \end{pmatrix} \begin{pmatrix} a & b \\ c & d \end{pmatrix} = \begin{pmatrix} 1 + dc\lambda & d^2\lambda \\ -c^2\lambda & 1 - dc\lambda \end{pmatrix} \in G_0(\tau),$$

$$(4.3) \quad \begin{pmatrix} a & b \\ c & d \end{pmatrix} \begin{pmatrix} 1 & 0 \\ m\lambda & 1 \end{pmatrix} \begin{pmatrix} a & b \\ c & d \end{pmatrix}^{-1} = \begin{pmatrix} 1 + bdm\lambda & -b^2m\lambda \\ d^2m\lambda & 1 - bdm\lambda \end{pmatrix} \in G_0(\tau),$$

$$(4.4) \quad \begin{pmatrix} a & b \\ c & d \end{pmatrix}^{-1} \begin{pmatrix} 1 & 0 \\ m\lambda & 1 \end{pmatrix} \begin{pmatrix} a & b \\ c & d \end{pmatrix} = \begin{pmatrix} 1 - abm\lambda & -b^2m\lambda \\ a^2m\lambda & 1 + abm\lambda \end{pmatrix} \in G_0(\tau).$$

Case 1. Suppose $c = 0$. By (4.1), we have $a = \pm\sqrt{n_2}$ and $d = \pm 1/\sqrt{n_2}$ ($ad = 1$), where $n_2 \in \mathbb{Z}[\lambda]$. By (4.4), $abm \in \mathbb{Z}[\lambda]$. Hence $b = \pm n_3/m\sqrt{n_2}$, where $n_3 \in \mathbb{Z}[\lambda]$. As a consequence,

$$A = \pm \begin{pmatrix} \sqrt{n_2} & n_3/m\sqrt{n_2} \\ 0 & 1/\sqrt{n_2} \end{pmatrix}.$$

It follows that $A = \pi A'$, where $\pi \in \mathbb{Q}(\lambda)^{1/2} = \{z \in \mathbb{R} : z^2 \in \mathbb{Q}(\lambda)\}$ and $A' \in PGL_2^+(\mathbb{Q}(\lambda))$ is primitive.

Case 2. Suppose $c \neq 0$. By (4.1), $c = \pm\sqrt{\tau n_1}$, where $n_1 \in \mathbb{Z}[\lambda]$. Since $1 - ac\lambda \in \mathbb{Z}[\lambda]$, $a = \pm n_3/\sqrt{\tau n_1}$, where $n_3 \in \mathbb{Z}[\lambda]$. By (4.2), $1 - dc\lambda \in \mathbb{Z}[\lambda]$. Hence $d = \pm n_4/\sqrt{\tau n_1}$, where $n_4 \in \mathbb{Z}[\lambda]$. Since the determinant of A is 1, we conclude that

$$b = \pm(n_3 n_4 / \tau n_1 - 1) / \sqrt{\tau n_1}.$$

Similar to Case 1, $A = \pi A'$, where $\pi \in \mathbb{Q}(\lambda)^{1/2} = \{z \in \mathbb{R} : z^2 \in \mathbb{Q}(\lambda)\}$ and $A' \in PGL_2^+(\mathbb{Q}(\lambda))$ is primitive.

Remark. Applying the proof we presented in this section, one can show that every member of the normalizer of $G_0(\tau)$ of G_q , $q \geq 4$, is of the form $A = \pi A'$, where $\pi \in \mathbb{Q}(\lambda)^{1/2} = \{z \in \mathbb{R} : z^2 \in \mathbb{Q}(\lambda)\}$ and $A' \in PGL_2^+(\mathbb{Q}(\lambda))$ is primitive.

5. SECOND REDUCTION: $N(G_0(\tau)) \leq G_5$

Throughout this section, m is the smallest positive rational integer in the nontrivial ideal (τ) and $A = \pi A' \in N(G_0(\tau))$, where $\pi \in \mathbb{Q}(\lambda)^{1/2}$ and $A' \in PGL_2^+(\mathbb{Q}(\lambda))$ is primitive. The main purpose of this section is to show that A is a member of $G = G_5$.

For each

$$A = \pi A' = \pi \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in N(G_0(\tau)),$$

let a'/c' be the reduced form of a/c . Proposition 6 of [LLT1] implies that G contains an element of the form

$$B^{-1} = \begin{pmatrix} a' & b' \\ c' & d' \end{pmatrix}.$$

It follows that

$$u(A) = BA = \pi BA' = \pi \begin{pmatrix} x & y \\ 0 & z \end{pmatrix},$$

where $x, y, z \in \mathbb{Z}[\lambda]$. Note that since $B \in G$ and A' is primitive, $\gcd(x, y, z) = 1$. Furthermore, since $B \in G$ and $A \in N(G_0(\tau))$,

$$(5.1) \quad u(A)\sigma u(A)^{-1} = BA\sigma(BA)^{-1} = \begin{pmatrix} r & s \\ t & u \end{pmatrix} \in G$$

for every σ in $G_0(\tau)$. In particular, r/t and s/u are in reduced form.

Lemma 5. x/z is a rational integer and m is a multiple of x/z .

Proof. Let $\sigma = \begin{pmatrix} 1 & \lambda \\ 0 & 1 \end{pmatrix}$. It follows by applying (5.1) that

$$u(A)\sigma u(A)^{-1} = \begin{pmatrix} 1 & x\lambda/z \\ 0 & 1 \end{pmatrix} \in G.$$

By Corollary 5 of [LLT1], $x/z \in \mathbb{Z}$. This completes the proof of the first part of the lemma. Let $\sigma = \begin{pmatrix} 1 & 0 \\ m\lambda & 1 \end{pmatrix}$. It follows that

$$(5.2) \quad u(A)\sigma u(A)^{-1} = \begin{pmatrix} 1 + my\lambda/x & -my^2\lambda/xz \\ mz\lambda/x & 1 - my\lambda/x \end{pmatrix} \in G.$$

Hence $mz\lambda/x \in \mathbb{Z}[\lambda]$. Since

- (i) $mz\lambda/x \in \mathbb{Z}[\lambda]$,
- (ii) both x/z and m are rational integers,

m is a multiple of x/z . This completes the proof of the lemma. $\square$

Lemma 6. $z = \pm 1$.

Proof. Replacing x by $-x$ if necessary, we may assume that $mz/x > 0$. By (5.2), $(1 + my\lambda/x)/(mz\lambda/x)$ and $(-1 + my\lambda/x)/(mz\lambda/x)$ are reduced forms (recall that the denominator of a reduced form is nonnegative). Note that

$$(1 + my\lambda/x)/(mz\lambda/x) - (-1 + my\lambda/x)/(mz\lambda/x) = 2/(mz\lambda/x).$$

By Corollary 3, we have

- (i) $(1 + my\lambda/x)/(mz\lambda/x) = k\lambda + 1/(mz\lambda/x)$,
- (ii) $(-1 + my\lambda/x)/(mz\lambda/x) = k\lambda - 1/(mz\lambda/x)$,

where $k \in \mathbb{Z}$. Hence $y = kz\lambda$. Using Lemma 5, we conclude that both x and y are multiples of z . Since $\gcd(x, y, z) = 1$, the above implies that $z = \pm 1$. $\square$

Applying Lemmas 5 and 6 to $u(A)$, we have (replace B by $-B$ if necessary)

$$u(A) = BA = \pi BA' = \pi \begin{pmatrix} x & k\lambda \\ 0 & 1 \end{pmatrix},$$

where $x, k \in \mathbb{Z}$. Multiplying $u(A)$ by

$$\begin{pmatrix} 1 & -k\lambda \\ 0 & 1 \end{pmatrix} \in G,$$

one has

$$d(A) = \begin{pmatrix} 1 & -k\lambda \\ 0 & 1 \end{pmatrix} u(A) = \pi \begin{pmatrix} x & 0 \\ 0 & 1 \end{pmatrix}.$$

Note that since $A \in N(G_0(\tau))$,

$$(5.3) \quad d(A)\sigma d(A)^{-1} = \begin{pmatrix} r' & s' \\ t' & u' \end{pmatrix} \in G$$

for every σ in $G_0(\tau)$. In particular, r'/t' and s'/u' are reduced forms.

Lemma 7. Let (τ) be a nontrivial ideal of $\mathbb{Z}[\lambda]$. Then $N(G_0(\tau)) \leq G = G_5$.

Proof. Let $A = \pi A' \in N(G_0(\tau))$, where $\pi \in \mathbb{Q}(\lambda)^{1/2}$ and $A' \in PGL_2^+(\mathbb{Q}(\lambda))$ is primitive. Since

$$(i) \quad d(A) = \pi \begin{pmatrix} x & 0 \\ 0 & 1 \end{pmatrix},$$

- (ii) A and $d(A)$ have the same determinant,

$\det A = \pi^2 x$. It follows that $\det A' = x$. Since $x/z \in \mathbb{Z}$ (Lemma 5) and $z = \pm 1$ (Lemma 6), $x \in \mathbb{Z}$. As members in $PGL_2^+(\mathbb{Q}(\lambda))$ admit positive determinant, $\det A' = x \in \mathbb{N}$. Suppose that $x \geq 2$. By Lemma 4, there exists some f such that $e(mx^{f+1}) > e(mx^f)$ (m is the smallest positive rational integer in (τ)). Let $n = x^{f+1}$. Since $-\lambda^{e(mn)}/\lambda^{e(mn)}mn$ is in reduced form (Section 3), by Proposition 6 of [LLT1], $G_0(\tau)$ has an element of the form

$$\sigma = \begin{pmatrix} -\lambda^{e(mn)} & * \\ \lambda^{e(mn)}mn & * \end{pmatrix} \in G_0(\tau).$$

By (5.3),

$$d(A)\sigma d(A)^{-1} = \begin{pmatrix} -\lambda^{e(mn)} & * \\ \lambda^{e(mn)}mn/x & * \end{pmatrix} \in G.$$

It follows that

$$-\lambda^{e(mn)}/(\lambda^{e(mn)}mn/x)$$

is in reduced form. This is a contradiction ($e(mn) = e(mx^{f+1}) > e(mx^f) = e(mn/x)$). Hence $x = 1$. As a consequence,

$$A' = B^{-1} \begin{pmatrix} 1 & k\lambda \\ 0 & 1 \end{pmatrix} \in G.$$

Since $1 = \det A = \pi^2 \det A' = \pi^2$ and $\pi \in \mathbb{Q}(\lambda)^{1/2}$, we conclude that $\pi = \pm 1$. Hence $A = \pm A'$. Since $A' \in G$, $A = \pm A' \in G = G_5$. This implies that $N(G_0(\tau)) \leq G = G_5$. $\square$

6. THE MAIN THEOREM

Theorem 8. *Let (τ) be an ideal of $\mathbb{Z}[\lambda]$ and let (τ_0) be the maximal ideal of $\mathbb{Z}[\lambda]$ such that $(\tau_0^2) \subseteq (\tau)$. Then $N(G_0(\tau)) \leq G_0(\tau_0)$.*

Proof. Let $A \in N(G_0(\tau))$. By Lemma 7, $A \in G$. The theorem now follows by applying (4.1). $\square$

Main Theorem. *Let (τ) be an ideal of $\mathbb{Z}[\lambda]$. Suppose that τ is square free. Then $G_0(\tau)$ is self-normalized in $PSL_2(\mathbb{R})$.*

Proof. Since τ is square free, $(\tau) = (\tau_0)$ in Theorem 8. This completes the proof of the theorem. $\square$

REFERENCES

- [AL] A. O. L. Atkin, J. Lehner, *Hecke operators on $\Gamma_0(m)$* , Math. Ann. 185, (1970), 134-160. MR **42**:3022
- [C] J. H. Conway, *Understanding Groups like $\Gamma_0(N)$* , Groups, difference sets and the monster (Columbus, Ohio, 1993), Ohio State Univ. Math. Res. Inst. Publ., 4, de Gruyter, Berlin, 1996, 327-343. MR **98b**:11041
- [CLLT] S. P. Chan, M. L. Lang, C. H. Lim, S. P. Tan, *The invariants of the congruence subgroups $G_0(P)$ of the Hecke group*, Illinois J. of Math. 38 (1994), 636-652. MR **95d**:11047
- [K] R. S. Kulkarni, *An Arithmetic-Geometric method in the study of subgroups of the Modular Groups*, Amer. Jour. Math. 113 (1991), 1053-1134. MR **92i**:11046
- [L1] A. Leutbecher, *Über die Heckschen Gruppen $G(\lambda)$* , Abh. Math. Sem. Hambg. 31 (1967), 199-205. MR **37**:4018
- [L2] A. Leutbecher, *Über die Heckschen Gruppen $G(\lambda)$, II*, Math. Ann. 211 (1974), 63-68. MR **50**:238
- [LT1] M. L. Lang, S. P. Tan, *Normalizer of the congruence subgroups of the Hecke groups G_4 and G_6* , preprint.
- [LT2] M. L. Lang, S. P. Tan, *Normalizer of the congruence subgroups of the Hecke group G_5* , Proc. Amer. Math. Soc. 127 (1999), 3131-3140. CMP 99:17
- [LLT1] M. L. Lang, C. H. Lim, S. P. Tan, *Independent generators for congruence subgroups of Hecke groups*, Math. Z. 220 (1995), 569-594. MR **96k**:11049
- [LLT2] M. L. Lang, C. H. Lim, S. P. Tan, *Principal congruence subgroups of the Hecke groups*, (submitted for publication).
- [LN] J. Lehner, M. Newman, *Weierstrass Points of $\Gamma_0(N)$* , Annals of Math. 79 (1964), 360-368. MR **28**:5045

- [P] L.A. Parson, *Generalized Kloosterman sums and the Fourier coefficients of cusp forms*, Trans. Amer. Math. Soc. 217 (1976), 329-350. MR **54**:241
- [R] D. Rosen, *A class of continued fractions associated with certain properly discontinuous groups*, Duke Math., 21(1954), 549-563. MR **16**:458d

DEPARTMENT OF MATHEMATICS, NATIONAL UNIVERSITY OF SINGAPORE, SINGAPORE 119260,
REPUBLIC OF SINGAPORE

E-mail address: `matlml@math.nus.edu.sg`

E-mail address: `mattansp@nus.edu.sg`