CHERN-OSSERMAN INEQUALITY FOR MINIMAL SURFACES IN $\mathbf{H}^n$

CHEN QING AND CHENG YI

(Communicated by Christopher Croke)

ABSTRACT. We obtain Chern-Osserman's inequality of a complete properly immersed minimal surface in hyperbolic n -space, provided the L^2 -norm of the second fundamental form of the surface is finite.

1. INTRODUCTION

Let M be a complete minimal surface in Euclidean space $\mathbf{R}^n$ with finite total Gaussian curvature. Then the total Gaussian curvature of M satisfies the Chern-Osserman inequality ([2], [6])

$$-\chi(M) \leq \frac{-1}{2\pi} \int_M K - k,$$

where K is the Gaussian curvature of M , $\chi(M)$ is the Euler characteristic of M and k is the number of ends of M . The explicit expression of the total Gaussian curvature was obtained by Jorge and Meeks:

$$\begin{aligned} -\chi(M) &= \frac{-1}{2\pi} \int_M K - \sup \frac{\text{area} M \cap B(t)}{\pi t^2} \\ &= \frac{1}{4\pi} \int_M |A|^2 - \sup \frac{\text{area} M \cap B(t)}{\pi t^2}, \end{aligned}$$

where A is the second fundamental form of M and $B(t)$ is the extrinsic distance ball of radius t from a fixed point.

In the paper we present an analogue of the Chern-Osserman inequality of complete minimal surfaces in hyperbolic n -space $\mathbf{H}^n$ of constant curvature -1 , namely

Theorem. *Let M be an oriented immersed complete minimal surface in $\mathbf{H}^n$, A the second fundamental form of M , r the distance of $\mathbf{H}^n$ from a fixed point and $M_t = \{x \in M : r(x) < t\}$. Suppose $\int_M |A|^2(x) < \infty$; then*

$$(1) \sup \frac{\text{area} M_t}{\cosh t - 1} < +\infty;$$

$$(2) -\chi(M) \leq \frac{1}{4\pi} \int_M |A|^2 - \sup \frac{\text{area}(M_t)}{2\pi(\cosh t - 1)},$$

where $\chi(M)$ is the Euler characteristic of M . Consequently, M has finite topological type.

Received by the editors September 14, 1998.

1991 *Mathematics Subject Classification.* Primary 53A20; Secondary 53C42.

Key words and phrases. Minimal surface, Chern-Osserman inequality, Euler characteristic.

The minimal surfaces in $\mathbf{H}^n$ have some properties similar to minimal surfaces in $\mathbf{R}^n$, such as the monotonicity formula (see Proposition 2.2 below). But there are many differences between minimal surfaces in $\mathbf{H}^n$ and those in $\mathbf{R}^n$, one of the differences is that the total Gaussian curvature of a complete minimal surface in $\mathbf{H}^n$ is always infinite (this can be seen from the correspondent Gauss equation). Another important difference comes from the fundamental work of M. T. Anderson [1] for the existence of minimal varieties in $\mathbf{H}^n$. By his results, minimal surfaces in the hyperbolic space are much richer than those in Euclidean space, and the asymptotic behavior of the surfaces is not “regular” in general; in particular, the Bernstein Theorem (a complete minimal graph in $\mathbf{R}^3$ is flat) does not hold in $\mathbf{H}^3$.

In [5] De Oliveira proved that if M is an immersed complete minimal surface in $\mathbf{H}^n$ with $\int_M |A|^2 < \infty$, then M is properly immersed and is conformally equivalent to a compact surface with a finite number of disks removed. M. Kokubu [4] established the Weierstrass type representation formula for minimal surfaces in hyperbolic space. By his result, the Gauss maps of minimal surfaces in hyperbolic space are neither holomorphic nor anti-holomorphic. So the method employed in [2] and [6] is not valid in our case.

2. PRELIMINARIES

Let $\mathbf{H}^n$ be a hyperbolic n -space of constant curvature -1 , and M a properly immersed complete minimal surface in $\mathbf{H}^n$. Denote the covariant derivative of $\mathbf{H}^n$ and M by D and ∇ respectively; the second fundamental form of M is defined by

$$\begin{aligned} A : TM \otimes TM &\rightarrow T^\perp M, \\ (2.1) \quad A(X, Y) &= D_X Y - \nabla_X Y, \text{ for } X, Y \in C^\infty(TM). \end{aligned}$$

For a smooth function f on $\mathbf{H}^n$, and any two tangent vector fields $X, Y \in C^\infty(TM)$,

$$\begin{aligned} (D^2 f)(X, Y) &= (Ddf)(X, Y) \\ &= X(df(Y)) - df(D_X Y) \\ &= X(df(Y)) - df(\nabla_X Y + A(X, Y)) \\ &= \nabla^2 f(X, Y) - \langle A(X, Y), Df \rangle, \end{aligned}$$

which, together with the fact that $(D^2 r)(X, X) = \coth r(\langle X, X \rangle - \langle X, Dr \rangle^2)$, implies

Proposition 2.1. *For any unit tangent vector e of M ,*

$$(\nabla^2 r)(e, e) = \coth r(1 - \langle e, \nabla r \rangle^2) + \langle A(e, e), \nabla^\perp r \rangle,$$

where $\nabla^\perp r$ is the projection of Dr onto the normal of M .

Let r be the distance function of H^n from a fixed point. By Sard's theorem, for almost all $t > 0$, $M_t = \{x \in M : r(x) < t\}$ is a related compact open subset of M with the boundary ∂M_t being a closed immersed curve of M . $\{M_t\}$ is a family of exhaustion of M . We will consider the growth of the area of M_t , and make use of the following notations for convenience:

$$v(t) = \text{area} M_t, \quad \text{and} \quad R(t) = \int_{M_t} |A|^2.$$

Proposition 2.2 ([1], Theorem 1). *$\frac{v(t)}{\cosh t - 1}$ is monotone nondecreasing in t , i.e., $v'(t) \cosh t - v(t) \sinh t \geq v'(t)$.*

Proposition 2.3. *Suppose M is an oriented and properly immersed complete minimal surface in $\mathbf{H}^n$. Then for almost all $t > 0$,*

$$v(t) + \frac{1}{2}R(t) + 2\pi\chi(M_t) = v'(t) \coth t - \int_{\partial M_t} \left\langle A\left(\frac{\nabla r}{|\nabla r|}, \frac{\nabla r}{|\nabla r|}\right), \frac{\nabla^\perp r}{|\nabla r|} \right\rangle.$$

Proof. Denote the geodesic curvature of ∂M_t in M by k_g^t , and by K the Gaussian curvature of M . Then the Gauss-Bonnet formula reads

$$(2.2) \quad \int_{\partial M_t} k_g^t + \int_{M_t} K = 2\pi\chi(M_t),$$

where $\chi(M_t)$ is the Euler characteristic of M_t , i.e. $\chi(M_t) = 2 - 2g - k$ with g being the genus of M_t and k being the number of components of ∂M_t .

Substituting the Gauss equation $K = -1 - \frac{1}{2}|A|^2$ into (2.2) we have

$$(2.3) \quad v(t) + \frac{1}{2}R(t) + 2\pi\chi(M_t) = \int_{\partial M_t} k_g^t.$$

Suppose e is the unit tangent vector of ∂M_t . Since the normal of ∂M_t in M is $\frac{\nabla r}{|\nabla r|}$,

$$\begin{aligned} k_g^t &= -\left\langle \nabla_e e, \frac{\nabla r}{|\nabla r|} \right\rangle \\ (2.4) \quad &= \frac{1}{|\nabla r|} (\nabla^2 r)(e, e) \\ &= \frac{1}{|\nabla r|} (\coth r + \langle A(e, e), \nabla^\perp r \rangle), \end{aligned}$$

where the last equality follows by Proposition 2.1.

Substituting (2.4) into (2.3), and using the fact that $v'(t) = \int_{\partial M_t} \frac{1}{|\nabla r|}$, and that $A(e, e) + A\left(\frac{\nabla r}{|\nabla r|}, \frac{\nabla r}{|\nabla r|}\right) = 0$, we complete the proof.

Lemma 2.4. *For $t > s > 0$,*

$$\frac{\int_{M_t} \cosh r}{\cosh^2 t} - \frac{\int_{M_s} \cosh r}{\cosh^2 s} = \int_{M_t - M(s)} \frac{1 + |\nabla^\perp r|^2 \sinh^2 r}{\cosh^3 r}.$$

Proof. By the minimality of M and Proposition 2.1, we observe that

$$\Delta r = \coth r (2 - |\nabla r|^2),$$

where Δ is the Laplacian of M . This yields

$$(2.5) \quad \Delta \cosh r = 2 \cosh r.$$

Integrating (2.2) over M_t , and using Green's formula,

$$(2.6) \quad 2 \int_{M_t} \cosh r = \int_{\partial M_t} |\nabla r| \sinh r.$$

By using the co-area formula ([7]), we have

$$\begin{aligned}
 \frac{d}{dt} \left(\frac{\int_{M_t} \cosh r}{\cosh^2 t} \right) &= \frac{1}{\cosh^3 t} \left(\cosh t \frac{d}{dt} \int_{M_t} \cosh r - 2 \sinh t \int_{M_t} \cosh r \right) \\
 &= \frac{1}{\cosh^3 t} \left(\cosh t \int_{\partial M_t} \frac{\cosh r}{|\nabla r|} - \sinh t \int_{\partial M_t} |\nabla r| \sinh r \right) \\
 (2.4) \quad &= \frac{1}{\cosh^3 t} \int_{\partial M_t} \left(\frac{\cosh^2 r}{|\nabla r|} - \sinh^2 r |\nabla r| \right) \\
 &= \frac{1}{\cosh^3 t} \int_{\partial M_t} \frac{1}{|\nabla r|} (1 + \sinh^2 r |\nabla^\perp r|^2).
 \end{aligned}$$

The proposition is then proved by integrating (2.6) from s to t and the co-area formula.

Lemma 2.5.

$$\int_0^t v'(s) \cosh s ds \geq \frac{\cosh t + 1}{2} v(t).$$

Proof. By Proposition 2.2, $v'(t) \geq \frac{v(t) \sinh t}{\cosh t - 1}$, so we have

$$\begin{aligned}
 \int_0^t v'(s) \cosh s ds &= v(t) \cosh t - \int_0^t v(s) \sinh s ds \\
 &\geq v(t) \cosh t - \int_0^t v'(s) (\cosh s - 1) ds \\
 &= v(t) (\cosh t + 1) - \int_0^t v'(s) \cosh s ds,
 \end{aligned}$$

which proves the lemma.

3. PROOF OF THE THEOREM

Suppose M is as in the Theorem. By the result of De Oliveira [5], M is properly immersed.

(1) By Proposition 2.3 we have

$$(3.1) \quad \frac{v'(t) \cosh t - v(t) \sinh t}{\sinh t} = \frac{1}{2} R(t) + 2\pi \chi(M_t) + \int_{\partial M_t} \langle A \left(\frac{\nabla r}{|\nabla r|}, \frac{\nabla r}{|\nabla r|} \right), \frac{\nabla^\perp r}{|\nabla r|} \rangle;$$

hence

$$\frac{d}{dt} \frac{v(t)}{\cosh t} \leq \frac{\sinh t}{\cosh^2 t} \left(\frac{1}{2} R(t) + 2\pi \chi(M_t) \right) + \int_{\partial M_t} \frac{|A|}{|\nabla r|} \frac{|\nabla^\perp r| \sinh t}{\cosh^2 t}.$$

Since $\chi(M_t) \leq 1$, integrating the above inequality from 0 to t and by using the co-area formula, we have

$$(3.2) \quad \frac{v(t)}{\cosh t} \leq 2 \int_0^t \left(\frac{1}{2} R(s) + 2\pi \right) e^{-s} ds + \int_{M_t} |A| \frac{|\nabla^\perp r| \sinh r}{\cosh^2 r}.$$

By using the Cauchy-Schwarz inequality,

$$\begin{aligned}
 \frac{v(t)}{\cosh t} &\leq C_1 + \left(\int_{M_t} \frac{|A|^2}{\cosh r} \right)^{\frac{1}{2}} \left(\int_{M_t} \frac{|\nabla^\perp r|^2 \sinh^2 r}{\cosh^3 r} \right)^{\frac{1}{2}} \\
 (3.3) \quad &\leq C_1 + C_2 \left(\frac{\int_{M_t} \cosh r}{\cosh^2 t} \right)^{\frac{1}{2}} \quad (\text{by Proposition 2.3}) \\
 &\leq C_1 + C_2 \left(\frac{v(t)}{\cosh t} \right)^{\frac{1}{2}},
 \end{aligned}$$

where C_1 and C_2 are two constants independent of t . By Proposition 2.2, $\frac{v(t)}{\cosh t}$ is monotone non-decreasing in t ; therefore, either $\sup \frac{v(t)}{\cosh t} \leq C_1^2$ or

$$\frac{v(t)}{\cosh t} \leq \left(\frac{v(t)}{\cosh t} \right)^{\frac{1}{2}} + C_2 \left(\frac{v(t)}{\cosh t} \right)^{\frac{1}{2}}$$

when t is large enough. It follows that

$$\sup \frac{v(t)}{\cosh t} \leq \max\{C_1^2, (1 + C_2)^2\}.$$

This proves (1) of the Theorem.

(2) By the arithmetic geometric mean inequality, (3.1) implies

$$\begin{aligned}
 (3.4) \quad \frac{v'(t) \cosh t - v(t) \sinh t}{\sinh t} &\leq \frac{1}{2} R(t) + 2\pi \chi(M_t) + \frac{1}{2} \int_{\partial M_t} \left(\frac{|A|^2}{|\nabla r|} + \frac{|\nabla^\perp r|^2}{|\nabla r|} \right) \\
 &\leq \frac{1}{2} R(t) + 2\pi \chi(M_t) + \frac{1}{2} R'(t) + \int_{\partial M_t} \frac{|\nabla^\perp r|^2}{|\nabla r|}.
 \end{aligned}$$

Since by Proposition 2.1, $\Delta \cosh r = 2 \cosh r$, Green's formula gives

$$\int_{\partial M_t} |\nabla r| \sinh r = \int_{M_t} 2 \cosh r;$$

then by the co-area formula we have

$$\begin{aligned}
 (3.5) \quad \int_{\partial M_t} \frac{|\nabla^\perp r|^2}{|\nabla r|} &= \int_{\partial M_t} \frac{1}{|\nabla r|} - |\nabla r| \\
 &= v'(t) - \frac{1}{\sinh t} \int_{\partial M_t} |\nabla r| \sinh r \\
 &= v'(t) - \frac{1}{\sinh t} \int_{M_t} 2 \cosh r \\
 &= v'(t) - \frac{2}{\sinh t} \int_0^t v'(s) \cosh s ds.
 \end{aligned}$$

By Lemma 2.5 we obtain

$$(3.6) \quad \int_{\partial M_t} \frac{|\nabla^\perp r|^2}{|\nabla r|} \leq v'(t) - \frac{\cosh t + 1}{\sinh t} v(t).$$

Substituting (3.6) into (3.4) we have

$$\begin{aligned} \frac{v'(t) \cosh t - v(t) \sinh t}{\sinh t} &\leq \frac{1}{2}(R(t) + R'(t)) + 2\pi\chi(M_t) \\ &\quad + \frac{v'(t) \sinh t - v(t) \cosh t - v(t)}{\sinh t}. \end{aligned}$$

This implies

$$\begin{aligned} (3.7) \quad -2\pi\chi(M_t) &\leq \frac{1}{2}(R(t) + R'(t)) + \frac{(v'(t) + v(t))(\sinh t - \cosh t) - v(t)}{\sinh t} \\ &\leq \frac{1}{2}(R(t) + R'(t)) - \frac{v(t)}{\sinh t}. \end{aligned}$$

Since $\int_0^\infty R'(t)dt < \infty$, there is a monotone increasing sequence $\{t_i\}$ diverging to infinity such that $R'(t_i) \rightarrow 0$ as $i \rightarrow \infty$. Taking $t = t_i$ in (3.7) and letting i tend to infinity, we prove the theorem.

ACKNOWLEDGMENT

The authors would like to thank the referee for useful comments.

REFERENCES

1. M. T. Anderson, *Complete minimal varieties in hyperbolic space*, Invent.math. **69** (1982), 477-494. MR **84c**:53005
2. S. S. Chern and R. Osserman, *Complete minimal surface in E^N* , J. d'Analyse Math. **19** (1967), 15-34. MR **37**:2103
3. L. P. Jorge and W. H. Meeks, *The topology of minimal surfaces of finite total Gaussian curvature*, Topology **22** (1983), 203-221. MR **84d**:53006
4. M. Kokubu, *Weierstrass representation for minimal surfaces in hyperbolic space*, Tohoku Math. J. **49** (1997), 367-377. MR **98f**:53008
5. G. De Oliveira, *Compactification of minimal submanifolds of hyperbolic space*, Comm. An. and Geom. **1** (1993), 1-29. MR **94h**:53080
6. R. Osserman, *A survey of minimal surfaces*, Van Norstrand Rienhold, New York, 1969. MR **41**:934
7. L. Simon, *Lectures on Geometric Measure Theory*, C.M.A. Australian National University Vol.3, 1983. MR **87a**:49001

DEPARTMENT OF MATHEMATICS, THE UNIVERSITY OF SCIENCE AND TECHNOLOGY OF CHINA, HEFEI, ANHUI 230026, PEOPLE'S REPUBLIC OF CHINA

E-mail address: qchen@ustc.edu.cn

E-mail address: chengy@ustc.edu.cn