WEAK TYPE ESTIMATES FOR CONE MULTIPLIERS ON H^p SPACES, $p < 1$

SUNGGEUM HONG

(Communicated by Christopher D. Sogge)

ABSTRACT. We consider operators T^δ associated with the Fourier multipliers

$$\left(1 - \frac{|\xi'|^2}{\xi_{n+1}^2}\right)_+^\delta, \quad (\xi', \xi_{n+1}) \in \mathbb{R}^n \times \mathbb{R},$$

and show that T^δ is of weak type (p, p) on $H^p(\mathbb{R}^{n+1})$, $0 < p < 1$, for the critical value $\delta = n(\frac{1}{p} - \frac{1}{2}) - \frac{1}{2}$.

1. INTRODUCTION

We consider the family of Fourier multipliers

$$m^\delta(\xi', \xi_{n+1}) = \left(1 - \frac{|\xi'|^2}{\xi_{n+1}^2}\right)_+^\delta, \quad (\xi', \xi_{n+1}) \in \mathbb{R}^n \times \mathbb{R},$$

with $m^\delta(\xi', \xi_{n+1}) = 0$ when $|\xi'| > \xi_{n+1}$. Let $\widehat{f}$ be the Fourier transform of a Schwartz function f on $\mathbb{R}^n \times \mathbb{R}$. Then define convolution operators T^δ by

$$\widehat{T^\delta f}(\xi', \xi_{n+1}) = m^\delta(\xi', \xi_{n+1}) \widehat{f}(\xi', \xi_{n+1}).$$

It is conjectured that T^δ is bounded on L^p and it remains open for any $n \geq 2$. No optimal L^p -bounds are known for $p > 1$. For partial results, see G. Mockenhaupt [3] and J. Bourgain [1]. See also Mockenhaupt, Seeger and Sogge [4] for the related results on the wave equation. When $\delta > (n-1)/2$, it is not hard to show that T^δ is of weak type $(1, 1)$ by using Calderón-Zygmund theory.

The purpose of this paper is to prove a sharp endpoint result on $H^p(\mathbb{R}^{n+1})$, $p < 1$. This estimate implies the known result due to Stein, Taibleson and Weiss [7] that the Bochner-Riesz means of the critical index $\delta_p = n(1/p - 1/2) - 1/2$ is of weak type (p, p) for functions in $H^p(\mathbb{R}^n)$ (see the Appendix). Here H^p is the standard real Hardy space as defined in [6] by E. Stein.

We prove here

Theorem 1. *Suppose $0 < p < 1$ and $\delta = n(1/p - 1/2) - 1/2$. Then T^δ maps $H^p(\mathbb{R}^{n+1})$ boundedly into weak- $L^p(\mathbb{R}^{n+1})$; i.e., there exists a constant $C = C(n, p)$*

Received by the editors September 24, 1998 and, in revised form, January 28, 1999.

2000 *Mathematics Subject Classification.* Primary 42B15, 42B30.

Key words and phrases. Cone multipliers, atomic decomposition of H^p spaces.

such that for all $f \in H^p(\mathbb{R}^{n+1})$

$$|\{(x, t) \in \mathbb{R}^n \times \mathbb{R} : |T^\delta f(x, t)| > \alpha\}| \leq C \left(\frac{\|f\|_{H^p(\mathbb{R}^{n+1})}}{\alpha} \right)^p$$

for all $\alpha > 0$, where $|E|$ denotes the Lebesgue measure of the set $E \subset \mathbb{R}^{n+1}$.

It is not known whether $T^{\frac{n-1}{2}}$ is of weak type $(1, 1)$ or just of weak type $(1, 1)$ on functions in H^1 .

With respect to the notation, we use C to denote a positive constant, whose value may be different for each occurrence.

2. KERNEL ESTIMATES

Let $\varphi, \psi \in C_0^\infty(\mathbb{R})$ be supported in $(1/2, 2)$ such that $\sum_{k \geq 1} \varphi(2^k s) = 1$ and $\sum_{l=-\infty}^\infty \psi(2^{-l} t) = 1$ for $0 < s < 1$, $t > 0$. We now fix k and l . We shall need pointwise estimates for the kernels of

$$T_{k,l}^\delta f(x, t) = (2\pi)^{-(n+1)} \int_{\mathbb{R}} \int_{\mathbb{R}^n} G_{k,l}(x - y, t - s) f(y, s) dy ds$$

where

$$(2.1) \quad G_{k,l}(x, t) = \int_{\mathbb{R}} \int_{\mathbb{R}^n} \varphi\left(2^k \left(1 - \frac{|\xi'|^2}{\xi_{n+1}^2}\right)\right) \left(1 - \frac{|\xi'|^2}{\xi_{n+1}^2}\right)_+^\delta \psi(2^{-l} \xi_{n+1}) \times e^{i\langle x, \xi' \rangle + it\xi_{n+1}} d\xi' d\xi_{n+1}.$$

For each k and l , the kernel $G_{k,l}$ has the property

$$(2.2) \quad G_{k,l}(\cdot, \cdot) = 2^{l(n+1)} \{G_{k,0}(2^l \cdot, 2^l \cdot)\},$$

and we write $\sum_{k \geq 1} G_{k,l} = G_l = 2^{l(n+1)} G_0(2^l \cdot, 2^l \cdot)$ and $\sum_{k \geq 1} T_{k,l}^\delta = T_l^\delta$.

Lemma 1. Suppose $2^l |x| \leq 2$ for fixed l . Then for each k there is an estimate as follows: for every N

$$(2.3) \quad |G_{k,l}(x, t)| \leq C 2^{l(n+1)} 2^{-k(\delta+1)} \min\{1, (2^{-k} 2^l |t|)^{-N}\}.$$

Thus,

$$(2.4) \quad |G_l(x, t)| \leq C 2^{l(n+1)} \frac{1}{(1 + 2^l |t|)^{\delta+1}}.$$

Proof. In view of (2.1) and (2.2), it suffices to show (2.3) and (2.4) for $l = 0$. We integrate by parts N times with respect to ξ_{n+1} in (2.1). In view of the support property of φ , (2.3) follows. Using (2.3) gives

$$\left\{ C \sum_{|t| \leq 2^k} 2^{-k(\delta+1)} + C \sum_{|t| > 2^k} 2^{-k(\delta+1-N)} |t|^{-N} \right\},$$

and thus (2.4) is established. $\square$

Now we estimate the kernel for the case $2^l |x| > 2$. We let $|x| = r$ and define $G_{k,l}(x, t) = K_{k,l}(|x|, t)$. By Bochner's formula [8] and change of variables, we have

$$(2.5) \quad \begin{aligned} K_{k,l}(r, t) &= 2^{l(n+1)} (2^l r)^{-(n-2)/2} \int_{\mathbb{R}} \int_0^1 J_{\frac{n-2}{2}}(\rho(2^l r)\xi_{n+1}) \varphi(2^k(1 - \rho^2)) \\ &\quad \times (1 - \rho^2)^\delta \psi(\xi_{n+1}) \xi_{n+1}^{(n+2)/2} \rho^{n/2} e^{i(2^l t)\xi_{n+1}} d\rho d\xi_{n+1}. \end{aligned}$$

Here J_μ is the Bessel function of order $\mu > -\frac{1}{2}$ defined by

$$(2.6) \quad J_\mu(t) = A_\mu t^\mu \int_{-1}^1 e^{it\sigma} (1 - \sigma^2)^{\mu - \frac{1}{2}} d\sigma$$

where $A_\mu = [2^\mu \Gamma(2\mu + 1) \Gamma(\frac{1}{2})]^{-1}$.

In order to estimate the kernel (2.5), we need some properties of Bessel functions [8]

$$(2.7) \quad \frac{d}{dt} \{t^{-\mu} J_\mu(t)\} = -t \{t^{-(\mu+1)} J_{\mu+1}(t)\}.$$

For the following lemma, we use dyadic decompositions of Bessel functions (2.6) following the article by Müller and Seeger [5].

Lemma 2. *Suppose that $2^l r > 2$ for fixed l . Then for each k there is an estimate as follows:*

$$(2.8) \quad |K_{k,l}(r, t)| \leq C 2^{l(n+1)} 2^{-k(\delta+1)} (2^l r)^{-(n-1)/2} \min\{1, (2^{-k} 2^l r)^{-N_1}\} \\ \times \left\{ \frac{1}{(1 + 2^l |t + r|)^N} + \frac{1}{(1 + 2^l |t - r|)^N} \right\}.$$

Moreover,

$$(2.9) \quad |K_l(r, t)| \leq C 2^{l(n+1)} (1 + 2^l r)^{-(n+1+2\delta)/2} \left\{ \frac{1}{(1 + 2^l |t + r|)^N} + \frac{1}{(1 + 2^l |t - r|)^N} \right\}.$$

Proof. Let $\eta \in C_0^\infty(\mathbb{R})$ be supported in $(-1/2, 2)$ and equal to 1 in $(-1/4, 1/4)$. Define $m = 0, 1, 2, \dots$ and

$$\eta_{mk}(\sigma, \nu) = \begin{cases} \eta(2^{-k} \nu (1 - \sigma^2)) & \text{if } m = 0, \\ \eta(2^{-k-m} \nu (1 - \sigma^2)) - \eta(2^{-k-m+1} \nu (1 - \sigma^2)) & \text{if } m > 0 \end{cases}$$

and set

$$J_{\mu,k}^m(\rho\nu) = A_\mu(\rho\nu)^\mu \int_{-1}^1 e^{i(\rho\nu)\sigma} (1 - \sigma^2)^{\mu-1/2} \eta_{mk}(\sigma, \nu) d\sigma.$$

Let $M > N + N_1 + (n-1)/2$ and set

$$\phi_{mk\nu}(\sigma) = \begin{cases} (1 - \sigma^2)^{(n-3)/2} \eta_{mk}(\sigma, \nu) & \text{if } m = 0, \\ \left(\frac{1}{i\rho\nu}\right)^M \left(\frac{d}{d\sigma}\right)^M [\eta_{mk}(\sigma, \nu) (1 - \sigma^2)^{(n-3)/2}] & \text{if } m > 0. \end{cases}$$

Then

$$(2.10) \quad J_{\frac{n-2}{2},k}^m(\rho\nu) = A_{\frac{n-2}{2}}(\rho\nu)^{\frac{n-2}{2}} \int_{-1}^1 e^{i(\rho\nu)\sigma} \phi_{mk\nu}(\sigma) d\sigma$$

by integration by parts if $m > 0$.

Fix k and set $l = 0, \nu = r\xi_{n+1}$. We may decompose the kernel (2.5) as

$$K_{k,0} = \sum_{m=0}^{\infty} K_{k,0}^m$$

where

$$\begin{aligned} K_{k,0}^m(r, t) &= r^{-(n-2)/2} \int_{\mathbb{R}} \int_0^1 J_{\frac{n-2}{2},k}^m(\rho\nu) \varphi(2^k(1-\rho^2)) \\ &\quad \times (1-\rho^2)^\delta \psi(\xi_{n+1}) \xi_{n+1}^{(n+2)/2} \rho^{n/2} e^{it\xi_{n+1}} d\rho d\xi_{n+1}. \end{aligned}$$

Formula (2.10) and straightforward computation imply that

$$\begin{aligned} K_{k,0}^m(r, t) &= A_{\frac{n-2}{2}} \int_{-1}^1 \phi_{mk\nu}(\sigma) \int_{\mathbb{R}} \int_0^1 \varphi(2^k(1-\rho^2)) \\ (2.11) \quad &\times (1-\rho^2)^\delta \psi(\xi_{n+1}) \xi_{n+1}^n \rho^{n-1} e^{i(t+\rho r\sigma)\xi_{n+1}} d\rho d\xi_{n+1} d\sigma. \end{aligned}$$

We integrate by parts with respect to ρ and ξ_{n+1} in (2.11) and by Fubini's theorem

$$\begin{aligned} |K_{k,0}^m(r, t)| &\leq C 2^{-k(n-1)/2} \int_{1/2}^2 \int_{-1}^1 \int_0^1 |\phi_{mk\nu}(\sigma)| (1+|\sigma\nu|)^{-N_1} \\ (2.12) \quad &\times (1+|t+\rho r\sigma|)^{-N} \left| \left(\frac{\partial}{\partial \rho} \right)^{N_1} \varphi(2^k(1-\rho^2)) (1-\rho^2)^\delta \rho^{(n-1)/2} \right| \\ &\times \left| \left(\frac{\partial}{\partial \xi_{n+1}} \right)^N \psi(\xi_{n+1}) \xi_{n+1}^n \right| d\rho d\sigma d\xi_{n+1}. \end{aligned}$$

Next note the size estimate

$$(2.13) \quad |\phi_{mk\nu}(\sigma)| \leq C 2^{-mM} (2^{m+k}\nu^{-1})^{(n-3)/2}.$$

Moreover, $\phi_{mk\nu}$ vanishes unless either $1-\sigma^2 \approx 2^{m+k}\nu^{-1}$ for $m > 0$, or $1-\sigma^2 \leq 2^k\nu^{-1}$ for $m = 0$. Hence if σ is in the support of $\phi_{mk\nu}$, then either $|\nu-\nu\sigma| \leq 2^{m+k}$ or $|\nu+\nu\sigma| \leq 2^{m+k}$. Then using the estimates (2.13), the integrand of (2.12) is bounded by

$$\begin{aligned} &C 2^{-k\delta} |\phi_{mk\nu}(\sigma)| \frac{1}{(1+2^{-k}|\sigma\nu|)^{N_1}} \frac{1}{(1+|t+\rho r\sigma|)^N} \\ &\leq C 2^{k\{(n-3)/2-\delta\}} 2^{m((n-3)/2+N+N_1-M)} \nu^{-(n-3)/2} \xi_{n+1}^{-N} \\ &\quad \times \frac{1}{(1+2^{-k}|\nu|)^{N_1}} \left\{ \frac{1}{(1+|t+\rho r|)^N} + \frac{1}{(1+|t-\rho r|)^N} \right\}. \end{aligned}$$

If we integrate over the support of $\varphi(2^k(1-\rho^2)) \otimes \phi_{mk\nu} \otimes \psi$ for $m \geq 0$ in (2.12), we gain an additional factor of $C 2^m r^{-1}$. Since $M > N + N_1 + (n-1)/2$, we may sum over m and the desired estimates (2.8) follow from (2.12). Hence we obtain

$$\begin{aligned} (1+||t|-r|)^{-N} &\left\{ C \sum_{r \leq 2^k} 2^{-k(\delta+1)} r^{-(n-1)/2} \right. \\ &\quad \left. + C \sum_{r > 2^k} 2^{-k(\delta+1-N_1)} r^{-(n-1)/2-N_1} \right\}, \end{aligned}$$

and thus (2.9) is established for $l = 0$. For the case $l \neq 0$, we use (2.2). □

In Section 4 we will need estimates for the derivatives of the kernels. When $|x| \leq 2$, straightforward computations in (2.1) give us Lemma 3(a). If $|x| = r > 2$, we use (2.7) to show Lemma 3(b).

Lemma 3. (a) Suppose that $|x| \leq 2$. Then for every N ,

$$|G_{k,0}^{(\gamma)}(x,t)| \leq C 2^{-k(\delta+1)} \min\{1, (2^{-k}|t|)^{-N}\}.$$

Moreover,

$$\begin{aligned} |G_0^{(\gamma)}(x,t)| &\leq C \sum_{|t| \leq 2^k} 2^{-k(\delta+1)} + C \sum_{|t| > 2^k} 2^{-k(\delta+1-N)} |t|^{-N} \\ &\leq C \frac{1}{(1+|t|)^{\delta+1}} \quad \text{for } \gamma \in \mathbb{N}^{n+1} \text{ and } |\gamma| = 1, 2, 3, \dots \end{aligned}$$

(b) Suppose that $r > 2$. Then

$$|K_{k,0}^{(\gamma)}(r,t)| \leq C 2^{-k(\delta+1)} r^{-(n-1)/2} \min\{1, (2^{-k}r)^{-N_1}\} (1+||t|-r|)^{-N}.$$

Moreover,

$$\begin{aligned} |K_0^{(\gamma)}(r,t)| &\leq (1+||t|-r|)^{-N} \left\{ C \sum_{r \leq 2^k} 2^{-k(\delta+1)} r^{-(n-1)/2} \right. \\ &\quad \left. + C \sum_{r > 2^k} 2^{-k(\delta+1-N_1)} r^{-(n-1)/2-N_1} \right\} \\ &\leq C (1+r)^{-(n+1+2\delta)/2} \left\{ \frac{1}{(1+|t+r|)^N} + \frac{1}{(1+|t-r|)^N} \right\} \end{aligned}$$

for $\gamma \in \mathbb{N}^2$ and $|\gamma| = 1, 2, 3, \dots$.

3. THE ATOMIC DECOMPOSITION OF H^p AND PRELIMINARY LEMMAS

Definition 1. Let $0 < p \leq 1$ and d be an integer that satisfies $d \geq (n+1)(1/p-1)$. Let Q be a cube in $\mathbb{R}^{n+1}$. We say that a is a (p, d) -atom associated with Q if a is supported on $Q \subset \mathbb{R}^{n+1}$ and satisfies

$$(3.1) \quad \begin{aligned} \text{(i)} \quad &|a(x)| \leq |Q|^{-1/p} \quad \text{almost everywhere} \\ \text{(ii)} \quad &\int_{\mathbb{R}^{n+1}} a(x) x^\beta dx = 0 \end{aligned}$$

where $\beta = (\beta_1, \beta_2, \dots, \beta_{n+1})$ is an $(n+1)$ -tuple of non-negative integers satisfying $|\beta| = \beta_1 + \beta_2 + \dots + \beta_{n+1} \leq d$, and $x^\beta = x^{\beta_1} x^{\beta_2} \dots x^{\beta_{n+1}}$.

If $\{a_i\}$ is a collection of (p, d) -atoms and $\{\lambda_i\}$ is a sequence of complex numbers with $\sum_i |\lambda_i|^p < \infty$, then the series

$$f = \sum_{i=1}^{\infty} \lambda_i a_i$$

converges in the sense of distributions, and its sum belongs to H^p with $\|f\|_{H^p} \leq C (\sum_i |\lambda_i|^p)^{1/p}$ and the converse inequality also holds (see [6]).

The following lemma is due to Stein, Taibleson and Weiss [7]. It is a version of the triangle inequality on $L^{p,\infty}$, $0 < p < 1$.

Lemma 4. Suppose $0 < p < 1$ and $\{f_i\}$ is a sequence of measurable functions such that

$$(3.2) \quad |\{x : |f_i(x)| > \alpha > 0\}| \leq \alpha^{-p}$$

for $i = 1, 2, 3, \dots$. If $\sum_{i=1}^{\infty} |\lambda_i|^p \leq 1$, then

$$(3.3) \quad \left| \left\{ x : \left| \sum_{i=1}^{\infty} \lambda_i f_i(x) \right| > \alpha \right\} \right| \leq \frac{2-p}{1-p} \alpha^{-p}.$$

Proof. See ([7]). $\square$

Lemma 5. For given $\beta > 0$, assume that $0 < p < 1$. Suppose that $\{g_l\}$ is a sequence of measurable functions such that

$$(3.4) \quad |\{x : |g_l(x)| > \alpha\}| \leq A^p 2^{-\beta l p} \alpha^{-p}$$

for $l \geq 0$ and all $\alpha > 0$. Then

$$\left| \left\{ x : \sum_{l \geq 0} |g_l(x)| > \alpha \right\} \right| \leq C \alpha^{-p}.$$

Proof. From (3.4), we have

$$\left| \left\{ x : |g_l(x)| \cdot \left(\frac{2^{\beta l}}{A} \right) > \alpha \right\} \right| \leq A^p 2^{-\beta l p} (A 2^{-\beta l} \alpha)^{-p} = \alpha^{-p}.$$

Moreover, $\sum_{l \geq 0} A^p 2^{-\beta l p} = A_{\beta}^p A^p$ where $A_{\beta}^p = \frac{1}{1-2^{-\beta p}}$.

By Lemma 4, we then obtain

$$\begin{aligned} \left| \left\{ x : \sum_{l \geq 0} |g_l(x)| > \alpha \right\} \right| &\leq \left| \left\{ x : \sum_{l \geq 0} \frac{A 2^{-\beta l}}{A_{\beta} A} \frac{|g_l(x)|}{A 2^{-\beta l}} > \frac{\alpha}{A_{\beta} A} \right\} \right| \\ &\leq \left(\frac{2-p}{1-p} \right) \left(\frac{\alpha}{A_{\beta} A} \right)^{-p} = C \alpha^{-p}. \end{aligned}$$

$\square$

The following technical estimates will be used in Section 4.

Lemma 6. Suppose $0 < p < 1$ and $\delta = n(1/p - 1/2) - 1/2$. Suppose

$$\begin{aligned} &\iint_{\{|x| > 2^{1-l}, |t| > 2^{1-l}; 2^{la} |x|^{-n/p} \chi_{\{|t|-|x| \leq 2^{-l}\}} > \alpha/16C\}} dx dt \\ &+ \iint_{\{|x| > 2^{1-l}, |t| > 2^{1-l}; 2^{lb} |x|^{-n/p} |t|^{-N} > \alpha/16C\}} dx dt \\ &+ \iint_{\{|x| > 2^{1-l}, |t| \leq 2^{1-l}; 2^{lb} |x|^{-n/p-N} > \alpha/16C\}} dx dt \\ &+ \iint_{\{|x| \leq 2^{1-l}, |t| > 2^{1-l}; 2^{lc} |t|^{-(\delta+1)} > \alpha/16C\}} dx dt \\ &\leq C 2^{ld} \alpha^{-p}. \end{aligned}$$

Then

- (i) if $a = n+1-n/p$, $b = n+1-n/p-N$ and $c = n-\delta$, then $d = (n+1)(p-1)$,
- (ii) if $a = n+N+2-n/p$, $b = n+2-n/p$ and $c = n+N+1-\delta$, then $d = (n+N+2)p - (n+1)$.

Proof. Applying Fubini's theorem to the first integral and Chebyshev's inequality to the third and last integrals yields the desired estimates. $\square$

4. WEAK TYPE ESTIMATES

We will show that $T^\delta f$ satisfies the uniform weak type estimates (3.2) when f is a (p, N) -atom ($N \geq (n+1)(1/p - 1)$), and prove Theorem 1.

Proposition 1. *Suppose f is a (p, N) -atom ($N \geq (n+1)(1/p - 1)$) on $\mathbb{R}^{n+1}$ and $\delta = n(1/p - 1/2) - 1/2$. Then there exists a constant $C = C(n, p)$ such that*

$$(4.1) \quad |\{(x, t) \in \mathbb{R}^n \times \mathbb{R} : |T^\delta f(x, t)| > \alpha\}| \leq C \alpha^{-p}$$

for all $\alpha > 0$.

Proof. Since T^δ is translation invariant, we can assume that f is supported in a cube Q of diameter 2^R centered at the origin. We observe

$$\begin{aligned} & |\{(x, t) : |T^\delta f(x, t)| > \alpha\}| \\ & \leq |\{(x, t) \in Q^* : |T^\delta f(x, t)| > \alpha/2\}| \\ & \quad + |\{(x, t) \in (Q^*)^c : |T^\delta f(x, t)| > \alpha/2\}| \end{aligned}$$

where Q^* is the cube concentric with Q and with sides of twice the length, and we will show that each term is bounded by $C \alpha^{-p}$.

Consider $(x, t) \in Q^*$. Suppose $0 < p < 2$ and $p/2 + 1/q = 1$. By Hölder's inequality and the Plancherel theorem, we have

$$\begin{aligned} \iint_{Q^*} |T^\delta f(x, t)|^p dx dt & \leq \left(\iint_{\mathbb{R}^{n+1}} |T^\delta f|^2 dx dt \right)^{p/2} |Q^*|^{1/q} \\ & \leq C. \end{aligned}$$

Hence for all $\alpha > 0$,

$$(4.2) \quad |\{(x, t) \in Q^* : |T^\delta f(x, t)| > \alpha/2\}| \leq C \alpha^{-p}.$$

Next, to establish the following estimate

$$(4.3) \quad |\{(x, t) \in (Q^*)^c : |T^\delta f(x, t)| > \alpha/2\}| \leq C \alpha^{-p}$$

for all $\alpha > 0$, we first assume that f is supported in the cube I of diameter 1 centered at the origin. We now consider the case $(x, t) \in (I^*)^c$ and claim that

$$\begin{aligned} (4.4) \quad |T_l^\delta f(x, t)| & \leq C \left\{ 2^{la} |x|^{-n/p} \chi_{\{|x| > 2^{1-l}, |t| > 2^{1-l}, ||t|-|x|| \leq 2^{-l}\}} \right. \\ & \quad + 2^{lb} |x|^{-n/p} |t|^{-N} \chi_{\{|x| > 2^{1-l}, |t| > 2^{1-l}\}} \\ & \quad + 2^{lb} |x|^{-n/p-N} \chi_{\{|x| > 2^{1-l}, |t| \leq 2^{1-l}\}} \\ & \quad \left. + 2^{lc} |t|^{-(\delta+1)} \chi_{\{|x| \leq 2^{1-l}, |t| > 2^{1-l}\}} \right\} \end{aligned}$$

where

- (i) $a = n + 1 - n/p$, $b = n + 1 - n/p - N$ and $c = n - \delta$ for $l \geq 0$,
- (ii) $a = n + N + 2 - n/p$, $b = n + 2 - n/p$ and $c = n + N + 1 - \delta$ for $l < 0$.

Fix $l \geq 0$. Since f is supported in the cube I of diameter 1, by (2.2) we have

$$|T_l^\delta f(x, t)| \leq 2^{l(n+1)} \iint_{I^*} |f(y, s)| |K_0(2^l|x-y|, 2^l(t-s))| dy ds.$$

Consider the case $2^l|x| > 2$, $2^l|t| > 2$ and $2^l||t| - |x|| \leq 1$. Then by Lemma 2, we have

$$(4.5) \quad |T_l^\delta f(x, t)| \leq C 2^{l(n+1-n/p)} |x|^{-n/p} \chi_{\{|t|-|x| \leq 2^{-l}\}}.$$

If $2^l|x| > 2$ and $2^l||t| - |x|| > 1$, then by Lemma 2, we have

$$(4.6) \quad |T_l^\delta f(x, t)| \leq C 2^{l(n+1-n/p-N)} \{ |x|^{-n/p-N} \chi_{\{|t| \leq 2^{1-l}\}} + |x|^{-n/p} |t|^{-N} \chi_{\{|t| > 2^{1-l}\}} \}.$$

Finally, when $2^l|x| \leq 2$, $2^l|t| > 2$, we use Lemma 1 and thus

$$(4.7) \quad |T_l^\delta f(x, t)| \leq C 2^{l(n-\delta)} |t|^{-(\delta+1)} \chi_{\{|x| \leq 2^{1-l}, |t| > 2^{1-l}\}}.$$

Combining the estimates (4.5)–(4.7), we obtain (4.4) for $l \geq 0$, and by Lemma 6,

$$\iint_{\{|x| > 2^{1-l}, |t| > 2^{1-l} : |T_l^\delta f(x, t)| > \alpha/4\}} dx dt \leq C 2^{l(n+1)(p-1)} \alpha^{-p}.$$

Now applying Lemma 5 with $\beta p = (n+1)(1-p)$, we obtain

$$(4.8) \quad \left| \left\{ (x, t) \in (I^*)^c : \sum_{l \geq 0} |T_l^\delta f(x, t)| > \alpha/4 \right\} \right| \leq C \alpha^{-p}.$$

We now fix $l < 0$. Let $P_{k,l,x}(y, s)$ denote the N -th order Taylor polynomial of the function $(y, s) \rightarrow K_{k,l}(|x-y|, t-s)$ expanded about the origin, the center of the cube. Now $P_{k,l,x} = 2^{l(n+1)} P_{k,0,x}(2^l \cdot, 2^l \cdot)$ for fixed k and l . Then using the moment conditions on f ,

$$\begin{aligned} T_{k,l}^\delta f(x, t) &= \iint_{I^*} f(y, s) 2^{l(n+1)} K_{k,0}(2^l|x-y|, 2^l(t-s)) dy ds \\ &= \iint_{I^*} f(y, s) 2^{l(n+1)} [K_{k,0}(2^l|x-y|, 2^l(t-s)) - P_{k,0,x}(2^l|y|, 2^l s)] dy ds. \end{aligned}$$

A straightforward calculation shows that the absolute value of the last term is dominated by

$$C \iint_{I^*} |f(y, s)| 2^{l(n+1)} \sum_{|\gamma|=N+1} |K_{k,0}^{(\gamma)}(2^l|x|, 2^l t)| |2^l(y, s)|^{N+1} dy ds.$$

We apply Lemma 3 and the same arguments used for (4.5)–(4.7) and thus obtain the bounds (4.4) for $l < 0$. Moreover, by Lemma 6 we see that

$$\iint_{\{|x| > 2^{1-l}, |t| > 2^{1-l} : |T_l^\delta f(x, t)| > \alpha/4\}} dx dt \leq C 2^{l\{(n+N+2)p-(n+1)\}} \alpha^{-p}.$$

Since $(n+1) < (n+N+2)p$, by Lemma 5 with $\beta p = (n+N+2)p - (n+1)$, we obtain

$$(4.9) \quad \left| \left\{ (x, t) \in (I^*)^c : \sum_{l < 0} |T_l^\delta f(x, t)| > \alpha/4 \right\} \right| \leq C \alpha^{-p}.$$

Putting together the estimates (4.8) and (4.9), we obtain (4.3) for the cube I .

Suppose now that f is a (p, N) -atom ($N \geq (n+1)(1/p-1)$), supported in a cube Q of diameter 2^R centered at (x_Q, t_Q) . By translation invariance we can assume

$(x_Q, t_Q) = (0, 0)$. Let $h(x, t) = 2^{R(n+1)/p} f(2^R x, 2^R t)$. Then h is an atom supported in the cube I centered $(0, 0)$. But this implies

$$\begin{aligned} T_l^\delta f(x, t) &= \int_R \int_{R^n} 2^{-R(n+1)/p} h\left(\frac{(x-y)}{2^R}, \frac{(t-s)}{2^R}\right) G_l(y, s) dy ds \\ &= 2^{-R(n+1)/p} 2^{R(n+1)} \left(G_l(2^R \cdot, 2^R \cdot) * h \right) \left(\frac{x}{2^R}, \frac{t}{2^R} \right) \\ &= 2^{-R(n+1)/p} \left(G_{l+R} * h \right) \left(\frac{x}{2^R}, \frac{t}{2^R} \right). \end{aligned}$$

If we repeat the same arguments used for (4.8) and (4.9), we get (4.3). This proves Proposition 1. $\square$

We now proceed with the proof of Theorem 1.

Proof. If $f = \sum_{i=1}^\infty \lambda_i f_i \in H^p(\mathbb{R}^{n+1})$, $T_l^\delta f$ is well defined since each $(T_l^\delta f_i)(x, t)$ is the convolution of the atom f_i with an integrable function G_l . Using the weak type (p, p) estimates of T_l^δ for each l and by Lemma 5, we obtain that $T^\delta f_i$ satisfies a uniform weak type estimate when f_i is a (p, N) -atom ($N \geq (n+1)(1/p - 1)$) in Proposition 1. Since $|T^\delta f(x, t)| \leq \sum_{i=1}^\infty |\lambda_i| |T^\delta f_i(x, t)|$ and $\sum_{i=1}^\infty |\lambda_i|^p < \infty$, Theorem 1 is a consequence of Lemma 4. $\square$

APPENDIX

Let $0 < p < 1$. Denote the quasi-norm $(\sup_{\alpha > 0} \alpha^p |\{x \in \mathbb{R}^n : |g(x)| > \alpha\}|)^{1/p}$ of g in $L^{p,\infty}$ by $\|g\|_{L^{p,\infty}}$. Let $T_m f = m^\vee * f$. We define the class of Fourier multipliers $\mathcal{M}(H^p, L^{p,\infty})(\mathbb{R}^n)$ to be the set of all bounded measurable functions m so that for all $f \in C_0^\infty(\mathbb{R}^n) \cap H^p(\mathbb{R}^n)$,

$$\|T_m f\|_{L^{p,\infty}} \leq C \|f\|_{H^p}.$$

The best constant C is the quasi-norm of the operator T_m , and we write $\|m\|_{\mathcal{M}}$ for this quantity.

Lemma 7. *Suppose f_ϵ and f are measurable functions on $\mathbb{R}^n$ and $f_\epsilon \rightarrow f$ almost everywhere. Assume that $\|f_\epsilon\|_{L^{p,\infty}} \leq M^{1/p}$ for some $M > 0$ and for all $\epsilon > 0$. Let $\alpha > 0$ be fixed. Then $\alpha^p |\{x \in \mathbb{R}^n : |f(x)| > \alpha\}| \leq M$.*

Proof. Suppose $\alpha^p |\{x : |f(x)| > \alpha\}| > M$. Then from the right continuity of $|\{x : |f(x)| > \alpha\}|$, there exists $\beta > \alpha$ such that $\alpha^p |\{x : |f(x)| > \beta\}| > M$. Define $E = \{x : |f(x)| > \alpha\}$. We know $|E| < \infty$. Then by Egoroff's theorem, for every $\eta > 0$, there exists $F \subset E$ such that $|E \setminus F| < \eta$ and $f_\epsilon \rightarrow f$ uniformly on F . We choose ϵ small so that $|f_\epsilon(x) - f(x)| < (\beta - \alpha)/2$ for $x \in F$. Moreover, on F the inequality $|f(x)| > \beta$ implies that $|f_\epsilon(x)| \geq |f(x)| - |f_\epsilon(x) - f(x)| > \beta - (\beta - \alpha)/2 > \alpha$.

From this, $\{x : |f(x)| > \beta\} \cap F \subset \{x : |f_\epsilon(x)| > \alpha\}$ and $\alpha^p |\{x : |f(x)| > \beta\} \cap F| \leq \alpha^p |\{x : |f_\epsilon(x)| > \alpha\}|$. Since $\{x : |f(x)| > \beta\} = (\{x : |f(x)| > \beta\} \cap F) \cup (\{x : |f(x)| > \beta\} \cap (E \setminus F))$,

$$\begin{aligned} &\alpha^p |\{x : |f(x)| > \beta\}| - \alpha^p |E \setminus F| \\ &\leq \alpha^p |\{x : |f(x)| > \beta\} \cap F| \\ &\leq \alpha^p |\{x : |f_\epsilon(x)| > \alpha\}| \leq M. \end{aligned}$$

When η is sufficiently small, this is a contradiction. $\square$

The following is based on de Leeuw's restriction theorem [2].

Theorem 2. *Let $m(\xi', \xi'')$ be contained in the class $\mathcal{M}(H^p, L^{p,\infty})(\mathbb{R}^{k+l})$ and be continuous. Then $m_{\xi''}(\xi') \equiv m(\xi', \xi'')$ is contained in the class $\mathcal{M}(H^p, L^{p,\infty})(\mathbb{R}^k)$ and the multiplier norm of $m_{\xi''}$ does not exceed that of m .*

Proof. Let $f_1 \in C_0^\infty(\mathbb{R}^k) \cap H^p(\mathbb{R}^k)$, $f_{2,\epsilon} \in C_0^\infty(\mathbb{R}^l)$ with $\widehat{f_{2,\epsilon}}(\xi'') = \epsilon^{l(1/p-1)} \phi(\frac{\xi''-a}{\epsilon})$, where ϕ is supported in $B(0, 1)$ (the unit ball about the origin). Define $f_\epsilon(x', x'') = (f_1 \otimes f_{2,\epsilon})(x', x'')$. From this $\|f_\epsilon\|_{H^p} \leq A_\phi \|f_1\|_{H^p}$. Since we have

$$\begin{aligned} & T_m(f_1 \otimes f_{2,\epsilon})(x', x'') \\ &= \frac{1}{(2\pi)^{k+l}} \int_{\xi''} \int_{\xi'} m(\xi', \xi'') \widehat{f_1}(\xi') \epsilon^{l(1/p-1)} \phi\left(\frac{\xi''-a}{\epsilon}\right) e^{i\langle x', \xi' \rangle + i\langle x'', \xi'' \rangle} d\xi' d\xi'' \\ &= T_{m^\epsilon}(f_1 \otimes \epsilon^{1/p} \check{\phi})(x', \epsilon x'') e^{i\langle x'', a \rangle} \end{aligned}$$

where $m^\epsilon(\xi', \xi'') = m(\xi', \epsilon \xi'' + a)$,

$$\|T_{m^\epsilon}(f_1 \otimes \check{\phi})\|_{L^{p,\infty}} \leq A_\phi \|m\|_{\mathcal{M}} \|f_1\|_{H^p}.$$

Then for all $\alpha > 0$, we have

$$\alpha^p |\{(x', x'') : |T_{m^\epsilon}(f_1 \otimes \check{\phi})(x', x'')| > \alpha\}| \leq A_\phi^p \|m\|_{\mathcal{M}}^p \|f_1\|_{H^p}^p$$

and by the Lebesgue Dominated Convergence Theorem, $T_{m^\epsilon}(f_1 \otimes \check{\phi})$ converges to $(T_{m_a} f_1) \otimes \check{\phi}$ as $\epsilon \rightarrow 0$ where $m_a(\xi') = m(\xi', a)$. So from Lemma 7, we have

$$\|(T_{m_a} f_1) \otimes \check{\phi}\|_{L^{p,\infty}} \leq A_\phi \|m\|_{\mathcal{M}} \|f_1\|_{H^p}.$$

Hence m_a is contained in the class $\mathcal{M}(H^p, L^{p,\infty})(\mathbb{R}^k)$ and thus

$$\|m_a\|_{\mathcal{M}} \leq \|m\|_{\mathcal{M}}.$$

□

Remark 1. Even if we replace the continuity assumption of m by almost everywhere conditions, Theorem 2 still holds.

ACKNOWLEDGEMENT

It is a pleasure to thank my advisor A. Seeger for his guidance and helpful observations.

REFERENCES

1. J. Bourgain, *Estimates for the cone multipliers*, Opera. Theory Adv. Appl. **77** (1995), 41–60. MR **96m**:42022
2. M. Jodeit Jr., *A note on the Fourier multipliers*, Proc. Amer. Math. Soc. **27** (1971), 423–424. MR **42**:4965
3. G. Mockenhaupt, *A note on the cone multipliers*, Proc. Amer. Math. Soc. **117** (1993), 145–152. MR **93c**:42015
4. G. Mockenhaupt, A. Seeger and C. D. Sogge, *Wave front sets, local smoothing and Bourgain's circular maximal theorem*, Ann. of Math. **136** (1992), 207–218. MR **93i**:42009
5. D. Müller and A. Seeger, *Inequalities for spherically symmetric solutions of the wave equation*, Math. Z. **3** (1995), 417–426. MR **96e**:35093
6. E. M. Stein, *Harmonic analysis : Real variable method, orthogonality and oscillatory integrals*, Princeton Univ. Press, 1993. MR **95c**:42002

7. E. M. Stein, M. H. Taibleson and G. Weiss, *Weak-type estimates for maximal operators on certain H^p spaces*, Rend. Circ. Mat. Palermo(2) suppl. **1** (1981), 81–97. MR **83c**:42017
8. E. M. Stein and G. Weiss, *Introduction to Fourier Analysis on Euclidean Spaces*, Princeton Univ. Press, 1971. MR **46**:4102

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF WISCONSIN-MADISON, MADISON, WISCONSIN 53706

E-mail address: hong@math.wisc.edu

Current address: Department of Mathematics, Seoul National University, Seoul, 151-742, Korea

E-mail address: shong@math.snu.ac.kr