THE ABELIANIZATION OF ALMOST FREE GROUPS

CHARLY BITTON

(Communicated by Carl G. Jockusch, Jr.)

ABSTRACT. We will construct an almost free non-free group G of cardinality $\aleph_1$ such that G/G' is a free abelian group.

1. INTRODUCTION

In this paper, we will deal with almost free non-free groups and almost free non-free abelian groups. A group G of cardinality λ is said to be almost free if every subgroup of G of cardinality less than λ is free. Similarly, an abelian group A of cardinality λ is said to be almost free if every subgroup of A of cardinality less than λ is free abelian. In 1951, Higman [2] constructed an almost free non-free group of cardinality $\aleph_1$; since then, almost free groups and almost free abelian groups have been investigated by a number of authors. In particular, Shelah [6] proved that if there exists an almost free non-free abelian group of cardinality λ , then there also exists an almost free non-free group of cardinality λ . The converse of this result is still open and only partial results are known in this direction (see [1]). There is a natural way to approach this question. Suppose that G is an almost free non-free group of cardinality λ . Then an easy argument shows that the quotient group G/G' is an almost free abelian group of cardinality λ . (Here G' denotes the commutator subgroup of G .) If it could be shown that G/G' is necessarily non-free, then the problem would be solved. In this paper, we will construct an almost free non-free group G such that G/G' is a free abelian group. Consequently, this naive approach to the problem cannot succeed.

2. THE ABELIANIZATION OF ALMOST FREE NON-FREE GROUPS

In this section, we will construct an almost free non-free group G of cardinality $\aleph_1$ such that G/G' is a free abelian group of cardinality $\aleph_1$.

Before starting the proof, we will review some group theoretic results and establish our notation. Suppose that G is a free group. Then it is well known that every subgroup H of G is free. We will write $H \mid G$ if H is a free factor of G ; i.e., if there exists a basis X of G and a subset $X' \subseteq X$ such that X' is a basis of H . In this case, if E is the subgroup generated by $X \setminus X'$, then $G = H * E$ is the free product of H and E . If Y is any subset of G , then $\langle Y \rangle$ denotes the subgroup of G which is

Received by the editors May 5, 1999 and, in revised form, October 5, 1999.

1991 *Mathematics Subject Classification*. Primary 03E75, 03E05, 20K27.

This is part of the author's Ph.D. thesis done under the supervision of Professor M. Magidor to whom the author is greatly indebted for his help.

generated by Y . If H is any subgroup of G , then H^G denotes the normal closure of H in G ; i.e., the smallest normal subgroup of G which contains H .

We will make repeated use of the following result.

Lemma 2.1. *Suppose that G is a free group and that $K \mid G$. If H is any subgroup of G , then $H \cap K \mid H$.*

Proof. For example, see Problem 32 in Section 2.4 of [4]. $\square$

We will also make use of the following characterization of the commutator subgroup of a free group.

Lemma 2.2. *Suppose that the group A is freely generated by the set $\mathcal{A}$. For each $x \in \mathcal{A}$, let $\ell_x^A : A \rightarrow \mathbb{Z}$ be the function such that $\ell_x^A(w)$ is the sum of the exponents of the occurrences of x in the word $w \in A$. Then $w \in A'$ iff $\ell_x^A(w) = 0$ for every $x \in \mathcal{A}$.*

Proof. For example, see Problem 2 in Section 2.2 of [4]. $\square$

Definition 2.3. If κ is an infinite cardinal, then we say that the principle $F(\kappa)$ holds iff there is a free group A of rank κ and a free subgroup B of rank κ such that:

- (I) If C is a free factor of B of rank less than κ , then C is also a free factor of A .
- (II) B is not a free factor of A .
- (III) $B \cap A' = B'$.

In [2], Higman used a pair of free groups of rank $\aleph_0$ satisfying conditions (I) and (II) of principle $F(\aleph_0)$ to construct an almost free non-free group of cardinality $\aleph_1$. We will essentially give his construction in the proof of Lemma 2.5. We have added condition (III) in order to ensure that B/B' is naturally embedded into A/A' via the map $bB' \mapsto bA'$. Later we will construct a pair of free groups (B, A) satisfying the conditions of principle $F(\aleph_0)$ in such a way that the natural embedding $B/B' \hookrightarrow A/A'$ is surjective. In this case, we will write $B/B' = A/A'$.

Theorem 2.4. *There exists an almost free non-free group G of cardinality $\aleph_1$ such that G/G' is a free abelian group of cardinality $\aleph_1$.*

We will break the proof of Theorem 2.4 into a series of lemmas. We will use the next lemma to construct the group G from a suitably chosen pair (A, B) of free groups exemplifying principle $F(\aleph_0)$.

Lemma 2.5. *Principle $F(\aleph_0)$ implies that there exists an almost free non-free group G of cardinality $\aleph_1$.*

Proof. Suppose that the free groups $B \leq A$ exemplify principle $F(\aleph_0)$. We will define a smooth strictly increasing chain of free group G_α by induction on $\alpha < \aleph_1$. Let G_0 be a free group of rank $\aleph_0$. Now suppose inductively that $\alpha > 0$ and that G_β has been defined for all $\beta < \alpha$. Suppose further that if $\gamma < \beta < \alpha$, then every finitely generated free factor of G_γ is also a free factor of G_β . First suppose that α has the form $\beta + 2$. In this case, we let $G_\alpha = G_{\beta+1} * \langle x_{\beta+1} \rangle$, where $\langle x_{\beta+1} \rangle$ is an infinite cyclic group such that $G_{\beta+1} \cap \langle x_{\beta+1} \rangle = \{e\}$. Next suppose that $\alpha = \delta + 1$ for some limit ordinal δ . Then we let G_α be a free group such that $G_\delta \leq G_\alpha$ and

the following diagram commutes:

$$\begin{array}{ccc} G_\delta & \xrightarrow{\text{isomorphism}} & B \\ \downarrow & & \downarrow \\ G_\alpha = G_{\delta+1} & \xrightarrow{\text{isomorphism}} & A \end{array}$$

Finally if α is a limit ordinal, then we let $G_\alpha = \bigcup_{\beta < \alpha} G_\beta$. In this case, it is necessary to check that G_α is a free group. To see this, first let $\langle g_i \mid i < \omega \rangle$ be an enumeration of the elements of G_α ; then choose inductively an increasing sequence of ordinals

$$\beta_0 \leq \beta_1 \leq \cdots \leq \beta_n \leq \cdots < \alpha$$

and an increasing sequence of subgroups of G_α

$$F_0 \leq F_1 \leq \cdots \leq F_n \leq \cdots$$

such that $\{g_0, \dots, g_n\} \subseteq F_n$ and F_n is a finitely generated free factor of G_{β_n} . By Lemma 2.1, F_n is a free factor of F_{n+1} for each $n < \omega$. Since $G_\alpha = \bigcup_{n < \omega} F_n$, it follows that G_α is a free group.

Now let $G = \bigcup_{\alpha < \aleph_1} G_\alpha$. Then it is clear that G is almost free. Suppose that G is free. Then we can express $G = \bigcup_{\alpha < \aleph_1} K_\alpha$ as the union of a smooth strictly increasing chain of countable subgroups such that $K_\alpha \mid G$ for all $\alpha < \aleph_1$. There exists a club C such that $K_\gamma = G_\gamma$ for all $\gamma \in C$; so by Lemma 2.1, $G_\gamma \mid G_{\gamma+1}$ for all $\gamma \in C$. But by taking $\gamma \in C$ to be a limit ordinal, we obtain a contradiction. $\square$

Lemma 2.6. *In the above construction, $G_\alpha \cap G' = G'_\alpha$.*

Proof. An easy induction shows that if $\alpha < \beta$, then $G_\alpha \cap G'_\beta = G'_\alpha$. (The only point where the induction could conceivably break down is when α is a limit ordinal and $\beta = \alpha + 1$. However, condition (III) of principle $F(\aleph_0)$ deals with this case.) The result follows. $\square$

Now we want to define a particular pair of free groups (B, A) exemplifying principle $F(\aleph_0)$ such that if G is the group constructed in Lemma 2.5, then G/G' is a free abelian group of cardinality $\aleph_1$. We shall make use of the following result of Levi. Recall that if H is any group, then the n^{th} derived subgroup $H^{(n)}$ of H is defined inductively by $H^{(0)} = H$ and $H^{(n+1)} = (H^{(n)})'$.

Lemma 2.7. *If H is a free group, then $\bigcap_{n < \omega} H^{(n)} = \{e\}$.*

Proof. This is an immediate consequence of Corollary 2.12 of [4]. $\square$

Definition 2.8. Let A be the group freely generated by the set $\{a_n \mid n \geq 1\}$ and let

$$B = \langle a_1^{-1}[a_2, a_3], a_2^{-1}[a_3, a_4], a_3^{-1}[a_4, a_5], \dots, a_n^{-1}[a_{n+1}, a_{n+2}], \dots \rangle.$$

Lemma 2.9. (1) *Every finitely generated free factor of B is also a free factor of A .*

(2) *B is not a free factor of A .*

(3) *$B \cap A' = B'$.*

(4) *$B/B' = A/A'$.*

Proof. First we will show that every finitely generated free factor of B is also a free factor of A . For each $n > 2$, define $A_n = \langle a_1, \dots, a_n \rangle$ and

$$B_n = \langle a_1^{-1}[a_2, a_3], a_2^{-1}[a_3, a_4], \dots, a_{n-2}^{-1}[a_{n-1}, a_n] \rangle.$$

Then it is easily checked that $A_n = B_n * \langle a_{n-1}, a_n \rangle$. Hence B_n is a free factor of A . If C is a finitely generated free factor of B , then there exists an integer $n > 2$ such that $C \leq B_n$. By Lemma 2.1, C is a free factor of B_n . Since B_n is a free factor of A , it follows that C is also a free factor of A .

Next we will prove that B is not a free factor of A . To prove this, it is enough to show that $H = A/B^A$ is not a free group. (Recall that B^A denotes the normal closure of B in A .) Note that H is generated by the elements $\{a_n \mid n \geq 1\}$ subject to the relations

$$a_1 = [a_2, a_3], a_2 = [a_3, a_4], a_3 = [a_4, a_5], \dots, a_n = [a_{n+1}, a_{n+2}], \dots$$

So we see that

$$a_1 = [a_2, a_3] \in H^{(1)}$$

and then that

$$a_1 = [a_2, a_3] = [[a_3, a_4], [a_4, a_5]] \in H^{(2)}.$$

Continuing in this fashion, we see that $a_1 \in \bigcap_{n < \omega} H^{(n)}$. It is easily seen that H is isomorphic to the direct limit of the system

$$H_2 \xrightarrow{h_2} H_3 \xrightarrow{h_3} \dots \xrightarrow{h_{n-1}} H_n \xrightarrow{h_n} H_{n+1} \xrightarrow{h_{n+1}} \dots$$

where H_n is the free group $\langle a_{n-1}, a_n \rangle$ and $h_n : H_n \rightarrow H_{n+1}$ is the embedding such that $a_{n-1} \mapsto [a_n, a_{n+1}]$ and $a_n \mapsto a_n$. The element $a_1 \in H$ corresponds to the constant sequence $(a_1, a_1, \dots)$ in the direct limit and so a_1 is a nonidentity element of H . By Lemma 2.7, H is not a free group.

Finally consider the canonical homomorphism $\pi : B \rightarrow A/A'$ such that for each $n \geq 1$,

$$\pi(a_n^{-1}[a_{n+1}, a_{n+2}]) = a_n^{-1}[a_{n+1}, a_{n+2}]A' = a_n^{-1}A'.$$

Clearly π is surjective and $\ker \pi = B \cap A'$. Note that

$$\begin{aligned} w \in \ker \pi &\text{ iff } \ell_x^A(w) = 0 \text{ for all } x \in \{a_n \mid n \geq 1\}, \\ &\text{ iff } \ell_y^B(w) = 0 \text{ for all } y \in \{a_n^{-1}[a_{n+1}, a_{n+2}] \mid n \geq 1\}, \\ &\text{ iff } w \in B'. \end{aligned}$$

(The last equivalence is a consequence of Lemma 2.2.) Thus $B \cap A' = B'$ and $B/B' = A/A'$. $\square$

Lemma 2.10. G/G' is free abelian of cardinality $\aleph_1$.

Proof. For each $\alpha < \aleph_1$, let $A_\alpha = G_\alpha/G'_\alpha$. By Lemma 2.6, we have that

$$\begin{aligned} G/G' &= \bigcup_{\alpha < \aleph_1} G_\alpha/G'_\alpha \cap G' \\ &= \bigcup_{\alpha < \aleph_1} G_\alpha/G'_\alpha = \bigcup_{\alpha < \aleph_1} A_\alpha. \end{aligned}$$

If α is a limit ordinal, then $G_\alpha/G'_\alpha = G_{\alpha+1}/G'_{\alpha+1}$ and so $A_\alpha = A_{\alpha+1}$. On the other hand, if α is a successor ordinal, then $A_{\alpha+1} = A_\alpha \oplus \langle \bar{x}_\alpha \rangle$, where $\bar{x}_\alpha$ denotes

the image of the element $x_\alpha \in G_{\alpha+1}$ in the quotient group $A_{\alpha+1}$. It follows that if $\alpha < \beta$, then A_α is a direct summand of A_β . It is now clear that G/G' is a free abelian group of cardinality $\aleph_1$. $\square$

This completes the proof of Theorem 2.4.

3. CONCLUDING REMARKS

Finally we will say a few words about extending our main result to cardinalities $\lambda > \aleph_1$. We need only consider regular cardinals λ , since Shelah [5] has proved that if G is an almost free group of singular cardinality, then G is free. (The corresponding result is also true for almost free abelian groups of singular cardinality.) In [6], Shelah discovered that the purely combinatorial principle $NPT(\lambda)$ is equivalent to the existence of an almost free non-free abelian group; in [3], it was shown in *ZFC* that $NPT(\lambda)$ holds for an unbounded set of cardinals below the first cardinal fixed point. If $NPT(\lambda)$ holds, then it is possible to generalize the proof of Theorem 2.4 and construct an almost free non-free group G of cardinality λ such that G/G' is a free abelian group. Of course, the real problem is to produce such a group when $NPT(\lambda)$ fails.

ACKNOWLEDGEMENT

The author wishes to express his appreciation to the referee for making many improvements and helpful comments.

REFERENCES

- [1] C. Bitton, *Problems in set theory arising from group theory*, PhD thesis, The Hebrew University, 1998.
- [2] G. Higman, *Almost free groups*, Proc. London Math. Soc. (3) **1** (1951), 284–290. MR **13**:430d
- [3] M. Magidor and S. Shelah, *When does almost free imply free?*, J. Amer. Math. Soc. **7** (1994), 769–830. MR **94m**:03081
- [4] W. Magnus, A. Karrass and D. Solitar, *Combinatorial Group Theory*, Dover Publications, New York, 1966. MR **34**:7617
- [5] S. Shelah, *A compactness theorem for singular cardinals, free algebras, Whitehead problem and transversals*, Israel J. Math **21** (1975), 319–349. MR **52**:10410
- [6] S. Shelah, *Incompactness in regular cardinals*, Notre Dame J. Formal Logic **26** (1985), 195–228. MR **87f**:03095

INSTITUTE OF MATHEMATICS, HEBREW UNIVERSITY, JERUSALEM, ISRAEL

Current address: Department of Mathematics, University of California, Irvine, California 92679

E-mail address: cbitton@math.uci.edu