THE FEFFERMAN-STEIN TYPE INEQUALITY FOR THE KAKEYA MAXIMAL OPERATOR

HITOSHI TANAKA

(Communicated by Christopher D. Sogge)

ABSTRACT. Let K_δ , $0 < \delta \ll 1$, be the Kakeya maximal operator defined as the supremum of averages over tubes of the eccentricity δ . We shall prove the so-called Fefferman-Stein type inequality for K_δ ,

$$\|K_\delta f\|_{L^p(\mathbf{R}^d, w)} \leq C_{d,p} \left(\frac{1}{\delta}\right)^{d/p-1} \left(\log\left(\frac{1}{\delta}\right)\right)^{\alpha(d)} \|f\|_{L^p(\mathbf{R}^d, K_\delta w)},$$

in the range $(1 < p \leq (d^2 - 2)/(2d - 3), d \geq 3)$, with some constants $C_{d,p}$ and $\alpha(d)$ independent of f and the weight w .

1. INTRODUCTION

The purpose of this note is to investigate the so-called Fefferman-Stein type inequality for the Kakeya maximal operator. Throughout this note $0 < \delta \ll 1$ will be a small parameter. For f a locally integrable function on $\mathbf{R}^d$, $d \geq 2$, define

$$(K_{h,\delta} f)(x) = \sup_T \frac{1}{|T|} \int_T |f(y)| dy,$$

where the supremum is taken over all tubes T containing $x \in \mathbf{R}^d$ with the length h and the radius of the cross section $h\delta$. We define the Kakeya maximal operator K_δ by

$$(K_\delta f)(x) = \sup_{h>0} (K_{h,\delta} f)(x).$$

We call a non-negative Borel measurable function w a weight if it is a locally integrable function on $\mathbf{R}^d$. By $w(A)$ we mean the $w(x)dx$ measure of a set A .

It is verified for $d = 2$ that in the range $1 < p \leq d$ the Fefferman-Stein type inequality

$$(1.1) \quad \left(\int_{\mathbf{R}^d} (K_\delta f)(x)^p w(x) dx \right)^{1/p} \leq C_{d,p,\epsilon} \left(\frac{1}{\delta}\right)^{d/p-1+\epsilon} \left(\int_{\mathbf{R}^d} |f(x)|^p (K_\delta w)(x) dx \right)^{1/p}$$

holds for all $\epsilon > 0$ (Müller and F. Solia, [MS]). But in higher dimensions this fact has been verified only in the range $1 < p \leq (d+1)/2$ (A. M. Vargas, [Va]). The main difficulty of this problem lies in making the exponent p as high as possible.

Bourgain proved that an unweighted version of (1.1) (putting $w \equiv 1$) holds in the range $1 < p \leq p_d$, where $(d+1)/2 < p_d < (d+2)/2$ is some exponent given by

Received by the editors December 15, 1999.

2000 *Mathematics Subject Classification*. Primary 42B25.

This work was supported by the Japan Society for the Promotion of Sciences and the Fūjyukai Foundation.

a recursive formula starting from $p_3 = 7/3$ [Bo1]. Wolff improved this result [Wo]. He proved that an unweighted version of (1.1) holds in the range $1 < p \leq (d+2)/2$. Recently, in higher dimensions Bourgain improved it further to $1 < p \leq (1/2 + c)d$ ($c > 0$ independent of d) [Bo2].

A different approach to this problem (an unweighted version) was given by Igari. He investigated the most difficult case $p = d$. He proved that an unweighted version of (1.1) with $p = d$ holds for a special basis [Ig]. He restricted the bases for taking the supremum to only tubes T of which the axis intersects a fixed line. The author proved the weighted version of this restricted result [Ta2]. In this note we shall improve Vargas's result by using this restricted estimates.

The main theorem of this note is the following.

Theorem 1. *Let $d \geq 3$. There exist constants $C_{d,p}$ and $\alpha(d)$ independent of δ , f , and w such that*

$$\|K_\delta f\|_{L^p(\mathbf{R}^d, w)} \leq C_{d,p} \left(\frac{1}{\delta}\right)^{d/p-1} (\log(\frac{1}{\delta}))^{\alpha(d)} \|f\|_{L^p(\mathbf{R}^d, K_\delta w)}$$

holds in the range $1 < p \leq (d^2 - 2)/(2d - 3)$.

By using sieve arguments and three-points interpolation lemma our result can be reduced to the discrete analogue as stated in the following theorem. (See [MS], [Va], and also [Ta2].)

Let $Q = (-1/2, 1/2)^d$ and $\tilde{Q} = (-2, 2)^d$. We divide $\tilde{Q}$ into δ -tubes, Q_i centered at $i \in \mathcal{I}$, where $\mathcal{I}$ is the set of lattice points with the δ -separation.

Theorem 2. *Let $d \geq 3$. For a measurable set $A \subset Q$ and $0 < \lambda \leq 1$ define*

$$I = \{i \in \mathcal{I} : (K_{1,\delta} \chi_A)(i) > \lambda\}.$$

Then

$$(1.2) \quad \sum_{i \in I} w(Q_i) \leq C_d \left(\frac{1}{\delta}\right)^{d-p(d)} \left(\frac{1}{\lambda}\right)^{p(d)} (\log(\frac{1}{\delta}))^{\beta(d)} (K_\delta w)(A),$$

where

$$p(d) = (d^2 - 2)/(2d - 3)$$

and

$$\beta(d) = (d+1)(d-2)/(2d-3).$$

In the following C 's will denote constants which may be different in each occasion but depend only on the dimension d .

2. PROOF OF THEOREM 2

2.1. Preliminaries. We summarize some known results for later use.

Given any line L in $\mathbf{R}^d$ define

$$(K_{1,\delta}^L f)(x) = \sup_T \frac{1}{|T|} \int_T |f(y)| dy,$$

where the supremum is taken over all δ -tubes T which contain x and of which the axis intersects L . Here, δ -tube is the tube with the length 1 and the radius of the cross section δ .

Lemma 3 (Theorem 2 in [Ta2]). *Let $d \geq 3$. Let $\lambda > 0$ and L be any line in $\mathbf{R}^d$. Then*

$$w(\{x \in \mathbf{R}^d : (K_{1,\delta}^L f)(x) > \lambda\}) \leq C\left(\frac{1}{\lambda}\right)^d (\log(\frac{1}{\delta}))^{d+1} \|f\|_{L^d(\mathbf{R}^d, K_\delta w)}^d.$$

Let $\mathcal{B}_{\leq 1/\delta}$ be the class of all rectangles in $\mathbf{R}^d$ which satisfy

$$1 \leq (\text{the length of longest sides})/(\text{the length of shortest sides}) \leq \frac{1}{\delta}.$$

The corresponding maximal operator associated to this base $\mathcal{B}_{\leq 1/\delta}$ will be denoted by $K_{\leq 1/\delta}$.

Lemma 4 (Theorem 3 in [Ta1]). *Let $d \geq 2$. There exist constants C_1 and C_2 depending only on d such that*

$$C_1(K_\delta f)(x) \leq (K_{\leq 1/\delta} f)(x) \leq C_2(K_\delta f)(x)$$

holds for every $x \in \mathbf{R}^d$.

2.2. Main argument. Write $W = K_\delta w$. Fix $A \subset Q$ and $0 < \lambda \leq 1$. Recall $I = \{i \in \mathcal{I} : (K_{1,\delta} \chi_A)(i) > \lambda\}$. Then for every $i \in I$ we can select a δ -tube T_i , which contains i , such that

$$(2.1) \quad |A \cap T_i| > C\delta^{d-1}\lambda.$$

Then, it follows from (2.1) and the Schwarz inequality that

$$\begin{aligned} & (C\delta^{d-1}\lambda \sum_{i \in I} w(Q_i))^2 \\ & \leq \left(\sum_{i \in I} w(Q_i) |A \cap T_i| \right)^2 = \left(\int_A \sum_{i \in I} w(Q_i) \chi_{T_i} \right)^2 \\ & \leq \left\{ \int_A \left(\sum_{i \in I} w(Q_i) \chi_{T_i} \right)^2 W^{-1} \right\} W(A) \\ & \leq \left\{ \sum_{i \in I} w(Q_i) \sum_{j \in I} w(Q_j) W^{-1}(T_i \cap T_j) \right\} W(A) \\ & \leq \max_{i \in I} \left(\sum_{j \in I} w(Q_j) W^{-1}(T_i \cap T_j) \right) \cdot \left(\sum_{i \in I} w(Q_i) \right) W(A). \end{aligned}$$

Hence

$$(2.2) \quad N^{-1} \sum_{i \in I} w(Q_i) \leq C \left(\frac{1}{\delta} \right)^{2(d-1)} \left(\frac{1}{\lambda} \right)^2 W(A),$$

which corresponds to low multiplicity of Wolff (see the proof of Lemma 3.1 in [Wo]), where

$$N = \max_{i \in I} \left(\sum_{j \in I} w(Q_j) W^{-1}(T_i \cap T_j) \right).$$

By the fact that $p(d) > 2$ and $\frac{1}{\lambda} \geq 1$, we may assume that

$$(2.3) \quad C\delta^{2(d-1)} \leq N.$$

The following proposition, corresponding to high multiplicity of Wolff, will be proven later.

Proposition 5. *With previous setup we have*

$$(2.4) \quad N \left\{ \sum_{i \in I} w(Q_i) \right\}^{(d-2)/(d-1)} \leq C \left(\frac{1}{\delta} \right)^{-d} \left\{ \left(\log \left(\frac{1}{\delta} \right) \right)^{d+1} \left(\frac{1}{\lambda} \right)^d W(A) \right\}^{(d-2)/(d-1)}.$$

Multiplying both sides of (2.2) and (2.4) together, we obtain the desired inequality (1.2).

2.3. Proof of Proposition 5. Take some $i_0 \in I$ so that

$$(2.5) \quad N = \sum_{j \in I} w(Q_j) W^{-1}(T_{i_0} \cap T_j).$$

Let

$$(2.6) \quad I_0 = \{j \in I : T_{i_0} \cap T_j \neq \emptyset\}$$

and

$$(2.7) \quad s_0 = \inf_{y \in T_{i_0}} W(y).$$

By the geometric observation of Córdoba [Co] one sees that

$$(2.8) \quad |T_{i_0} \cap T_j| \leq C \frac{\delta^d}{\delta + \text{dist}(T_{i_0}, j)}.$$

From (2.5)–(2.8) we have

$$(2.9) \quad N \leq C(s_0)^{-1} \sum_{j \in I_0} w(Q_j) \frac{\delta^d}{\delta + \text{dist}(T_{i_0}, j)}.$$

Define the subset of I_0 as

$$\sigma_k = \{j \in I_0 : (k-1)\delta \leq \text{dist}(T_{i_0}, j) < k\delta\}, \quad k = 1, 2, \dots,$$

and rewrite

$$(s_0)^{-1} \sum_{j \in I_0} w(Q_j) \frac{\delta^d}{\delta + \text{dist}(T_{i_0}, j)} = (s_0)^{-1} \sum_k \sum_{j \in \sigma_k} w(Q_j) \frac{\delta^d}{\delta + \text{dist}(T_{i_0}, j)}.$$

Then

$$(2.10) \quad N \leq C(s_0)^{-1} \delta^{d-1} \sum_k \sum_{j \in \sigma_k} \frac{w(Q_j)}{k}.$$

It follows for some k_0 to be specified later that

$$\begin{aligned} & \delta^{d-1} \sum_{k=1}^{k_0} \sum_{j \in \sigma_k} \frac{w(Q_j)}{k} \\ &= \delta^{d-1} \sum_{k=1}^{k_0} \sum_{j \in \sigma_k} w(Q_j) \left(\sum_{l=k}^{k_0} \frac{1}{l(l+1)} + \frac{1}{k_0+1} \right) \\ &= \left\{ \delta^{d-1} \sum_{k=1}^{k_0} \sum_{j \in \sigma_k} w(Q_j) \left(\sum_{l=k}^{k_0} \frac{1}{l(l+1)} \right) \right\} + \left\{ \frac{\delta^{d-1}}{k_0+1} \sum_{k=1}^{k_0} \sum_{j \in \sigma_k} w(Q_j) \right\} \\ &= \text{I} + \text{II}. \end{aligned}$$

By reversing the order of summation we have

$$\begin{aligned} \text{I} &= \delta^{d-1} \sum_{l=1}^{k_0} \left(\sum_{k=1}^l \sum_{j \in \sigma_k} w(Q_j) \right) / (l(l+1)) \\ &\leq C \delta^{2d-2} \sum_{l=1}^{k_0} l^{d-3} \left(\sum_{k=1}^l \sum_{j \in \sigma_k} w(Q_j) \right) / ((l\delta)^{d-1}). \end{aligned}$$

By using Lemma 4 we see that

$$(s_0)^{-1} \left(\sum_{k=1}^l \sum_{j \in \sigma_k} w(Q_j) \right) / ((l\delta)^{d-1}) \leq C (s_0)^{-1} \frac{\int_{R_l} w}{|R_l|} \leq C,$$

where

$$R_l = \{x \in \mathbf{R}^d : \text{dist}(T_{i_0}, x) \leq l\delta\}.$$

Hence

$$(s_0)^{-1} \text{I} \leq C \delta^{2d-2} \sum_{l=1}^{k_0} l^{d-3} \leq C \delta^d (k_0 \delta)^{d-2}.$$

Combining these inequalities we obtain

$$(2.11) \quad N \leq C \delta^d \{ (k_0 \delta)^{d-2} + (k_0 \delta)^{-1} (s_0)^{-1} \sum_{j \in I_0} w(Q_j) \}.$$

Now, we can choose some k_0 so that

$$(k_0 \delta)^{d-1} \sim (s_0)^{-1} \sum_{j \in I_0} w(Q_j)$$

by (2.3) and (2.10). Then the two terms in the right-hand side of (2.11) balance and hence

$$(2.12) \quad N \leq C \delta^d \{ (s_0)^{-1} \sum_{j \in I_0} w(Q_j) \}^{(d-2)/(d-1)}.$$

Applying Lemma 3 with $L =$ the axis of T_{i_0} and $f = \chi_A$, we clearly obtain

$$(2.13) \quad \sum_{j \in I_0} w(Q_j) \leq C \left(\frac{1}{\lambda} \right)^d \left(\log \left(\frac{1}{\delta} \right) \right)^{d+1} W(A).$$

Thus, from (2.12) and (2.13) we have

$$(2.14) \quad N (s_0)^{(d-2)/(d-1)} \leq C \delta^d \left\{ \left(\frac{1}{\lambda} \right)^d \left(\log \left(\frac{1}{\delta} \right) \right)^{d+1} W(A) \right\}^{(d-2)/(d-1)}.$$

Finally, again by Lemma 4 we observe that

$$(2.15) \quad \sum_{i \in I} w(Q_i) \leq C \frac{w(\tilde{Q})}{|\tilde{Q}|} \leq C' s_0.$$

Thus, from (2.14) and (2.15) we obtain

$$N \left\{ \sum_{j \in I} w(Q_j) \right\}^{(d-2)/(d-1)} \leq C \delta^d \left\{ \left(\frac{1}{\lambda} \right)^d \left(\log \left(\frac{1}{\delta} \right) \right)^{d+1} W(A) \right\}^{(d-2)/(d-1)}. \quad \square$$

REFERENCES

- [Bo1] J. Bourgain, *Besicovitch type maximal operators and applications to Fourier analysis*, Geom. Funct. Anal., **1** (1990), 147–187. MR **92g**:42010
- [Bo2] J. Bourgain, *On the dimension of Keakeya sets and related maximal inequalities*, Geom. Funct. Anal., **9** (1999), 256–282. MR **2000b**:42013
- [Co] A. Córdoba, *The Keakeya maximal function and the spherical summation multiplier*, Amer. J. Math., **99** (1977), 1–22. MR **56**:6259
- [Ig] S. Igari, *The Keakeya maximal operator with a special base*, Approx. Theory and its Appl., **13** (1997), 1–7. MR **99d**:42030
- [MS] D. Müller and F. Soria, *A double-weight L^2 inequality for the Keakeya maximal function*, Fourier Anal. Appl., Kahane Special Issue (1995), 467–478. MR **96k**:42026
- [Ta1] H. Tanaka, *Some weighted inequalities for the Keakeya maximal operator on functions of product type*, J. Math. Sci. Univ. Tokyo, **6** (1999), 315–333. CMP 99:17
- [Ta2] H. Tanaka, *A weighted inequality for the Keakeya maximal operator with a special base*, Tokyo J. Math., to appear.
- [Va] A. M. Vargas, *A weighted inequality for the Keakeya maximal operator*, Proc. Amer. Math. Soc., **120** (1994), 1101–1105. MR **94f**:42023
- [Wo] T. Wolff, *An improved bound for Keakeya type maximal functions*, Rev. Mat. Iberoamericana, **11** (1995), 651–674. MR **96m**:42034

DEPARTMENT OF MATHEMATICS, GAKUSHUIN UNIVERSITY, 1-5-1 MEJIRO, TOSHIMA-KU, TOKYO 171-8588, JAPAN

E-mail address: hitoshi.tanaka@gakushuin.ac.jp