SOME CLASSES OF TOPOLOGICAL QUASI *-ALGEBRAS

F. BAGARELLO, A. INOUE, AND C. TRAPANI

(Communicated by David R. Larson)

ABSTRACT. The completion $\overline{\mathcal{A}}[\tau]$ of a locally convex *-algebra $\mathcal{A}[\tau]$ with not jointly continuous multiplication is a *-vector space with partial multiplication xy defined only for x or $y \in \mathcal{A}_0$, and it is called a topological quasi *-algebra. In this paper two classes of topological quasi *-algebras called strict CQ*-algebras and HCQ*-algebras are studied. Roughly speaking, a strict CQ*-algebra (resp. HCQ*-algebra) is a Banach (resp. Hilbert) quasi *-algebra containing a C*-algebra endowed with another involution $\#$ and C*-norm $\|\cdot\|_{\#}$. HCQ*-algebras are closely related to left Hilbert algebras. We shall show that a Hilbert space is a HCQ*-algebra if and only if it contains a left Hilbert algebra with unit as a dense subspace. Further, we shall give a necessary and sufficient condition under which a strict CQ*-algebra is embedded in a HCQ*-algebra.

1. INTRODUCTION

Topological quasi *-algebras were first introduced by Lassner [6] for the mathematical description of some quantum physical models, and after that, they have been studied by Lassner [6, 7], Trapani [10] and Bagarello-Trapani [3, 4], etc. In this paper we shall study two classes of topological quasi *-algebras called strict CQ*-algebras and HCQ*-algebras from a mathematical point of view but also in the perspective of possible physical applications. Let $\mathcal{A}$ be a *-algebra with two involutions $*$ and $\#$ and two norms $\|\cdot\|$ and $\|\cdot\|_{\#}$ satisfying $\|x^*\| = \|x\|$, $\|x\| \leq \|x\|_{\#}$ and $\|x^{\#}x\|_{\#} = \|x\|_{\#}^2$ for each $x, y \in \mathcal{A}$. Then the completion $\overline{\mathcal{A}}[\|\cdot\|, \|\cdot\|_{\#}]$ of $\mathcal{A}[\|\cdot\|, \|\cdot\|_{\#}]$ is a topological quasi *-algebras containing (under natural assumptions) two C*-algebras $\mathcal{A}_{\#}[\|\cdot\|_{\#}]$ and $\mathcal{A}_{\flat}[\|\cdot\|_{\flat}]$ with different involutions $\#$ and $\flat$, respectively, which are connected by the isometric involution $J : x \rightarrow x^*$. This is called a *pseudo CQ*-algebra*. If $\|x\|_{\#} = \sup\{\|xy\|; \|y\| \leq 1\}$, then $\overline{\mathcal{A}}[\|\cdot\|, \|\cdot\|_{\#}]$ is a particular kind of CQ*-algebra as defined and studied in [1, 2], and it is called a *strict CQ*-algebra*, and denoted by $(\overline{\mathcal{A}}[\|\cdot\|, \|\cdot\|_{\#}], \#, \|\cdot\|_{\#})$. Let $\overline{\mathcal{A}}[\|\cdot\|, \|\cdot\|_{\#}]$ be a topological quasi *-algebra with isometric involution $J : x \rightarrow x^*$ and Hilbertian norm $\|\cdot\|$. If $\mathcal{A}$ has another involution $\#$ satisfying $\|x\| \leq \|L_x\|$ and $L_x^* = L_{x^{\#}}$ for each $x \in \mathcal{A}$, where L_x is the bounded linear operator on the Hilbert space $\overline{\mathcal{A}}[\|\cdot\|, \|\cdot\|_{\#}]$ defined by $L_x y = xy, y \in \mathcal{A}$, then $\overline{\mathcal{A}}[\|\cdot\|, \|\cdot\|_{\#}]$ is a strict CQ*-algebra with involution $\#$ and C*-norm $\|x\|_{\#} \equiv \|L_x\|, x \in \mathcal{A}$, and it is called a *HCQ*-algebra* and denoted by $(\overline{\mathcal{A}}[\|\cdot\|, \|\cdot\|_{\#}], \#)$. HCQ*-algebras are closely related to left Hilbert algebras. Let $(\overline{\mathcal{A}}[\|\cdot\|, \|\cdot\|_{\#}], \#)$ be an HCQ*-algebra. Then $\mathcal{A}$ is

Received by the editors February 20, 2000.

2000 *Mathematics Subject Classification*. Primary 46K70.

Key words and phrases. Topological quasi *-algebras, CQ*-algebras, HCQ*-algebras.

a left Hilbert algebra in the Hilbert space $\overline{\mathcal{A}}[\|\cdot\|]$ with involution $\#$, and the full left Hilbert algebra $\mathcal{A}''$ of $\mathcal{A}$ has unit. But, the isometric involution J does not necessarily coincide with the modular conjugation operator $J_{\mathcal{A}}$ of the left Hilbert algebra $\mathcal{A}$. If $J_{\mathcal{A}} = J$, then the HCQ*-algebra $(\overline{\mathcal{A}}[\|\cdot\|], \#)$ is said to be *standard*. Suppose that $(\overline{\mathcal{A}}[\|\cdot\|], \#)$ is standard. Then $\mathcal{A}$ is contained in the maximal Tomita algebra $(\mathcal{A}'')_0$ of $\mathcal{A}''$ and $(\overline{(\mathcal{A}'')_0}[\|\cdot\|], \#)$ is a standard HCQ*-algebra with the one-parameter group $\{\Delta_{\mathcal{A}}^{it}\}_{t \in \mathbb{R}}$ of *-automorphisms, where $\Delta_{\mathcal{A}}$ is the modular operator of $\mathcal{A}$. From these results, it is shown that a Hilbert space is a standard HCQ*-algebra if and only if it contains a left Hilbert algebra as dense subspace. Finally, we give a necessary and sufficient condition under which a strict CQ*-algebra is embedded into a standard HCQ*-algebra using the GNS-construction of positive sesquilinear form on the strict CQ*-algebra $\overline{\mathcal{A}}[\|\cdot\|]$.

2. STRICT CQ*-ALGEBRAS AND HCQ*-ALGEBRAS

Let $\mathcal{A}[\|\cdot\|]$ be a normed *-algebra with isometric involution $*$ and separately (but not jointly) continuous multiplication. Then the completion, $\overline{\mathcal{A}}[\|\cdot\|]$, of $\mathcal{A}[\|\cdot\|]$ is a topological quasi *-algebra that we call, as is natural, a *Banach quasi *-algebra*. In particular, if $\|\cdot\|$ is a Hilbertian norm, then $\overline{\mathcal{A}}[\|\cdot\|]$ is called a *Hilbert quasi *-algebra*. For any $a \in \overline{\mathcal{A}}[\|\cdot\|]$ we put

$$L_a x = ax \text{ and } R_a x = xa, \quad x \in \mathcal{A}.$$

Then L_a and R_a are linear maps of $\mathcal{A}$ into $\overline{\mathcal{A}}[\|\cdot\|]$. In particular, if $a \in \mathcal{A}$, then L_a and R_a can be extended to bounded linear operators on the Banach space $\overline{\mathcal{A}}[\|\cdot\|]$ and they are denoted by the same symbols L_a and R_a .

Let $\overline{\mathcal{A}}[\|\cdot\|]$ be a Banach quasi *-algebra and assume that the *-algebra $\mathcal{A}$ has another norm $\|\cdot\|_{\#}$ and another involution $\#$ satisfying the following conditions:

- (a.1) $\|x^{\#}x\|_{\#} = \|x\|_{\#}^2, \quad \forall x \in \mathcal{A}.$
- (a.2) $\|x\| \leq \|x\|_{\#}, \quad \forall x \in \mathcal{A}.$
- (a.3) $\|xy\| \leq \|x\|_{\#}\|y\|, \quad \forall x, y \in \mathcal{A}.$

Then by (a.2), the identity map $i : \mathcal{A}[\|\cdot\|_{\#}] \rightarrow \mathcal{A}[\|\cdot\|]$ has a continuous extension $\hat{i}$ from the completion $\mathcal{A}_{\#}$ of $\mathcal{A}[\|\cdot\|_{\#}]$ ($\mathcal{A}_{\#}$ is, of course, a C*-algebra) into $\overline{\mathcal{A}}[\|\cdot\|]$. If $\hat{i}$ is injective, then $\mathcal{A}_{\#}$ is (identified with) a dense subspace of $\overline{\mathcal{A}}$. This happens if, and only if,

- (a.4) two norms $\|\cdot\|$ and $\|\cdot\|_{\#}$ are compatible in the following sense [5]: for any sequence $\{x_n\} \subset \mathcal{A}$ such that $\|x_n\| \rightarrow 0$ and $x_n \rightarrow x$ in $\mathcal{A}_{\#}[\|\cdot\|_{\#}]$, $x = 0$ results, i.e. if $\hat{i}^{-1} : \mathcal{A}[\|\cdot\|] \rightarrow \mathcal{A}_{\#}[\|\cdot\|_{\#}]$ is closable.

Definition 2.1. A Banach quasi *-algebra $\overline{\mathcal{A}}[\|\cdot\|]$ is said to be a *psuedo CQ*-algebra* if the *-algebra $\mathcal{A}$ has a another norm $\|\cdot\|_{\#}$ and another involution $\#$ satisfying the conditions (a.1)–(a.4) above. Furthermore, if $\|x\|_{\#} = \|L_x\| \equiv \sup\{\|xy\|; y \in \mathcal{A} \text{ s.t. } \|y\| \leq 1\}$ for each $x \in \mathcal{A}$, then $\overline{\mathcal{A}}[\|\cdot\|]$ is said to be a *strict CQ*-algebra*.

A pseudo CQ*-algebra $\overline{\mathcal{A}}[\|\cdot\|]$ is fully determined by the involution $\#$ and the C*-norm $\|\cdot\|_{\#}$, and so it will often be denoted by $(\overline{\mathcal{A}}[\|\cdot\|], \#, \|\cdot\|_{\#})$. On the other hand, a strict CQ*-algebra is fully determined when the new involution $\#$ is known; so it can be simply denoted as $(\overline{\mathcal{A}}[\|\cdot\|], \#)$, making lighter in this way the notation introduced by two of us in [1, 2]. Let $(\overline{\mathcal{A}}[\|\cdot\|], \#, \|\cdot\|_{\#})$ be a pseudo CQ*-algebra and, as above, let $\mathcal{A}_{\#}$ be the C*-algebra obtained by completing the #-algebra $\mathcal{A}$ with respect to the C*-norm $\|\cdot\|_{\#}$. Let J be the involution $*$ of the Banach quasi

-algebra $\overline{\mathcal{A}}[\|\cdot\|]$. Then $\mathcal{A}_{\flat} \equiv J\mathcal{A}_{\#}$ is a C-algebra equipped with the operations $x^* + y^* \equiv (x + y)^*$, $\lambda x^* \equiv (\overline{\lambda}x)^*$, $x^*y^* \equiv (yx)^*$, the involution $(x^*)^{\flat} \equiv x^{\#*}$ and the C*-norm $\|x^*\|_{\flat} \equiv \|x\|_{\#}$, $\forall x, y \in \mathcal{A}_{\flat}$.

Proposition 2.2. *A pseudo CQ*-algebra $(\overline{\mathcal{A}}[\|\cdot\|], \#, \|\cdot\|_{\#})$ contains two C*-algebras $\mathcal{A}_{\#}$ and $\mathcal{A}_{\flat} \equiv J\mathcal{A}_{\#}$ with different involutions $\#$ and $\flat$, respectively, as dense subalgebra. In particular, if $(\overline{\mathcal{A}}[\|\cdot\|], \#, \|\cdot\|_{\#})$ is a strict CQ*-algebra, then $L_{\mathcal{A}_{\#}}$ and $R_{\mathcal{A}_{\flat}}$ are C*-algebras, $L_x R_y = R_y L_x$ for each $x \in \mathcal{A}_{\#}$ and $y \in \mathcal{A}_{\flat}$ and $R_{\mathcal{A}_{\flat}} = J L_{\mathcal{A}_{\#}} J$.*

By Proposition 2.2, every strict CQ*-algebra is a CQ*-algebra in the sense of [1, 2] but the converse is not true in general ($(\mathcal{A}_{\#} \cap \mathcal{A}_{\flat})$ is not required to be $\#$ -invariant).

We summarize the situation with the following scheme:

$$\begin{array}{ccccc} & & \mathcal{A}_{\#} & & \\ & \subset & \downarrow J & \subset & \\ \mathcal{A}[\|\cdot\|] & & \mathcal{A}_{\flat} & & \overline{\mathcal{A}}[\|\cdot\|] \\ & \subset & & \subset & \\ \text{normed } *- \text{algebra} & & \text{C}^* \text{-algebras} & & \text{CQ}^* \text{-algebra,} \end{array}$$

which summarizes the situation: the *-algebra $\mathcal{A}[\|\cdot\|]$ is contained in its closures, $\mathcal{A}_{\#} = \overline{\mathcal{A}}[\|\cdot\|_{\#}]$ and $\mathcal{A}_{\flat} = \overline{\mathcal{A}}[\|\cdot\|_{\flat}] = J\mathcal{A}_{\#}$. These C*-algebras, moreover, are both contained in $\overline{\mathcal{A}}[\|\cdot\|]$.

Definition 2.3. A Hilbert quasi *-algebra $\overline{\mathcal{A}}[\|\cdot\|]$ is said to be a *HCQ*-algebra* if there is another involution $\#$ of $\mathcal{A}$ such that $L_x^* = L_{x\#}$ and $\|x\| \leq \|L_x\|$ for each $x \in \mathcal{A}$. Here we denote it by $(\overline{\mathcal{A}}[\|\cdot\|], \#)$.

HCQ*-algebras are closely related to left Hilbert algebras. Before going forth, for the reader's convenience, we briefly review the definitions and the basic properties of left Hilbert algebras. A *-algebra $\mathfrak{A}$ with involution $\#$ is said to be a *left Hilbert algebra* if it is a dense subspace in a Hilbert space $\mathcal{H}$ with inner product $(\cdot | \cdot)$ satisfying the following conditions:

- (i) For any $x \in \mathfrak{A}$ the map $y \in \mathfrak{A} \rightarrow xy \in \mathfrak{A}$ is continuous.
- (ii) $(xy|z) = (y|x^{\#}z)$, $\forall x, y, z \in \mathfrak{A}$.
- (iii) $\mathfrak{A}^2 \equiv \{xy; x, y \in \mathfrak{A}\}$ is total in $\mathcal{H}$.
- (iv) The involution $x \rightarrow x^{\#}$ is closable in $\mathcal{H}$.

By (i), for any $x \in \mathfrak{A}$ we denote by $\pi_{\mathfrak{A}}(x)$ the unique continuous linear extension to $\mathcal{H}$ of the map $y \in \mathfrak{A} \rightarrow xy \in \mathfrak{A}$; then $\pi_{\mathfrak{A}}$ is a *-representation of $\mathfrak{A}$ on $\mathcal{H}$.

We denote by $S_{\mathfrak{A}}$ the closure of the involution $\#$. Let $S_{\mathfrak{A}} = J_{\mathfrak{A}} \Delta_{\mathfrak{A}}^{\frac{1}{2}}$ be the polar decomposition of $S_{\mathfrak{A}}$. Then $J_{\mathfrak{A}}$ is an isometric involution on $\mathcal{H}$ and $\Delta_{\mathfrak{A}}$ is a non-singular positive self-adjoint operator in $\mathcal{H}$ such that $S_{\mathfrak{A}} = J_{\mathfrak{A}} \Delta_{\mathfrak{A}}^{\frac{1}{2}} = \Delta_{\mathfrak{A}}^{-\frac{1}{2}} J_{\mathfrak{A}}$ and $S_{\mathfrak{A}}^* = J_{\mathfrak{A}} \Delta_{\mathfrak{A}}^{-\frac{1}{2}} = \Delta_{\mathfrak{A}}^{\frac{1}{2}} J_{\mathfrak{A}}$, and $J_{\mathfrak{A}}$ is called the *modular conjugation operator* of $\mathfrak{A}$ and $\Delta_{\mathfrak{A}}$ is called the *modular operator* of $\mathfrak{A}$. We define the commutant $\mathfrak{A}'$ of $\mathfrak{A}$ as follows: For any $y \in \mathcal{D}(S_{\mathfrak{A}}^*)$ we put $\pi'_{\mathfrak{A}}(y)x = \pi_{\mathfrak{A}}(x)y$, $x \in \mathfrak{A}$ and put $\mathfrak{A}' = \{y \in \mathcal{D}(S_{\mathfrak{A}}^*); \pi'_{\mathfrak{A}}(y) \text{ is bounded}\}$. Then $\mathfrak{A}'$ is a left Hilbert algebra in $\mathcal{H}$ with involution $S_{\mathfrak{A}}^*$ and multiplication $y_1 y_2 \equiv \pi'_{\mathfrak{A}}(y_2) y_1$. Similarly, the commutant $\mathfrak{A}''$ of $\mathfrak{A}'$ is defined by $\mathfrak{A}'' = \{x \in \mathcal{D}(S_{\mathfrak{A}}); y \in \mathfrak{A}' \rightarrow xy \text{ is continuous}\}$. For any $x \in \mathfrak{A}''$ we denote by $\pi_{\mathfrak{A}}(x)$ the unique continuous linear operator on $\mathcal{H}$ such that $\pi_{\mathfrak{A}}(x)y = \pi'_{\mathfrak{A}}(y)x$, $y \in \mathfrak{A}'$. Then $\mathfrak{A}''$ is a left Hilbert algebra in $\mathcal{H}$ with involution $S_{\mathfrak{A}}$ and multiplication $x_1 x_2 \equiv \pi_{\mathfrak{A}}(x_1) x_2$ containing $\mathfrak{A}$. A left Hilbert algebra $\mathfrak{A}$ is said to be *full* if $\mathfrak{A} = \mathfrak{A}''$. It is well-known as the Tomita fundamental theorem that

$J_{\mathfrak{A}}\pi_{\mathfrak{A}}(\mathfrak{A})''J_{\mathfrak{A}} = \pi_{\mathfrak{A}}(\mathfrak{A})'$ and $\Delta_{\mathfrak{A}}^{it}\pi_{\mathfrak{A}}(\mathfrak{A})''\Delta_{\mathfrak{A}}^{-it} = \pi_{\mathfrak{A}}(\mathfrak{A})'', \forall t \in \mathbb{R}$. Let $\mathfrak{A}$ be a full left Hilbert algebra in $\mathcal{H}$, and $\mathfrak{A}_0 \equiv \{x \in \bigcap_{\alpha \in \mathbb{C}} \mathcal{D}(\Delta_{\mathfrak{A}}^{\alpha}); \Delta_{\mathfrak{A}}^{\alpha}x \in \mathfrak{A}, \forall \alpha \in \mathbb{C}\}$. Then $\mathfrak{A}_0$ is a left Hilbert subalgebra in $\mathcal{H}$ such that $\mathfrak{A}_0'' = \mathfrak{A}$, $J_{\mathfrak{A}}\mathfrak{A}_0 = \mathfrak{A}_0$ and $\{\Delta_{\mathfrak{A}}^{\alpha}; \alpha \in \mathbb{C}\}$ is a complex one-parameter group of automorphisms of $\mathfrak{A}_0$ such that $(\Delta_{\mathfrak{A}}^{\alpha}x)^{\#} = \Delta_{\mathfrak{A}}^{-\bar{\alpha}}x^{\#}$ and $(\Delta_{\mathfrak{A}}^{\alpha}x)^{*} = \Delta_{\mathfrak{A}}^{-\bar{\alpha}}x^{*}$ for each $\alpha \in \mathbb{C}$ and $x \in \mathfrak{A}_0$. This $\mathfrak{A}_0$ is called the *maximal Tomita algebra* of $\mathfrak{A}$. For more details, see [8, 9, 11].

Proposition 2.4. *Suppose that $(\overline{\mathcal{A}}[\|\cdot\|], \#)$ is a HCQ^{*} -algebra. Then the following statements hold:*

- (i) $(\overline{\mathcal{A}}[\|\cdot\|], \#)$ is a strict CQ^{*} -algebra with the C^{*} -norm $\|x\|_{\#} = \|L_x\|, x \in \mathcal{A}$.
- (ii) $\mathcal{A}$ is a left Hilbert algebra in the Hilbert space $\mathcal{H} \equiv \overline{\mathcal{A}}[\|\cdot\|]$ whose full left Hilbert algebra $\mathcal{A}''$ has a unit u .

Proof. (i) The proof is mostly trivial. We prove only that the condition (a.4) is satisfied in this case. Indeed, if $\{x_n\} \subset \mathcal{A}$ is a sequence such that $\|x_n\| \rightarrow 0$ and $x_n \rightarrow x$ in $\mathcal{A}_{\#}[\|\cdot\|]$, then by the assumption $L_{x_n} \rightarrow L_x$ with respect to the operator norm. The continuity of the multiplication in $\mathcal{A}[\|\cdot\|]$ easily implies that $L_x = 0$; thus $\|x\|_{\#} = 0$ and $x = 0$.

(ii) We first show that $\mathcal{A}$ is a left Hilbert algebra in $\mathcal{H}$ with involution $\#$. Since the C^{*} -algebra $\mathcal{A}_{\#}$ has an approximate identity $\{u_{\alpha}\}$, $\mathcal{A}$ is dense in the C^{*} -algebra $\mathcal{A}_{\#}$ and $\|x\| \leq \|x\|_{\#}$ for each $x \in \mathcal{A}$, and then it follows that $\mathcal{A}^2$ is total in $\mathcal{A}[\|\cdot\|]$. The assumption $L_x^{*} = L_{x^{\#}} (\forall x \in \mathcal{A})$ implies that $(xy|z) = (y|x^{\#}z)$ for each $x, y, z \in \mathcal{A}$, where $(\cdot|\cdot)$ is the inner product defined by the Hilbertian norm $\|\cdot\|$. Further, we have $\pi_{\mathcal{A}}(x) = L_x, \forall x \in \mathcal{A}$ and $\pi_{\mathcal{A}}(x)$ is bounded. Take any sequence $\{x_n\}$ in $\mathcal{A}$ such that $\lim_{n \rightarrow \infty} \|x_n\| = 0$ and $\lim_{n \rightarrow \infty} \|x_n^{\#} - y\| = 0$. Then it follows that $(y|x_1x_2^{\flat}) = \lim_{n \rightarrow \infty} (x_n^{\#}|x_1x_2^{\flat}) = \lim_{n \rightarrow \infty} (x_2x_1^{\flat}|x_n) = 0$ for each $x_1, x_2 \in \mathcal{A}$, which implies that $x \in \mathcal{A} \mapsto x^{\#} \in \mathcal{A}$ is closable. Thus $\mathcal{A}$ is a left Hilbert algebra in $\mathcal{H}$ with the involution $\#$. We next show that the full left Hilbert algebra $\mathcal{A}''$ has a unit u . For any $\varepsilon > 0$ and for any finite subsets $\{x_1, \dots, x_m\}$ and $\{y_1, \dots, y_m\}$ of $\mathcal{A}$, we define the set

$$\begin{aligned} K(\varepsilon; \{x_1, \dots, x_m\}, \{y_1, \dots, y_m\}) \\ = \{a \in \mathcal{H}; \|a\| \leq 1, |(ax_k - x_k|y_k)| \leq \varepsilon \\ \text{and } |(x_ka - x_k|y_k)| \leq \varepsilon, k = 1, \dots, m\}. \end{aligned}$$

Since the C^{*} -algebra $\mathcal{A}_{\#}$ has an approximate identity and $\|x\| \leq \|x\|_{\#}$ for each $x \in \mathcal{A}_{\#}$, it follows that $K(\varepsilon; \{x_1, \dots, x_m\}, \{y_1, \dots, y_m\}) \neq \emptyset$. Now let $\mathcal{K}$ be the family of all subsets $K(\varepsilon; \{x_1, \dots, x_m\}, \{y_1, \dots, y_m\})$ where $\varepsilon > 0$ and $\{x_1, \dots, x_m\}, \{y_1, \dots, y_m\}$ are finite subsets. Then $\mathcal{K}$ is a family of non-empty weakly closed subsets of the weakly compact set $\mathcal{H}_1 \equiv \{a \in \mathcal{H}; \|a\| \leq 1\}$. Hence, the intersection of all the sets in $\mathcal{K}$ is non-empty. Hence, an element u of this intersection is such that u is a *quasi-unit* of the topological quasi $*$ -algebra $\overline{\mathcal{A}}[\|\cdot\|]$, that is, $u \in \overline{\mathcal{A}}[\|\cdot\|]$ and $ux = xu = x$ for each $x \in \mathcal{A}$. Since

$$(S_{\mathcal{A}}x|u) = (x^{\#}|u) = (u|L_xu) = (u|x)$$

for each $x \in \mathcal{A}$, it follows that $u \in \mathcal{D}(S_{\mathcal{A}}^{*})$ and $\pi'_{\mathcal{A}}(u) = I$. Hence, $u \in \mathcal{A}'$ and $S_{\mathcal{A}}^{*}u = u$, which implies that

$$(S_{\mathcal{A}}^{*}y|u) = (\pi'_{\mathcal{A}}(S_{\mathcal{A}}^{*}y)|u) = (u|\pi'_{\mathcal{A}}(y)u) = (u|y)$$

for each $y \in \mathcal{A}'$. Hence, we have $u \in \mathcal{A}''$ and $S_{\mathcal{A}}u = u$. This completes the proof. $\square$

By Proposition 2.4, the situation of HCQ*-algebras can be schematized with the following diagram:

$$\begin{array}{ccccccc} & \subset & \mathcal{A}_{\#} & \subset & \mathcal{A}'' = L''_{\mathcal{A}}u & \subset & \mathcal{D}(S_{\mathcal{A}}) & \subset \\ \mathcal{A} & & \uparrow J & & \uparrow J_{\mathcal{A}} & & \uparrow J_{\mathcal{A}} & \overline{\mathcal{A}}[\|\ \|\] \\ & \subset & \mathcal{A}_{\flat} & \subset & \mathcal{A}' = L'_{\mathcal{A}}u & \subset & \mathcal{D}(S_{\mathcal{A}}^*) & \subset \end{array}$$

We now look for conditions under which $J = J_{\mathcal{A}}$.

Lemma 2.5. *Let $(\overline{\mathcal{A}}[\|\ \|\], \#)$ be a HCQ*-algebra. Then the following statements are equivalent:*

- (i) $J = J_{\mathcal{A}}$.
- (ii) $(x^{\#}|x^*) \geq 0$ for each $x \in \mathcal{A}$.

Proof. (i) $\Rightarrow$ (ii) This follows from

$$(x^{\#}|x^*) = (J_{\mathcal{A}}\Delta_{\mathcal{A}}^{\frac{1}{2}}x|J_{\mathcal{A}}x) = (x|\Delta_{\mathcal{A}}^{\frac{1}{2}}x) \geq 0, \forall x \in \mathcal{A}.$$

(ii) $\Rightarrow$ (i) By the assumption (ii) we have $S_{\mathcal{A}} = J(JJ_{\mathcal{A}}\Delta_{\mathcal{A}}^{\frac{1}{2}})$ and $JJ_{\mathcal{A}}\Delta_{\mathcal{A}}^{\frac{1}{2}} \geq 0$. The uniqueness of the polar decomposition of $S_{\mathcal{A}}$ implies $J = J_{\mathcal{A}}$. $\square$

If any one of the two equivalent statements of Lemma 2.5 holds, we say that the HCQ*-algebra $(\overline{\mathcal{A}}[\|\ \|\], \#)$ is *standard*.

Remark 2.6. Let $(\overline{\mathcal{A}}[\|\ \|\], \#)$ be a HCQ*-algebra. If it is standard, then $R'_{\mathcal{A}} = L''_{\mathcal{A}}$. Conversely, if $R'_{\mathcal{A}} = L''_{\mathcal{A}}$, then $JJ_{\mathcal{A}} = J_{\mathcal{A}}J$, but we don't know whether $J = J_{\mathcal{A}}$.

Since two HCQ*-algebras $(\overline{\mathcal{A}}[\|\ \|\], \#)$ with $(\overline{\mathcal{B}}[\|\ \|\], \#)$, $\overline{\mathcal{A}}[\|\ \|\] = \overline{\mathcal{B}}[\|\ \|\]$ as Hilbert spaces, need not coincide as HCQ*-algebras, we introduce the following notion:

Definition 2.7. A HCQ*-algebra $\overline{\mathcal{A}}[\|\ \|\]$ is said to be an *extension* of a HCQ*-algebra $\overline{\mathcal{B}}[\|\ \|\]$ if $\mathcal{B}$ is a dense *-subalgebra of $\mathcal{A}$ and $S_{\mathcal{A}} = S_{\mathcal{B}}$.

Proposition 2.8. *Let $(\overline{\mathcal{A}}[\|\ \|\], \#)$ be a standard HCQ*-algebra, and $\mathcal{B} \equiv (\mathcal{A}'')_0$ the maximal Tomita algebra of the full left Hilbert algebra $\mathcal{A}''$. Then $(\overline{\mathcal{B}}[\|\ \|\], S_{\mathcal{A}})$ is a standard HCQ*-algebra and it is an extension of $(\overline{\mathcal{A}}[\|\ \|\], S_{\mathcal{A}})$. Further, $\{\Delta_{\mathcal{A}}^{it}\}_{t \in \mathbb{R}}$ is a one-parameter group of *-automorphisms of the Hilbert quasi *-algebra $\overline{\mathcal{B}}[\|\ \|\]$, that is, $\Delta_{\mathcal{A}}^{it}\mathcal{B} = \mathcal{B}$, $(\Delta_{\mathcal{A}}^{it}a)^* = \Delta_{\mathcal{A}}^{it}a^*$, $\Delta_{\mathcal{A}}^{it}(ax) = (\Delta_{\mathcal{A}}^{it}a)(\Delta_{\mathcal{A}}^{it}x)$ and $\Delta_{\mathcal{A}}^{it}(xa) = (\Delta_{\mathcal{A}}^{it}x)(\Delta_{\mathcal{A}}^{it}a)$ for all $a \in \overline{\mathcal{B}}[\|\ \|\]$, $x \in \mathcal{B}$ and $t \in \mathbb{R}$.*

Proof. It is almost clear that $\overline{\mathcal{B}}[\|\ \|\]$ is a Hilbert quasi *-algebra with the involution $J_{\mathcal{A}} = J_{\mathcal{B}}$ and further $(\overline{\mathcal{B}}[\|\ \|\], S_{\mathcal{A}})$ is a standard HCQ*-algebra. Since $\{\Delta_{\mathcal{A}}^{it}\}_{t \in \mathbb{R}}$ is a one-parameter group of *-automorphisms of the Tomita algebra $\mathcal{B}$, it follows that $\{\Delta_{\mathcal{A}}^{it}\}_{t \in \mathbb{R}}$ is also a one-parameter group of *-automorphisms of the Hilbert quasi *-algebra $\overline{\mathcal{B}}[\|\ \|\]$. $\square$

Finally, we consider the question of when a Hilbert space can be regarded as a standard HCQ*-algebra. By Proposition 2.4, 2.8 and [9], Theorem 13.1, we have the following:

Theorem 2.9. *Let $\mathcal{H}$ be a Hilbert space. The following statements are equivalent:*

- (i) $\mathcal{H}$ is a standard HCQ^* -algebra.
- (ii) $\mathcal{H}$ contains a left Hilbert algebra with unit as dense subspace.
- (iii) There exists a von Neumann algebra on $\mathcal{H}$ with a cyclic and separating vector.

It is worth noticing, in particular, that the implication (iii) $\Rightarrow$ (i) shows that the class of standard HCQ^ -algebras is rather rich.*

3. THE STRUCTURE OF STRICT CQ^* -ALGEBRAS

In this section we study when a strict CQ^* -algebra is embedded in a standard HCQ^* -algebra. For that, we need a GNS-like construction for a class of positive sesquilinear forms on strict CQ^* -algebras $(\overline{\mathcal{A}}[\|\cdot\|], \#, \|\cdot\|_{\#})$. A sesquilinear form φ on $\overline{\mathcal{A}}[\|\cdot\|] \times \overline{\mathcal{A}}[\|\cdot\|]$ is said to be *positive* if $\varphi(a, a) \geq 0$ for all $a \in \overline{\mathcal{A}}[\|\cdot\|]$, and φ is said to be *faithful* if $\varphi(a, a) = 0, a \in \overline{\mathcal{A}}[\|\cdot\|]$, implies $a = 0$. Further, we need the following notion:

Definition 3.1. Let $(\overline{\mathcal{A}}[\|\cdot\|], \#, \|\cdot\|_{\#})$ and $(\overline{\mathcal{B}}[\|\cdot\|_1], \#, \|\cdot\|_{\#_1})$ be strict CQ^* -algebras. A linear map $\Phi : \overline{\mathcal{A}}[\|\cdot\|] \rightarrow \overline{\mathcal{B}}[\|\cdot\|_1]$ is said to be a **-homomorphism* of $(\overline{\mathcal{A}}[\|\cdot\|], \#, \|\cdot\|_{\#})$ into $(\overline{\mathcal{B}}[\|\cdot\|_1], \#, \|\cdot\|_{\#_1})$ if (i) Φ is a *-homomorphism of the quasi *-algebra $\overline{\mathcal{A}}[\|\cdot\|]$ into the quasi *-algebra $\overline{\mathcal{B}}[\|\cdot\|_1]$, that is, $\Phi(\mathcal{A}) \subset \mathcal{B}$ and $\Phi(a)^* = \Phi(a^*), \Phi(ax) = \Phi(a)\Phi(x)$ and $\Phi(xa) = \Phi(x)\Phi(a)$ for all $a \in \overline{\mathcal{A}}[\|\cdot\|]$ and $x \in \mathcal{A}$; (ii) $\Phi|_{\mathcal{A}_{\#}}$ is a *-homomorphism of the C^* -algebra $\mathcal{A}_{\#}$ into the C^* -algebra $\mathcal{B}_{\#_1}$. A bijective (resp. injective) *-homomorphism Φ such that $\Phi(\mathcal{A}) = \mathcal{B}$ and $\Phi(\mathcal{A}_{\#}) = \mathcal{B}_{\#_1}$ is called a **-isomorphism* of $(\overline{\mathcal{A}}[\|\cdot\|], \#, \|\cdot\|_{\#})$ onto (resp. into) $(\overline{\mathcal{B}}[\|\cdot\|_1], \#, \|\cdot\|_{\#_1})$. A *-homomorphism Φ is said to be *contractive* if $\|\Phi(a)\|_1 \leq \|a\|$ for all $a \in \overline{\mathcal{A}}[\|\cdot\|]$. A contractive *-isomorphism whose inverse is also contractive is called an *isometric *-isomorphism*.

Theorem 3.2. *Let $(\overline{\mathcal{A}}[\|\cdot\|], \#, \|\cdot\|_{\#})$ be a strict CQ^* -algebra with quasi-unit u . Then the following statements are equivalent:*

- (i) *There exists a contractive *-homomorphism (resp. *-isomorphism) of the strict CQ^* -algebra $(\overline{\mathcal{A}}[\|\cdot\|], \#, \|\cdot\|_{\#})$ into a HCQ^* -algebra $(\overline{\mathcal{B}}[\|\cdot\|_1], \#_1)$.*
- (ii) *There exists a (resp. faithful) positive sesquilinear form φ on $\overline{\mathcal{A}}[\|\cdot\|] \times \overline{\mathcal{A}}[\|\cdot\|]$ satisfying*
 - (ii)₁ $\varphi(x, y) = \varphi(u, x^{\#}y), \forall x, y \in \mathcal{A}$;
 - (ii)₂ $|\varphi(x, y)| \leq \|x\|\|y\|, \forall x, y \in \mathcal{A}$;
 - (ii)₃ $\varphi(x, y) = \varphi(y^*, x^*), \forall x, y \in \mathcal{A}$;*Further, $(\overline{\mathcal{B}}[\|\cdot\|_1], \#_1)$ is standard if and only if*
 - (ii)₄ $\varphi(x^*, x^{\#}) \geq 0, \forall x \in \mathcal{A}$.

Proof. (i) $\Rightarrow$ (ii) We put

$$\varphi(a, b) = (\Phi(a)|\Phi(b)), \quad a, b \in \overline{\mathcal{A}}[\|\cdot\|],$$

where $(\cdot | \cdot)$ is the inner product defined by the Hilbertian norm $\|\cdot\|_1$ on $\overline{\mathcal{B}}[\|\cdot\|_1]$. Then it is easily shown that φ is a positive sesquilinear form on $\overline{\mathcal{A}}[\|\cdot\|] \times \overline{\mathcal{A}}[\|\cdot\|]$ satisfying the condition (ii)₁ $\sim$ (ii)₃. If $(\overline{\mathcal{B}}[\|\cdot\|_1], \#_1)$ is standard, then (ii)₄ follows from Lemma 2.5.

(ii) $\Rightarrow$ (i) We put $\mathcal{N}_{\varphi} = \{a \in \overline{\mathcal{A}}[\|\cdot\|]; \varphi(a, a) = 0\}$. Then $\mathcal{N}_{\varphi}$ is a subspace of $\overline{\mathcal{A}}[\|\cdot\|]$ and, due to the positivity of φ , which implies $\varphi(a, b) = \overline{\varphi(b, a)}$ for each $a, b \in \overline{\mathcal{A}}[\|\cdot\|]$,

it follows that the quotient space $\lambda_\varphi(\overline{\mathcal{A}}[\|\cdot\|]) \equiv \overline{\mathcal{A}}[\|\cdot\|]/\mathcal{N}_\varphi = \{\lambda_\varphi(a) \equiv a + \mathcal{N}_\varphi; a \in \overline{\mathcal{A}}[\|\cdot\|]\}$ is a pre-Hilbert space with inner product $(\lambda_\varphi(a)|\lambda_\varphi(b))_\varphi = \varphi(a, b)$, $a, b \in \overline{\mathcal{A}}[\|\cdot\|]$. We denote by $\|\cdot\|_\varphi$ the norm defined by the inner product $(\cdot|\cdot)_\varphi$ and by $\mathcal{H}_\varphi$ the completion of $\lambda_\varphi(\overline{\mathcal{A}}[\|\cdot\|])[\|\cdot\|_\varphi]$. Since $\mathcal{A}$ is $\|\cdot\|$ -dense in $\overline{\mathcal{A}}[\|\cdot\|]$, it follows that

$$(ii)'_2 \quad |\varphi(a, b)| \leq \|a\| \|b\|, \quad \forall a, b \in \overline{\mathcal{A}}[\|\cdot\|];$$

$$(ii)'_3 \quad \varphi(a, b) = \varphi(b^*, a^*), \quad \forall a, b \in \overline{\mathcal{A}}[\|\cdot\|],$$

and since $(ii)'$ and $\|x\| \leq \|x\|_\#$, $\forall x \in \mathcal{A}$, it follows that

$$(ii)'_1 \quad \varphi(x, y) = \varphi(u, x^\# y), \quad \forall x, y \in \mathcal{A}_\#.$$

By $(ii)'_2$ $\mathcal{A}_\varphi \equiv \lambda_\varphi(\mathcal{A})$ is a dense subspace of the Hilbert space $\mathcal{H}_\varphi$ and further, it is a *-algebra equipped with the multiplication $\lambda_\varphi(x)\lambda_\varphi(y) = L_{\lambda_\varphi(x)}\lambda_\varphi(y) \equiv \lambda_\varphi(xy)$ and the involution $\lambda_\varphi(x)^* \equiv \lambda_\varphi(x^*)$. By $(ii)'_3$ the involution $\lambda_\varphi(x) \rightarrow \lambda_\varphi(x)^*$ can be extended to the isometric involution J_φ on $\mathcal{H}_\varphi$. By $(ii)'_1$ the linear functional on the C*-algebra $\mathcal{A}_\# : x \rightarrow \varphi(x, u)$ is positive, and so $\varphi(y^\#(x^\# x)y, u) \leq \|x\|_\#^2 \varphi(y, y)$ for each $x, y \in \mathcal{A}$. Hence it follows from $(ii)_1$ that

$$\|\lambda_\varphi(x)\lambda_\varphi(y)\|_\varphi^2 = \varphi(xy, xy) = \varphi(y^\# x^\# xy, u) \leq \|x\|_\#^2 \|\lambda_\varphi(y)\|_\varphi^2$$

for each $x, y \in \mathcal{A}$, so that $L_{\lambda_\varphi(x)}$ is bounded and $\|L_{\lambda_\varphi(x)}\| \leq \|x\|_\#$ for each $x \in \mathcal{A}$. Thus $\mathcal{H}_\varphi = \overline{\mathcal{A}_\varphi}[\|\cdot\|_\varphi]$ is a Hilbert quasi *-algebra. Further, the map $\lambda_\varphi(x) \rightarrow \lambda_\varphi(x)^\# \equiv \lambda_\varphi(x^\#)$ is an involution of $\mathcal{A}_\varphi$ and by $(ii)_1$ $L_{\lambda_\varphi(x)}^* = L_{\lambda_\varphi(x)^\#}$ for each $x \in \mathcal{A}$. Hence, $(\overline{\mathcal{A}_\varphi}[\|\cdot\|_\varphi], \#_1)$ is a HCQ*-algebra. Here we put $\Phi(a) = \lambda_\varphi(a)$, $a \in \overline{\mathcal{A}}[\|\cdot\|]$. Then it is easily shown that Φ is a *-homomorphism of the strict CQ*-algebra into the HCQ*-algebra $(\overline{\mathcal{A}_\varphi}[\|\cdot\|_\varphi], \#_1)$ satisfying $\Phi(\mathcal{A}_0) = \mathcal{A}_\varphi$, and by $(ii)'_2$ it is contractive. Suppose that φ is faithful. Then the *-representation of the C*-algebra $\mathcal{A}_\#$ on $\mathcal{H}_\varphi$ defined by $x \rightarrow L_{\lambda_\varphi(x)}$, $x \in \mathcal{A}_\#$ is faithful, which implies that $\|L_{\lambda_\varphi(x)}\| = \|x\|_\#$ for each $x \in \mathcal{A}_\#$. Further, since $\Phi(\mathcal{A}_0) = \mathcal{A}_\varphi$, it follows that $\Phi(\mathcal{A}_\#) = (\mathcal{A}_\varphi)_{\#_1}$ and $\Phi[\mathcal{A}_\#]$ is a *-isomorphism of the C*-algebra $\mathcal{A}_\#$ onto the C*-algebra $(\mathcal{A}_\varphi)_{\#_1}$. Hence Φ is a *-isomorphism of $(\overline{\mathcal{A}}[\|\cdot\|], \#, \|\cdot\|_\#)$ into $(\overline{\mathcal{A}_\varphi}[\|\cdot\|_\varphi], \#_1)$. By Lemma 2.5, the HCQ*-algebra $(\overline{\mathcal{A}_\varphi}[\|\cdot\|_\varphi], \#_1)$ is standard if and only if $(ii)_4$ holds. This completes the proof. $\square$

Now the question arises as to whether positive sesquilinear forms as described in (ii) do really exist. The answer is certainly positive due to the existence of standard HCQ*-algebras stated in Theorem 2.9. Indeed, the inner product $\langle \cdot, \cdot \rangle$ of a left Hilbert algebra satisfies conditions $(ii)_1$ – $(ii)_4$.

Furthermore, Theorem 3.2 answers the main question in this section: any form φ over a strict CQ*-algebra $(\overline{\mathcal{A}}[\|\cdot\|], \#, \|\cdot\|_\#)$ with quasi-unit, can be used to construct a HCQ*-algebra where $\overline{\mathcal{A}}$ is contractively embedded.

ACKNOWLEDGMENT

We acknowledge the financial support of the Gruppo Nazionale per l'Analisi Funzionale e le Applicazioni del C.N.R. and of the Italian Ministry of Scientific Research, and of the Japan Private School Promotion Foundation.

REFERENCES

1. F.Bagarello and C.Trapani, *States and representations of CQ*-algebras*, Ann. Inst. H. Poincaré **61**, 103-133 (1994) MR **95j**:46062
2. F.Bagarello and C.Trapani, *CQ*-algebras: structure properties*, Publ. Res. Inst. Math. Sci., Kyoto Univ. **32**, 85-116 (1996) MR **97d**:46064

3. F.Bagarello and C.Trapani, *L^p -spaces as quasi $*$ -algebras*, J. Math. Anal. Appl. **197**, 810-824 (1996) MR **96k**:46064
4. F.Bagarello and C.Trapani, *The Heisenberg dynamics of spin systems:a quasi $*$ -algebras approach*, J. Math. Phys. **37**, 4219-4234 (1996) MR **97j**:82005
5. I.M.Gelfand and N.Ya.Vilenkin, *Generalized functions Vol. 4*, Academic Press, New York and London, 1964 MR **55**:8786d
6. G.Lassner, *Algebras of unbounded operators and quantum dynamics*, Physica, **124A**, 471-479 (1984) CMP 17:01
7. G.Lassner and G.A.Lassner, *Qu^* -algebras and twisted product*, Publ. Res. Inst. Math. Sci., Kyoto Univ. **25**, 279-299 (1989) MR **90h**:47087
8. S.Stratila and L.Szido, *Lectures on von Neumann Algebras*, Revision of 1975 original, Abacus Press, Tunbridge Wells, 1979 MR **81j**:46089
9. M.Takesaki, *Tomita's theory of modular Hilbert algebras and its applications*, Lecture Notes in Mathematics, **128** Springer-Verlag, 1970 MR **42**:5061
10. C.Trapani, *Quasi $*$ -algebras of operators and their applications*, Rev. Math. Phys. **7** 1303-1332 (1995) MR **97a**:47070
11. A.Van Daele, *A new approach to the Tomita-Takesaki theory of generalized Hilbert algebras*, J. Functional Analysis **15**, 378-393 (1974) MR **49**:11264

DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DI PALERMO, I-90128 PALERMO, ITALY
E-mail address: bagarello@www.unipa.it

DEPARTMENT OF APPLIED MATHEMATICS, FUKUOKA UNIVERSITY, J-814-80 FUKUOKA, JAPAN
E-mail address: a-inoue@fukuoka-u.ac.jp

DIPARTIMENTO DI SCIENZE FISICHE ED ASTRONOMICHE, UNIVERSITÀ DI PALERMO, I-90123 PALERMO, ITALY
E-mail address: trapani@unipa.it