A UNIVERSAL COANALYTIC LINEAR ORDERING

ABHIJIT DASGUPTA

(Communicated by Carl G. Jockusch, Jr.)

ABSTRACT. We construct a Π_1^1 linear ordering in which every Π_1^1 (coanalytic) linear ordering can be order embedded.

1. INTRODUCTION

We say that $(A, <_A)$ is a *Borel (resp. coanalytic) linear ordering* if A is a set of reals, $<_A$ is a linear (total) ordering of A , and both A and $<_A$ are Borel (resp. coanalytic).

By the structure theorem on Borel orderings in [2], given any Borel linear ordering $(A, <_A)$, there is a Borel linear ordering $(B, <_B)$ such that $(A, <_A)$ can be order embedded in $(B, <_B)$, but $(B, <_B)$ cannot be order embedded in $(A, <_A)$. It follows that there is no universal Borel linear ordering with respect to order embeddability.

In this paper we show that among coanalytic linear orderings, on the other hand, there is a universal one:

Theorem 1.1. *There exists a Π_1^1 set P of reals and a Π_1^1 relation $<_P \subseteq P \times P$ which linearly orders P and satisfies the following: if A is a set of reals and $<_A \subseteq A \times A$ is any Π_1^1 relation which linearly orders A , then $(A, <_A)$ is order embeddable in $(P, <_P)$.*

2. PRELIMINARIES

Unless otherwise noted, we use notation, terminology, and results of modern descriptive set theory as can be found in a standard textbook like [3], [4], or [6]. The facts about linear orderings used below can be found in [5]; here we review those which we will need. We also use results from [1] and [2] about Borel orderings.

We use the standard notation of set theory for ordinals: ω , the smallest infinite (countable) ordinal, is the set of all natural numbers (finite ordinals); and ω_1 , the least uncountable ordinal, is the set of all countable ordinals.

A *linear ordering* is an ordered pair $(A, <_A)$ where A is a set and $<_A \subseteq A \times A$ is a relation on A such that $<_A$ is transitive and satisfies the rule of trichotomy on A : For any $x, y \in A$, exactly one of the conditions $x = y$, $x <_A y$, and $y <_A x$ holds. In this case we say that $<_A$ linearly orders A .

Received by the editors November 22, 1999 and, in revised form, May 1, 2000.

2000 *Mathematics Subject Classification.* Primary 03E15, 04A15; Secondary 06A05.

Key words and phrases. Descriptive set theory, coanalytic sets, total order, linear order.

Given linear orderings $(A, <_A)$, $(B, <_B)$ and a function $f: A \rightarrow B$, we say that f is (strictly) *increasing* or *order preserving* if for all $x, y \in A$, if $x <_A y$, then $f(x) <_B f(y)$. We say that $(A, <_A)$ is *embeddable* in $(B, <_B)$ if there is an order preserving $f: A \rightarrow B$.

If $(A, <_A)$ is a linear ordering and $B, C \subseteq A$, we write $B <_A C$ to mean $x <_A y$ for all $x \in B$ and $y \in C$.

A linear ordering is *short* if it does not contain any increasing or decreasing ω_1 -chain.

If ξ is an ordinal and $(A, <_A)$ is a linear ordering, then we define the ξ -fold *lexicographic power* of $(A, <_A)$ as the ordering on the set A^ξ (of all ξ -sequences of members of A) where for $x, y \in A^\xi$, x precedes y iff there exists $\alpha < \xi$ such that $x(\alpha) <_A y(\alpha)$ and for all $\beta < \alpha$, $x(\beta) = y(\beta)$; it is easily verified that this indeed defines a linear ordering. In particular, $2 = \{0, 1\}$ is ordered naturally, and 2^ξ denotes its ξ -fold lexicographic power. Let $2^{<\omega_1}$ denote the subordering of 2^{ω_1} consisting of those elements x of 2^{ω_1} for which there is $\zeta < \omega_1$ with $x(\zeta) = 1$ and $x(\beta) = 0$ for all β with $\zeta < \beta < \omega_1$ (i.e. binary ω_1 -sequences with a last 1). $(\omega^\omega, <_\mathbb{N})$ denotes the ω -fold lexicographic power of $(\omega, <)$.

A linear ordering $(A, <_A)$ is an η_1 -ordering if for any countable subsets $B, C \subseteq A$ with $B <_A C$, there exists $x \in A$ with $B <_A \{x\} <_A C$.

A linear ordering $(A, <_A)$ is *quasi-short* if there are subsets $A_\alpha \subseteq A$ for $\alpha < \omega_1$ such that $A = \bigcup \{A_\alpha \mid \alpha < \omega_1\}$, and for each $\alpha < \omega_1$, the restriction $<_A \upharpoonright A_\alpha$ of $<_A$ to A_α is short.

Lemma 2.1. 1. *A subordering of a short linear ordering is short.*

2. *The union of countably many short linear orderings is short. More precisely, if $(A, <_A)$ is a linear ordering with $A = \bigcup \{A_n \mid n \in \omega\}$ and $(A_n, <_A \upharpoonright A_n)$ is short for each n , then $(A, <_A)$ is short.*
3. *Suppose that $(A, <_A)$ is a linear ordering, $B \subseteq A$ is a short subordering, and whenever $x, y, z \in A$ and $x <_A y <_A z$, there is $w \in B$ with $x <_A w <_A z$. Then $(A, <_A)$ is short.*
4. *If ξ is a countable ordinal, then the ξ -fold lexicographic power of a short linear ordering is short. In particular, 2^ξ is short for countable ξ .*
5. *$2^{<\omega_1}$ is an η_1 -ordering.*

Proof. The first three statements are obvious, and the fourth is verified by induction on ξ using the second and third claims. The last statement is an immediate consequence of Theorem 9.24 on page 165 of [5]. $\square$

3. EMBEDDING QUASI-SHORT LINEAR ORDERINGS INTO η_1 -ORDERINGS

Theorem 3.1. *Every quasi-short linear ordering can be embedded into any η_1 -ordering.*

Proof. This follows from the following lemma.

Lemma 3.2 (Hausdorff). *If $(A, <_A)$ is a short linear ordering, $(B, <_B)$ is an η_1 -ordering, $S \subseteq A$, and $f: S \rightarrow B$ is strictly increasing, then f can be extended to a strictly increasing map from all of A into B .*

Proof. Let $\mathcal{F}$ be the family of all strictly increasing functions extending f (mapping a subset of A into B). Partially order $\mathcal{F}$ under extension. Then by Zorn's lemma, there is a maximal member g of $\mathcal{F}$. We claim that $\text{dom}(g) = A$. Otherwise there

would exist $a \in A \setminus \text{dom}(g)$. Let $L = \{x \in \text{dom}(g) \mid x <_A a\}$ and $R = \{x \in \text{dom}(g) \mid a <_A x\}$. Since $(A, <_A)$ is short, there exist countable sets P and Q with $P \subseteq L$ and $Q \subseteq R$ such that: (a) for all $x \in L$, there is $y \in P$ with $x <_A y$ or $x = y$; and (b) for all $x \in R$, there is $y \in Q$ with $y <_A x$ or $x = y$. Put $C = \{g(x) \mid x \in P\}$ and $D = \{g(x) \mid x \in Q\}$. Then C and D are countable subsets of B with $C <_B D$. Since $(B, <_B)$ is an η_1 -ordering, there is $b \in B$ with $C <_B \{b\} <_B D$. Define an extension h of g with $\text{dom}(h) = \text{dom}(g) \cup \{a\}$ by setting $h(a) = b$. Then h is a strictly increasing proper extension of g , contradicting the maximality of g . $\square$

To finish the proof of the theorem, let $(A, <_A)$ be a quasi-short linear ordering, and let $A_\alpha \subseteq A$ for $\alpha < \omega_1$ be such that $A = \bigcup \{A_\alpha \mid \alpha < \omega_1\}$, and for each $\alpha < \omega_1$, the restriction $<_A \upharpoonright A_\alpha$ of $<_A$ to A_α is short. Without loss of generality, we assume that the sets A_α increase with α . Now let $(B, <_B)$ be any η_1 -ordering. Using Lemma 3.2 and transfinite induction, we can build strictly increasing functions $f_\alpha: A_\alpha \rightarrow B$, $\alpha < \omega_1$, such that if $\alpha < \beta < \omega_1$, then f_β extends f_α . Then the common extension of all the functions f_α is a strictly increasing map from $(A, <_A)$ to $(B, <_B)$. $\square$

4. COANALYTIC ORDERINGS ARE QUASI-SHORT

We will use the fact that Borel linear orderings are short ([1], [2]) in the proof of the following theorem.

Theorem 4.1. *Let A be a set of reals and $<_A$ be a linear ordering of A such that $<_A$, as a subset of the plane, is Π_1^1 (coanalytic). Then $(A, <_A)$ is quasi-short.*

Proof. We assume that A is uncountable. A is Σ_2^1 , since $<_A$ is Π_1^1 and $x \in A$ iff $\exists y(x <_A y \vee y <_A x)$. So we can express A as a union of ω_1 Borel sets:

$$A = \bigcup_{\xi < \omega_1} A_\xi$$

where each A_ξ is a Borel set. For each $\xi < \omega_1$, let

$$\begin{aligned} <_\xi &\stackrel{\text{def}}{=} \{(x, y) \in A_\xi \times A_\xi \mid x <_A y\}, \\ >_\xi &\stackrel{\text{def}}{=} \{(x, y) \in A_\xi \times A_\xi \mid y <_A x\}, \\ =_\xi &\stackrel{\text{def}}{=} \{(x, y) \in A_\xi \times A_\xi \mid x = y\}. \end{aligned}$$

Since linear orderings satisfy the rule of trichotomy, the three Π_1^1 sets $<_\xi$, $>_\xi$, and $=_\xi$ form a partition of the Borel set $A_\xi \times A_\xi$. So by Suslin's theorem, $<_\xi$, $>_\xi$, and $=_\xi$ are all Borel. So $(A_\xi, <_\xi)$ is a Borel linear ordering, and therefore short, for each $\xi < \omega_1$. Hence $(A, <_A)$ is quasi-short. $\square$

5. A Π_1^1 ORDERING CONTAINING AN η_1 -ORDERING

Theorem 5.1. *There exist a Π_1^1 set P of reals and a Π_1^1 linear ordering $<_P$ of P such that $2^{<\omega_1}$ can be embedded in $(P, <_P)$.*

Proof. Throughout this proof we use terminology and results from [4]. We work in the space of irrationals $\mathcal{N} = \omega^\omega$. For $i, j \in \omega$, let $\langle i, j \rangle \in \omega$ be defined as $\langle i, j \rangle = 2^i(2j + 1) - 1$. For $x \in \omega^\omega$, and $i \in \omega$, define $(x)_i \in \omega^\omega$ by

$$(x)_i(n) = x(\langle i, n \rangle) \quad (n \in \omega).$$

For $x \in \omega^\omega$ and $n \in \omega$, let

$$\begin{aligned}
 \leq_x &\stackrel{\text{def}}{=} \{(i, j) \mid x(\langle i, j \rangle) = 1\}, \\
 <_x &\stackrel{\text{def}}{=} \{(i, j) \mid i \leq_x j \wedge i \neq j\}, \\
 \Phi[x] &\stackrel{\text{def}}{=} \text{the field of } \leq_x \\
 &= \{i \mid \exists j(i \leq_x j \vee j \leq_x i)\}, \\
 \Omega[x] &\stackrel{\text{def}}{=} \text{the subset of } \omega \text{ coded by } x \\
 &= \{i \mid x(i) = 1\}, \\
 <_x[n] &\stackrel{\text{def}}{=} \text{the } <_x\text{-predecessors of } n \\
 &= \{i \mid i <_x n\}.
 \end{aligned}$$

The following relations on ω^ω are arithmetical:

$$\begin{aligned}
 \text{LO}(x) &\stackrel{\text{def}}{\longleftrightarrow} x \text{ codes a linear ordering} \\
 &\longleftrightarrow (\forall n \in \Phi[x])(n \leq_x n) \\
 &\quad \wedge (\forall m, n)((m \leq_x n \wedge n \leq_x m) \longrightarrow m = n) \\
 &\quad \wedge (\forall m, n, k)((m \leq_x n \wedge n \leq_x k) \longrightarrow m \leq_x k) \\
 &\quad \wedge (\forall m, n \in \Phi[x])(m \leq_x n \vee n \leq_x m), \\
 \text{ISO}(f, x, y) &\stackrel{\text{def}}{\longleftrightarrow} \text{LO}(x) \text{ and } \text{LO}(y) \text{ and } (f \upharpoonright \Phi[x] \text{ maps } \Phi[x] \\
 &\quad \text{bijectively onto an initial segment of } <_y \text{ preserving} \\
 &\quad \text{order from } <_x \text{ to } <_y \text{ and } f = 0 \text{ outside } \Phi[x]) \text{ or} \\
 &\quad (\exists n \in \Phi[x])(f \upharpoonright <_x[n] \text{ maps } <_x[n] \text{ bijectively} \\
 &\quad \text{onto } \Phi[y] \text{ preserving order from } <_x \text{ to } <_y \text{ and } f = 0 \\
 &\quad \text{outside } <_x[n]).
 \end{aligned}$$

So if $\text{LO}(x)$, then $<_x$ is a linear ordering of a subset of ω . If $\text{LO}(x)$ and $n \in \omega$, then let $|x|$ denote the order-type of $<_x$, and $|x|_{<_n}$ the order-type of $<_x[n]$. Let

$$\text{WO}(x) \stackrel{\text{def}}{\longleftrightarrow} \text{LO}(x) \text{ and } <_x \text{ is a well-ordering.}$$

Then WO is Π_1^1 , and so are the relations “ $|x| = |y|$ ” and “ $|x| < |y|$ ” defined on WO . Note that

$$\text{WO}(x) \wedge \text{WO}(y) \longrightarrow (\exists! f)(\text{ISO}(f, x, y)).$$

We now define the Π_1^1 set P by

$$\begin{aligned}
 P(x) &\stackrel{\text{def}}{\longleftrightarrow} (x)_0 \text{ codes a successor ordinal and } (x)_1 \text{ codes} \\
 &\quad \text{a subset of } \Phi[(x)_0] \text{ “containing” } \sup |(x)_0| \\
 &\longleftrightarrow \text{WO}((x)_0) \wedge \Omega[(x)_1] \subseteq \Phi[(x)_0] \\
 &\quad \wedge (\exists n \in \Phi[(x)_0])(n \in \Omega[(x)_1] \wedge (\forall i \in \Phi[(x)_0])(i \leq_x n)).
 \end{aligned}$$

Notice that P codes $2^{\aleph_{\omega_1}}$, i.e. there is a natural surjection from P onto $2^{\aleph_{\omega_1}}$: If $x \in P$, then $(x)_0$ codes a countable successor ordinal $|x|_{<_0}$ with greatest element $\sup |(x)_0|$, and $(x)_1$ codes a subset of $|x|_{<_0}$ containing $\sup |(x)_0|$. So for $x \in P$, we can define $a_x \in 2^{\omega_1}$ by

$$a_x(\zeta) = \begin{cases} 1 & \text{if } (\exists n)(\zeta = |x|_{<_0} \wedge (x)_1(n) = 1), \\ 0 & \text{otherwise,} \end{cases} \quad (\zeta < \omega_1).$$

Define a linear ordering $<_P$ on P by

$$x <_P y \stackrel{\text{def}}{\longleftrightarrow} a_x < a_y \text{ in } 2^{\aleph_{\omega_1}} \vee (a_x = a_y \wedge x <_{\mathcal{N}} y).$$

Recall that $<_{\mathcal{N}}$ is the ω -fold lexicographic power of $(\omega, <)$ (here, in place of $<_{\mathcal{N}}$, any arithmetical linear ordering of ω^ω will do).

Note that by definition of $<_P$, $2^{\aleph_{\omega_1}}$ can be embedded in $(P, <_P)$.

It remains to show that $<_P$ is Π_1^1 . First:

$$\begin{aligned} a_x < a_y &\longleftrightarrow P(x) \wedge P(y) \wedge (\forall f)(\text{ISO}(f, (x)_0, (y)_0) \\ &\longrightarrow [(\exists n \in \Phi[(x)_0])((x)_1(n) \neq 1 \wedge (y)_1(f(n)) = 1 \wedge \\ &\quad (\forall i \in \Phi[(x)_0])(i <_x n \longrightarrow ((x)_1(i) = 1 \leftrightarrow (y)_1(f(i)) = 1)))] \\ &\quad \vee [(\forall i \in \Phi[(x)_0])((x)_1(i) = 1 \leftrightarrow (y)_1(f(i)) = 1) \\ &\quad \wedge |(x)_0| < |(y)_0|]). \end{aligned}$$

Next,

$$\begin{aligned} a_x = a_y &\longleftrightarrow P(x) \wedge P(y) \wedge |(x)_0| = |(y)_0| \wedge (\forall f)(\text{ISO}(f, (x)_0, (y)_0) \\ &\longrightarrow (\forall i \in \Phi[(x)_0])((x)_1(i) = 1 \leftrightarrow (y)_1(f(i)) = 1)). \end{aligned}$$

Finally, $<_{\mathcal{N}}$ is an arithmetical linear ordering of ω^ω .

It follows that $<_P$ is Π_1^1 . □

Theorem 1.1 is now immediate from Theorem 4.1, Theorem 3.1, and Theorem 5.1.

6. REMARKS

Let us call a linear ordering *very short* if it can be embedded in 2^α for some countable ordinal α . Thus very short linear orderings are short.

Harrington and Shelah in [1] first proved that Borel linear orderings are short. Harrington, Marker, and Shelah in [2] show that Borel linear orderings are very short. Since for each countable ordinal α there is a Borel linear ordering order isomorphic to 2^α , it also follows that there is no universal Borel linear ordering. Note that the authors in [2] studied *pre-orderings* (i.e. reflexive transitive relations), whereas here we looked only at (strict) *linear orderings* (i.e. relations which are asymmetric, transitive, and satisfy the rule of trichotomy).

On the other hand, there are short coanalytic linear orderings which are not very short. To see this, define an ordering $<_S$ on P :

$$x <_S y \stackrel{\text{def}}{\longleftrightarrow} P(x) \wedge P(y) \wedge ((x)_0 <_{\mathcal{N}} (y)_0 \vee ((x)_0 = (y)_0 \wedge a_x < a_y)).$$

REFERENCES

1. L. Harrington and S. Shelah, *Counting equivalence classes for co- κ -Souslin relations*, Logic Colloquium 1980, Eds. D. Van Dalen, D. Lascar, and T. J. Smiley, North-Holland, 1982. MR **84c**:03088
 2. L. Harrington, D. Marker, and S. Shelah, *Borel Orderings*, Trans. Amer. Math. Soc. **310**, 1 (1988), 293–302. MR **90c**:03041
 3. A. S. Kechris, *Classical Descriptive Set Theory*, Springer-Verlag, 1994.
 4. Y. N. Moschovakis, *Descriptive Set Theory*, North-Holland, 1980. MR **82e**:03002
 5. J. G. Rosenstein, *Linear Orderings*, Academic Press, 1982. MR **84m**:06001
 6. S. M. Srivastava, *A Course on Borel Sets*, Springer-Verlag, 1998. MR **99d**:04002
- E-mail address:* takdoom@yahoo.com