NEW PROOF OF TWO BERSTEIN-HILTON THEOREMS

CARMEN ELVIRA AND JOSE L. NAVARRO

(Communicated by Ralph L. Cohen)

ABSTRACT. With arguments of Homotopy Theory and without the assumption of finite type, we give short proofs of two classic theorems which deal with the co- H -type of a suspension and the homotopy class of a suspension map.

1. INTRODUCTION

In this paper we work in the based category of spaces having the homotopy type of CW -complexes with compactly generated topologies, so $\dim X \leq n$ means that there is an n -dimensional CW -complex in the homotopy type of X . The flat product $A \flat B$ is the homotopy fibre of the inclusion map $j : A \vee B \longrightarrow A \times B$. Clearly, $\zeta = M(\Omega i_1 \times \Omega i_2)$ is an explicit coretraction for Ωj , where M denotes the loop multiplication on $\Omega(A \vee B)$. In the sequel we shall use the same symbol for a map and its homotopy class, thus we mark as equal homotopic maps. The class $[*]$ of the nullhomotopic maps will be denoted by 0.

Since $(\Omega j)(1 - \zeta(\Omega j)) = 0$, there is a map $\xi : \Omega(A \vee B) \longrightarrow \Omega(A \flat B)$, unique up to homotopy, such that $(\Omega \pi)\xi = 1 - \zeta(\Omega j)$ and the map $J(f) : \Omega X \longrightarrow \Omega(A \flat B)$ given by the composition $J(f) = \xi(\Omega f)$ is called the co-Hopf construction associated to the map $f : X \longrightarrow A \vee B$.

A comultiplication on a space X is a map $\mu : X \longrightarrow X \vee X$ such that $j\mu = \Delta$, and we say that μ is homotopy associative if $(\mu \vee 1)\mu = (1 \vee \mu)\mu$. We denote by σ the standard comultiplication on any suspension space. A map $f : (X, \mu) \longrightarrow (Y, \nu)$ is a co- H -map, with respect to μ and ν , if $\nu f = (f \vee f)\mu$ or, equivalently, if $(i_1 + i_2)f = i_1 f + i_2 f$ in $[X, Y \vee Y]$. Plainly, the suspension Σg of any map $g : A \longrightarrow B$ is a co- H -map, with respect to the standard comultiplications.

With arguments of Homology Theory, and restricted to CW -complexes of finite type (i.e. CW -complexes with finitely generated homology groups), Bernstein and Hilton proved in [1] the two following results:

Theorem A. *Let (X, μ) be an $(n - 1)$ -connected co- H -space, $n \geq 2$, such that $\dim X \leq 3n - 3$. Then X has the co- H -type of a suspension.*

Theorem B. *Let B be an $(n - 1)$ -connected space, $n \geq 1$, and let A be such that $\dim A \leq 3n - 2$. Then every co- H -map $f : \Sigma A \longrightarrow \Sigma B$ is homotopic to a suspension map.*

Received by the editors March 20, 2000.

2000 *Mathematics Subject Classification.* Primary 55P40, 55P45; Secondary 55P30.

Key words and phrases. Co- H -space, suspension, coprojective plane.

The first author was partially supported by DGS, project PD96-0740.

In the introduction of [1] the authors suggest that a simpler proof of their results should be possible. In this paper we give short proofs of these theorems in the same spirit as the original proofs, but with arguments of Homotopy Theory that make unnecessary the assumption of finite type.

2. THE COPROJECTIVE PLANE OF A CO- H -SPACE

Let (X, μ) be a co- H -space. In [4] Hilton defined the coprojective plane XC_2 of X as the homotopy fibre of the co-Hopf construction $J(\mu)$. Let $f : (X, \mu) \rightarrow (Y, \nu)$ be a co- H -map. Since the co-Hopf construction is natural, the right square D in the following diagram is homotopy commutative:

$$\begin{array}{ccccc} XC_2 & \xrightarrow{p} & \Omega X & \xrightarrow{J(\mu)} & \Omega(X \wr X) \\ & & \downarrow \Omega f & D & \downarrow \Omega(f \wr f) \\ YC_2 & \xrightarrow{p} & \Omega Y & \xrightarrow{J(\nu)} & \Omega(Y \wr Y) \end{array}$$

Therefore $J(\nu)(\Omega f)p = 0$ or, equivalently, there is a map $f_2 : XC_2 \rightarrow YC_2$ such that $pf_2 = (\Omega f)p$. In particular, the coprojective plane is an invariant of the co- H -type in the 1-connected CW -type.

Composition of $(\Omega\pi)\xi = 1 - \zeta(\Omega j)$ with $\Omega\mu$ yields $(\Omega\pi)J(\mu) = \Omega\mu - \zeta(\Omega j)(\Omega\mu) = \Omega\mu - \zeta(\Omega\Delta) = \Omega(i_1 + i_2) - (\Omega i_1 + \Omega i_2)$. Let $f : (\Sigma A, \sigma) \rightarrow (X, \mu)$ be a map. Then

$$\begin{aligned} (\Omega\pi)J(\mu)(\Omega f)e &= \Omega((i_1 + i_2)f)e - (\Omega(i_1 f)e + \Omega(i_2 f)e) \\ &= \mathcal{A}((i_1 + i_2)f - (i_1 f + i_2 f)), \end{aligned}$$

where $\mathcal{A}$ is the adjointness isomorphism and $e : A \rightarrow \Omega\Sigma A$ is the unit. Since $(\Omega\pi)_*$ is a monomorphism, one has the following result which appears in [4] without proof.

Proposition 2.1. *A map $f : \Sigma A \rightarrow X$ is a co- H -map, with respect to σ and μ , if and only if $J(\mu)\mathcal{A}(f) = 0$.*

In other words, f is a co- H -map if and only if its adjoint $\mathcal{A}(f)$ factors through XC_2 . Obviously, $p : XC_2 \rightarrow \Omega X$ factors through XC_2 and, since $\mathcal{A}^{-1}(p) = \varepsilon(\Sigma p)$, one has

Corollary 2.2. *The composition $\varepsilon(\Sigma p)$ is a co- H -map.*

There is a triple $\Omega X \xrightarrow{J(\mu)} \Omega(X \wr X) \xrightarrow{q} X$, where $q = 0$, which is a word-by-word dual of the Sugawara fibration. However, Hilton proved that this triple is not a cofibration. Consider the commutative diagram

$$\begin{array}{ccccccc} XC_2 & \xrightarrow{p} & \Omega X & \xrightarrow{\quad} & C_p & \xrightarrow{q} & \Sigma(XC_2) \\ & & \searrow J(\mu) & & \downarrow \phi & & \downarrow \zeta \\ & & & & \Omega(X \wr X) & \xrightarrow{j} & Q \end{array}$$

where Q is the homotopy cofibre of $J(\mu)$. If X is $(n-1)$ -connected, $n \geq 2$, then $\Omega(X \wr X)$ is $(2n-3)$ -connected, and $J(\mu)$ is an $(n-1)$ -equivalence. By a result of Ganea ([2], Theorem 1.1), the extension ϕ of $J(\mu)$ is a $(3n-3)$ -equivalence and

likewise for ζ , by the standard 5-lemma argument. Let $k : Q \rightarrow \Sigma\Omega X$ be the next map in the Puppe sequence of $J(\mu)$.

$$\begin{array}{ccc} \Sigma(XC_2) & \xrightarrow{\Sigma p} & \Sigma\Omega X \\ \downarrow \zeta & \nearrow k & \downarrow \varepsilon \\ Q & \dashrightarrow & X \end{array}$$

Then $k\zeta = \Sigma p$ and, if $h : Q \rightarrow X$ is defined by composition $h = \varepsilon k$, Theorem 3.4 in [5] implies that h is a $(3n-3)$ -equivalence. Since $h\zeta = \varepsilon k\zeta = \varepsilon(\Sigma p)$, we then have the following.

Proposition 2.3. *Let X be an $(n-1)$ -connected co- H -space, $n \geq 2$. Then the composition $\varepsilon(\Sigma p)$ is a $(3n-3)$ -equivalence.*

3. PROOF OF THEOREMS A AND B

Let X be a 0-connected CW -complex and let $\phi : \tilde{H}_n(X) \rightarrow F$ be an epimorphism of $\tilde{H}_n(X)$ into the free abelian group F , where $n \geq 2$. Then there is a subcomplex $X(n, F)$ of X with the same $(n-1)$ -skeleton of X , and such that

- (1) $u_* : \tilde{H}_q(X(n, F)) \rightarrow \tilde{H}_q(X)$ is an isomorphism for every $q < n$,
- (2) $\phi u_* : \tilde{H}_n(X(n, F)) \rightarrow F$ is an isomorphism,
- (3) $\tilde{H}_q(X(n, F)) = 0$ for every $q > n$,

where $u : X(n, F) \rightarrow X$ denotes the inclusion map.

Note that, if $F = 0$ is the trivial group, then $X(n, 0) = X(n-1)$ is the homology $(n-1)$ -section of X . This (modified) homology decomposition was given in [1].

Assume that (X, μ) is an $(n-1)$ -connected co- H -space, $n \geq 2$, such that $\dim X \leq 3n-3$, and let Y be a CW -complex in the homotopy type of XC_2 . If we denote $W = Y(3n-4, F)$ and $u : W \rightarrow Y$ is the inclusion map, $F = \tilde{H}_{3n-3}(X)$ being a free abelian group, then Corollary 2.2 and Proposition 2.3 imply that the composition

$$\Sigma W \xrightarrow{\Sigma u} \Sigma Y \simeq \Sigma(XC_2) \xrightarrow{\Sigma p} \Sigma\Omega X \xrightarrow{\varepsilon} X$$

is a co- H -equivalence and Theorem A is proved.

Now, let $f : \Sigma A \rightarrow \Sigma B$ be a co- H -map with respect to σ_A and σ_B . Then it follows that $J(\sigma_B)(\Omega f)e_A = 0$, and so there is a map $u : A \rightarrow (\Sigma B)C_2$ such that $pu = (\Omega f)e_A$.

$$\begin{array}{ccccc} A & \xrightarrow{e_A} & \Omega\Sigma A & \xrightarrow{J(\sigma_A)} & \Omega(\Sigma A \flat \Sigma A) \\ & \searrow u & \downarrow \Omega f & & \downarrow \Omega(f \flat f) \\ g \downarrow & & (\Sigma B)C_2 & \xrightarrow{p} & \Omega\Sigma B \\ & \nearrow \phi & & & \\ B & \xrightarrow{e_B} & \Omega\Sigma B & \xrightarrow{J(\sigma_B)} & \Omega(\Sigma B \flat \Sigma B) \end{array}$$

On the other hand, since $1 : \Sigma B \rightarrow \Sigma B$ is obviously a co- H -map, Proposition 2.1 implies that $J(\sigma_B)e_B = 0$. Then there is a map $\phi : B \rightarrow (\Sigma B)C_2$ such that $p\phi = e_B$. Assume that B is $(n-1)$ -connected; then ΣB is n -connected and, by

applying Proposition 2.3 to ΣB , we infer that $\varepsilon(\Sigma p)$ is a $3n$ -equivalence. Note that $\varepsilon(\Sigma p)(\Sigma \phi) = \varepsilon(\Sigma e_B) = 1$, so $\Sigma \phi$ is a coretraction of $\varepsilon(\Sigma p)$. Thus, $\Sigma \phi$ is a $(3n - 1)$ -equivalence, and so ϕ is a $(3n - 2)$ -equivalence. Hence, the induced map $\phi_* : [A, B] \longrightarrow [A, (\Sigma B)C_2]$ is bijective if $\dim A < 3n - 2$, and surjective if $\dim A = 3n - 2$. Therefore, provided $\dim A \leq 3n - 2$, there is a map $g : A \longrightarrow B$ such that $\phi g = u$. Thus, $e_B g = p \phi g = pu = (\Omega f)e_A = \mathcal{A}(f)$ and, by taking adjoints, we have $f = \mathcal{A}^{-1}(e_B g) = \Sigma g$. This proves Theorem B.

4. REMARK

It has been known for a long time that any $(n - 1)$ -connected space such that $\dim X \leq 2n - 1$, $n \geq 1$, has the homotopy type of a suspension. A proof, by induction on the geometrical dimension of a CW -complex, was given in [3]. After the modified homology decomposition of Bernstein-Hilton it is possible to give a shorter proof. Indeed, if $n \geq 2$ the hypotheses imply that $\varepsilon : \Sigma \Omega X \longrightarrow X$ is a $(2n - 1)$ -equivalence, and that $F = \tilde{H}_{2n-1}(X)$ is a free abelian group, respectively. Let Y be a CW -complex in the homotopy type of ΩX . If $W = Y(2n - 2, F)$ and $u : W \longrightarrow Y$ is the inclusion map, then the composition

$$\Sigma W \xrightarrow{\Sigma u} \Sigma Y \simeq \Sigma \Omega X \xrightarrow{\varepsilon} X$$

is a homotopy equivalence in the 1-connected CW -type. If $n = 1$, note that $\dim X \leq 1$ implies that X is contractible or X has the homotopy type of a wedge of 1-spheres, so the suspension of a discrete space.

REFERENCES

1. I. Bernstein and P. Hilton, *On suspensions and comultiplications*, *Topology* **2** (1963), 73–82 MR **27**:762
2. T. Ganea, *A generalization of the homology and homotopy suspension*, *Comment. Math. Helv.* **39** (1965), 295–322 MR **31**:4033
3. P. Hilton, *On divisors and multiples of continuous maps*, *Fundamenta Math.* **43** (1956), 358–386 MR **18**:814b
4. P. Hilton, *Note on a theorem of Stasheff*, *Bull. Polish Acad. Sci. Math.* **10** (1962), 127–131 MR **25**:4536
5. M. Walker, *Homotopy pullbacks and applications to duality*, *Canad. J. Math.* **29** (1977), 45–64 MR **55**:11245

DEPARTAMENTO DE ANALISIS ECONOMICO, UNIVERSIDAD DE ZARAGOZA, 50005-ZARAGOZA, SPAIN

E-mail address: celvira@posta.unizar.es

DEPARTAMENTO DE MATEMATICAS, UNIVERSIDAD DE ZARAGOZA, 50009-ZARAGOZA, SPAIN

E-mail address: jlnava@posta.unizar.es