A GLOBAL PINCHING THEOREM FOR SURFACES WITH CONSTANT MEAN CURVATURE IN S^3

YI-JUNG HSU AND TAI-HO WANG

(Communicated by Christopher Croke)

ABSTRACT. Let M be a compact immersed surface in the unit sphere S^3 with constant mean curvature H . Denote by ϕ the linear map from $T_p(M)$ into $T_p(M)$, $\phi = A - \frac{H}{2}I$, where A is the linear map associated to the second fundamental form and I is the identity map. Let Φ denote the square of the length of ϕ . We prove that if $\|\Phi\|_{L^2} \leq C$, then M is either totally umbilical or an $H(r)$ -torus, where C is a constant depending only on the mean curvature H .

1. INTRODUCTION

Let M be a compact immersed hypersurface in the unit sphere S^{n+1} with constant mean curvature H . Denote by $h = [h_{ij}]$ the second fundamental form of M and by ϕ the tensor $\phi_{ij} = h_{ij} - \frac{H}{n}\delta_{ij}$. Let Φ denote the square of the length of ϕ . It is well known that if $H = 0$ and $0 \leq \Phi \leq n$, then M is either the equatorial sphere or a Clifford torus [3]. Recently, H. Alencar and M. do Carmo extended the above result to a hypersurface M with constant mean curvature H [1]. They proved that M is either totally umbilical or an $H(r)$ -torus if Φ satisfies a certain pointwise pinching condition. In 1989, C. L. Shen proved that a minimal hypersurface M is totally geodesic if M is of nonnegative sectional curvature, and Φ satisfies a certain global pinching condition [8]. Later, the first author improved a result of Shen in the case of $n = 2$ and found a sharp bound concerning the global pinching condition [6]. The purpose of this paper is to extend our global theorem to a surface M with constant mean curvature H and obtain the best constant.

Before stating our main result, let B be the constant $B = 2 + \frac{H^2}{2}$ and $m(B)$ be the maximum value of the function $q(x) = 2\sqrt{2} \frac{\sqrt{B}x^2 + 2(B-2)x + B\sqrt{B}}{(\sqrt{B}+x)^2((x^2-B)^2 + 8x^2)}$ on $[0, \infty)$. The following is our main result.

Theorem 1.1. *Let M be a compact immersed surface in the unit sphere S^3 with constant mean curvature H . Then*

$$\|\Phi\|_2 \geq 2\pi \sqrt{\frac{2g}{M(B)}},$$

Received by the editors April 17, 1997 and, in revised form, May 10, 2000.

2000 *Mathematics Subject Classification.* Primary 53C40, 53C42.

Key words and phrases. Mean curvature, sphere, totally umbilical.

where g is the genus of M and $\|\cdot\|_2$ is the L^2 -norm. The equality holds if and only if M is either totally umbilical or an $H(r)$ -torus. In particular, if $\|\Phi\|_2 \leq 2\pi\sqrt{\frac{2}{m(B)}}$, then M is either totally umbilical or an $H(r)$ -torus.

It should be noted that for M to be either totally umbilical or an $H(r)$ -torus, $\|\Phi\|_2 = 2\pi\sqrt{2g(\frac{B}{2})^{\frac{3}{4}}}$. It turns out that there exists a constant $H_0 \geq 2$ such that $m(B) = (\frac{2}{B})^{\frac{3}{2}}$ for all $|H| \leq H_0$. As a immediate consequence of the above result, we state

Corollary 1.2. *Let M be a compact immersed surface in the unit sphere S^3 with constant mean curvature H , $|H| \leq H_0$. Then*

$$\|\Phi\|_2 \geq 2\pi\sqrt{2g(\frac{B}{2})^{\frac{3}{4}}},$$

where g is the genus of M and $\|\cdot\|_2$ is the L^2 -norm. The equality holds if and only if M is either totally umbilical or an $H(r)$ -torus. In particular, if $\|\Phi\|_2 \leq 2\pi\sqrt{2(\frac{B}{2})^{\frac{3}{4}}}$, then M is either totally umbilical or an $H(r)$ -torus.

For the proof of the main theorem, we shall need the following Bernstein-Hopf theorem (see [2], [5]).

Theorem 1.3. *Let M be a compact immersed surface in the unit sphere S^3 with constant mean curvature H . If M is a topological sphere, then M is totally umbilical.*

2. NOTATIONS AND AUXILIARY RESULTS

Let M be a compact connected immersed surface in the unit sphere S^3 . Following the notations of [1] and [3],

Lemma 2.1. $\frac{1}{2}\Delta\Phi = \Phi(B - \Phi) + \sum \phi_{ijk}^2$ where ϕ_{ijk} denote the covariant derivative of ϕ_{ij} .

Lemma 2.2. $|\nabla\Phi|^2 = 2\Phi \sum \phi_{ijk}^2$.

Lemma 2.3. *If $\Phi \geq B$, then Φ is a constant function, $\Phi \equiv B$, and M is an $H(r)$ -torus.*

Proof. By Lemmas 2.1 and 2.2, we have $\frac{1}{2}\Delta \log \Phi = B - \Phi$ at the points where Φ is positive. It follows that $\Phi = B$ on M . $\square$

According to Lemma 2.1 and Lemma 2.3, we see that if Φ is a constant function, then either $\Phi = 0$ or $\Phi = B$.

In the minimal case, H. B. Lawson proved that the set of all zeros of Φ is either the whole space M or at most a finite set of points [7]. We need the following analogous result for the case that M is with constant mean curvature.

Lemma 2.4. *The set of all zeros of Φ is either the whole M or at most a finite set of points.*

Proof. The proof of the lemma is similar to that of Lawson. Let us sketch the proof for completeness. We use an isothermal coordinate (u, v) on a neighborhood D in M . Denote the position vector of this immersion by X and the unit normal of M in S^3 by N . Then the mean curvature H and the Gaussian curvature K are given

by $\frac{1}{F^2}(X_{uu} + X_{vv}) \cdot N$ and $\frac{1}{F^4}[(X_{uu} \cdot N)(X_{vv} \cdot N) - (X_{uv} \cdot N)^2]$ respectively, where $F^2 = X_u \cdot X_u = X_v \cdot X_v$.

Since M is of constant mean curvature, the Weingarten equations imply that $X_{uv} \cdot N$ is harmonic in the (u, v) coordinate, $X_{uu} \cdot N$ and $X_{vv} \cdot N$ differ by a constant if $X_{uv} \cdot N$ is constant on D , and the zero set of $X_{uv} \cdot N$ and that of $(X_{uu} - X_{vv}) \cdot N$ intersect transversely at the points where the gradient of $X_{uv} \cdot N$ does not vanish. Let G be the set of all points where the gradient of $X_{uv} \cdot N$ vanishes. Since $X_{uv} \cdot N$ is harmonic, G is either isolated or the whole D . For G being isolated, the set of all zeros of Φ is isolated. In the other case, the set of all zeros of Φ is either empty or the whole D . $\square$

Lemma 2.5. *If M is not totally umbilical, then*

$$\lim_{\epsilon \rightarrow 0} \sum_{i=1}^k \int_{\partial B_\epsilon(p_i)} \frac{\Phi_r}{\Phi} = 16\pi(g-1)$$

where $p_1, p_2, \dots, p_k$ constitute all the zeros of Φ and Φ_r denotes the derivative of Φ on $\partial B_\epsilon(p_i)$ in the radial direction from p_i . In particular, if Φ is positive on M , then M is a topological torus.

Proof. At the points where Φ is positive, by Lemma 2.2, we get

$$(2.1) \quad \Delta \log \Phi = B - \Phi.$$

Integrating (2.1) over $M_\epsilon = M \setminus \bigcup_{i=1}^k B_\epsilon(p_i)$, we get, from the Gauss equation

$$(2.2) \quad 2K = B - \Phi,$$

where K is the Gaussian curvature of M , the assertion by Stokes's theorem and the theorem of Gauss-Bonnet. $\square$

Lemma 2.6.

$$\int_M \left(\frac{\pi}{2} + \tan^{-1} \frac{H - \sqrt{2\Phi}}{2} - \tan^{-1} \frac{H + \sqrt{2\Phi}}{2} \right) (B - \Phi) + 2\sqrt{2\Phi} \geq 4\pi^2(1+g).$$

Proof. Regard M as an immersed surface of $\mathbb{R}^4$. Then the total absolute curvature of M in the sense of [4] is given by

$$\begin{aligned} T(M) &= \int_M \int_0^{2\pi} \left| \left(\sin \theta + \frac{H + \sqrt{2S - H^2}}{2} \cos \theta \right) \left(\sin \theta + \frac{H - \sqrt{2S - H^2}}{2} \cos \theta \right) \right| d\theta dV \\ &= \int_M \left(\frac{\pi}{2} + \tan^{-1} \frac{H - \sqrt{2\Phi}}{2} - \tan^{-1} \frac{H + \sqrt{2\Phi}}{2} \right) (B - \Phi) + 2\sqrt{2\Phi}. \end{aligned}$$

By the well-known inequality of Chern-Lashof [4], we have

$$T(M) \geq \frac{\pi^2}{2}(b_0 + b_1 + b_2),$$

where b_i is the i th Betti number relative to the real field, for $i = 0, 1, 2$. Since M is two-dimensional, $b_0 = 1$, $b_1 = 2g$ and $b_2 = 1$. $\square$

3. PROOF OF MAIN RESULTS

We are now in position to prove the main result of Theorem 1.1. We may assume that Φ is positive except possibly at a finite set of points (see Lemma 2.4). By Lemmas 2.2 and 2.6, we get

$$\begin{aligned}
& \int_M 2\sqrt{2B} + m(B) \sum \phi_{ijk}^2 - \left(\frac{\pi}{2} + \tan^{-1} \frac{H - \sqrt{2\Phi}}{2} - \tan^{-1} \frac{H + \sqrt{2\Phi}}{2}\right)(B - \Phi) - 2\sqrt{2\Phi} \\
&= \int_M m(B) \sum \phi_{ijk}^2 + \left[\frac{2\sqrt{2}}{\sqrt{B} + \sqrt{\Phi}} - \left(\frac{\pi}{2} + \tan^{-1} \frac{H - \sqrt{2\Phi}}{2} - \tan^{-1} \frac{H + \sqrt{2\Phi}}{2}\right)\right](B - \Phi) \\
&= \lim_{\epsilon \rightarrow 0} \int_{M_\epsilon} m(B) \frac{|\nabla \Phi|^2}{2\Phi} + \frac{1}{2} \left[\frac{2\sqrt{2}}{\sqrt{B} + \sqrt{\Phi}} - \left(\frac{\pi}{2} + \tan^{-1} \frac{H - \sqrt{2\Phi}}{2} - \tan^{-1} \frac{H + \sqrt{2\Phi}}{2}\right)\right] \Delta \log \Phi \\
&= \lim_{\epsilon \rightarrow 0} \int_{M_\epsilon} m(B) \frac{|\nabla \Phi|^2}{2\Phi} + \frac{1}{2} \nabla \left[\frac{2\sqrt{2}}{\sqrt{B} + \sqrt{\Phi}} - \left(\frac{\pi}{2} + \tan^{-1} \frac{H - \sqrt{2\Phi}}{2} - \tan^{-1} \frac{H + \sqrt{2\Phi}}{2}\right)\right] \nabla \log \Phi \\
&\quad - \lim_{\epsilon \rightarrow 0} \int_{\partial M_\epsilon} \left[\frac{2\sqrt{2}}{\sqrt{B} + \sqrt{\Phi}} - \left(\frac{\pi}{2} + \tan^{-1} \frac{H - \sqrt{2\Phi}}{2} - \tan^{-1} \frac{H + \sqrt{2\Phi}}{2}\right)\right] \frac{\Phi_r}{2\Phi} \\
&= 8\pi(1-g) \left(\frac{2\sqrt{2}}{\sqrt{B}} - \frac{\pi}{2}\right) + \int_M [m(B) - 2\sqrt{2} \frac{\sqrt{B}\Phi^2 + 2(B-2)\Phi + B\sqrt{B}}{(\sqrt{B} + \Phi)^2((\Phi^2 - B)^2 + 8\Phi^2)}] \sum \phi_{ijk}^2 \\
&\geq 8\pi(1-g) \left(\frac{2\sqrt{2}}{\sqrt{B}} - \frac{\pi}{2}\right),
\end{aligned}$$

where the equality holds if and only if Φ is constant. On the other hand, according to Lemma 2.6, we get

$$(3.1) \quad 2\sqrt{2B} \text{Area}(M) + m(B) \int_M \sum \phi_{ijk}^2 \geq 8\pi^2 g - 2\sqrt{\frac{2}{B}} 8\pi(g-1).$$

By combining (2.2) with the inequality (3.1), it follows from Lemma 2.1 that

$$m(B) \int_M \Phi^2 \geq 8\pi^2 g + (Bm(B) - 2\sqrt{\frac{2}{B}}) \int_M \Phi \geq 8\pi^2 g.$$

It remains to show that the second assertion holds. Suppose now that $\|\Phi\|_2 \leq 2\pi\sqrt{\frac{2}{m(B)}}$. The first assertion implies that $g = 0$ or 1 . If $g = 1$, then $\|\Phi\|_2 = 2\pi\sqrt{\frac{2}{m(B)}}$ and Φ is a constant function. If $g = 0$, then by Theorem 1.3, M is totally umbilical. This completes the proof of Theorem 1.1.

REFERENCES

- [1] H. Alencar and M. do Carmo, *Hypersurface with constant mean curvature in spheres*, Proc. Amer. Math. Soc. 120 (1994), 1223-1229. MR **94f**:53108
- [2] S. S. Chern, *On surface of constant mean curvature in a three-dimensional space of constant curvature*, Geometric Dynamics (Rio de Janeiro, 1981), Lecture Notes in Math., 1007, Springer-Verlag, Berlin, New York, 1983, pp. 104-108. MR **86b**:53058
- [3] S. S. Chern, M. do Carmo and S. Kobayashi, *Minimal submanifold of a sphere with second fundamental form of constant length*, Functional Analysis and Related Fields, Springer-Verlag, 1970, pp. 59-75. MR **42**:8424
- [4] S. S. Chern and R. K. Lashof, *On the total curvature of immersed manifolds. II*, Michigan Math. J. 5(1985), 5-12. MR **20**:4301
- [5] H. Hopf, *Differential geometry in the large*, Lecture Notes in Math., 1000, Springer-Verlag, 1983, pp. 136-146. MR **85b**:53001
- [6] Y. J. Hsu, *A global pinching theorem for compact minimal surfaces in S^3* , Proc. Amer. Math. Soc. 113(1991), 1041-1044. MR **92c**:53040

- [7] H. B. Lawson, *Complete minimal surfaces in S^3* , Ann. of Math. 92(1970), 335-374. MR **42**:5170
- [8] C. L. Shen, *A global pinching theorem of minimal hypersurfaces in the sphere*, Proc. Amer. Math. Soc. 105(1989), 192-198. MR **90c**:53162

DEPARTMENT OF APPLIED MATHEMATICS, NATIONAL CHIAO TUNG UNIVERSITY, HSINCHU, TAI-
WAN

E-mail address: `yjhsu@math.nctu.edu.tw`

DEPARTMENT OF APPLIED MATHEMATICS, NATIONAL CHIAO TUNG UNIVERSITY, HSINCHU, TAI-
WAN

E-mail address: `teich@math.sinica.edu.tw`