a -WEYL'S THEOREM FOR OPERATOR MATRICES

YOUNG MIN HAN AND SLAVIŠA V. DJORDJEVIĆ

(Communicated by Joseph A. Ball)

ABSTRACT. If $M_C = \begin{pmatrix} A & C \\ 0 & B \end{pmatrix}$ is a 2×2 upper triangular matrix on the Hilbert space $H \oplus K$, then a -Weyl's theorem for A and B need not imply a -Weyl's theorem for M_C , even when $C = 0$. In this note we explore how a -Weyl's theorem and a -Browder's theorem survive for 2×2 operator matrices on the Hilbert space.

1. INTRODUCTION

Throughout this note let H and K be Hilbert spaces, let $B(H, K)$ denote the set of bounded linear operators from H to K , and abbreviate $B(H, H)$ to $B(H)$ and let $K(H)$ denote the ideal of compact operators acting on H . If $T \in B(H)$ write $N(T)$ and $R(T)$ for the null space and range of T ; $\alpha(T) = \dim N(T)$. An operator $T \in B(H)$ is called upper semi-Fredholm if $R(T)$ is closed with finite dimensional null space and lower semi-Fredholm if $R(T)$ is closed with its range of finite co-dimension. If T is both upper semi- and lower semi-Fredholm, we call it *Fredholm*. The *index* of a Fredholm operator $T \in B(H)$ is the integer $i(T) = \alpha(T) - \alpha(T^*)$. An operator $T \in B(H)$ is called *Weyl* if it is Fredholm of index zero and is called *Browder* if it is Fredholm of “finite ascent and descent”. If $T \in B(H)$ write $\sigma(T)$ for the spectrum of T ; $\sigma_a(T)$ for the approximate point spectrum of T ; $\pi_0(T)$ for the set of eigenvalues of T ; $\pi_{00}(T)$ for the isolated points of $\sigma(T)$ which are eigenvalues of finite multiplicity; $\pi_{00}^a(T)$ for the isolated points of $\sigma_a(T)$ which are eigenvalues of finite multiplicity. The essential spectrum $\sigma_e(T)$, the Weyl spectrum $\omega(T)$ and the Browder spectrum $\sigma_b(T)$ of $T \in B(H)$ are defined by ([10], [11])

$$\begin{aligned}\sigma_e(T) &= \{\lambda \in \mathbb{C} : T - \lambda I \text{ is not Fredholm}\}, \\ \omega(T) &= \{\lambda \in \mathbb{C} : T - \lambda I \text{ is not Weyl}\}, \\ \sigma_b(T) &= \{\lambda \in \mathbb{C} : T - \lambda I \text{ is not Browder}\};\end{aligned}$$

evidently

$$\sigma_e(T) \subseteq \omega(T) \subseteq \sigma_b(T) = \sigma_e(T) \cup \text{acc } \sigma(T),$$

Received by the editors February 29, 2000 and, in revised form, August 25, 2000.

2000 *Mathematics Subject Classification*. Primary 47A50, 47A53.

Key words and phrases. Weyl spectrum, essential approximate point spectrum, Browder essential approximate point spectrum, a -Weyl's theorem, Weyl's theorem, a -Browder's theorem, Browder's theorem.

This work was supported by the Brain Korea 21 Project (through Seoul National University).

where we write $\text{acc } X$ for the accumulation points of $X \subseteq \mathbb{C}$. We say that *Weyl's theorem holds for $T \in B(H)$* if there is equality

$$(1.1) \quad \sigma(T) \setminus \omega(T) = \pi_{00}(T).$$

This obviously implies ([12]) that *Browder's theorem holds for T* :

$$(1.2) \quad \sigma(T) = \omega(T) \cup \pi_{00}(T).$$

By definition, $\sigma_{ea}(T) = \bigcap \{\sigma_a(T + K) : K \in K(H)\}$ is the essential approximate point spectrum and $\sigma_{ab}(T) = \bigcap \{\sigma_a(T + K) : TK = KT \text{ and } K \in K(H)\}$ is the Browder essential approximate point spectrum. We say that *a -Weyl's theorem holds for $T \in B(H)$* if there is equality

$$(1.3) \quad \sigma_a(T) \setminus \sigma_{ea}(T) = \pi_{00}^a(T),$$

and that *a -Browder's theorem holds for $T \in B(H)$* if there is equality

$$(1.4) \quad \sigma_{ea}(T) = \sigma_{ab}(T).$$

We notice that a -Weyl's theorem and a -Browder's theorem are kinds of "approximate point spectrum" versions of Weyl's theorem and Browder's theorem, respectively. It is known ([5], [7], [18]) that if $T \in B(H)$, then we have

$$(1.5) \quad a\text{-Weyl's theorem} \implies a\text{-Browder's theorem} \implies \text{Browder's theorem}.$$

V. Rakočević ([18]) has shown that the equality (1.3) holds for cohyponormal operators. Recently S.V. Djordjević and D.S. Djordjević ([5]) showed that if T^* is quasihyponormal, then T obeys a -Weyl's theorem.

2. a -WEYL'S THEOREM FOR 2×2 DIAGONAL, SKEW-DIAGONAL OPERATOR MATRICES

a -Weyl's theorem may or may not hold for a direct sum of operators for which a -Weyl's theorem holds. For example, if $U \in B(l_2)$ is the unilateral shift, then a -Weyl's theorem holds for both U and U^* , while it does not hold for $U \oplus U^*$. In fact, $\sigma_a(U \oplus U^*) = \{z \in \mathbb{C} : |z| \leq 1\}$ and $\sigma_{ea}(U \oplus U^*) = \{z \in \mathbb{C} : |z| = 1\}$. Therefore a -Weyl's theorem does not hold for $U \oplus U^*$. An operator $T \in B(H)$ is called *approximate-isoloid* (abbrev. a -isoloid) if every isolated point of $\sigma_a(T)$ is an eigenvalue of T and an operator $T \in B(H)$ is called *isoloid* if every isolated point of $\sigma(T)$ is an eigenvalue of T . Clearly, if T is a -isoloid, then it is isoloid. However the converse is not true. Consider the following example: let $T = T_1 \oplus T_2$, where T_1 is the unilateral shift on l_2 and T_2 is an injective quasinilpotent on l_2 . Then $\sigma(T) = \{z \in \mathbb{C} : |z| \leq 1\}$ and $\sigma_a(T) = \{z \in \mathbb{C} : |z| = 1\} \cup \{0\}$. Therefore T is isoloid but is not a -isoloid.

We consider the sets

$$\begin{aligned} \Phi_+(H) &= \{T \in B(H) : R(T) \text{ is closed and } \alpha(T) < \infty\}, \\ \Phi_-(H) &= \{T \in B(H) : R(T) \text{ is closed and } \alpha(T^*) < \infty\}, \\ \Phi_+^-(H) &= \{T \in B(H) : T \in \Phi_+(H) \text{ and } i(T) \leq 0\}. \end{aligned}$$

It is known ([18]) that if $T \in B(H)$, then $\sigma_{ea}(T) = \{\lambda \in \mathbb{C} : T - \lambda \notin \Phi_+^-(H)\}$.

When $A \in B(H)$ and $B \in B(K)$ are given we denote by M_C an operator acting on $H \oplus K$ of the form

$$M_C = \begin{pmatrix} A & C \\ 0 & B \end{pmatrix},$$

where $C \in B(K, H)$.

We begin with:

Lemma 2.1. *For a given pair (A, B) of operators, if $\begin{pmatrix} A & 0 \\ 0 & B \end{pmatrix} \in \Phi_+^-(H \oplus K)$, then $M_C \in \Phi_+^-(H \oplus K)$ for every $C \in B(K, H)$. Hence, in particular, we have*

$$(2.1.1) \quad \sigma_{ea}(M_C) \subset \sigma_{ea} \begin{pmatrix} A & 0 \\ 0 & B \end{pmatrix} \subset \sigma_{ea}(A) \cup \sigma_{ea}(B).$$

Proof. Suppose $\begin{pmatrix} A & 0 \\ 0 & B \end{pmatrix} \in \Phi_+^-(H \oplus K)$. Then $\begin{pmatrix} A & 0 \\ 0 & B \end{pmatrix} \in \Phi_+(H \oplus K)$ and $i \begin{pmatrix} A & 0 \\ 0 & B \end{pmatrix} \leq 0$. So A and B are upper semi-Fredholm and $i(A) + i(B) \leq 0$.

Observe that

$$(2.1.2) \quad M_C = \begin{pmatrix} I & 0 \\ 0 & B \end{pmatrix} \begin{pmatrix} I & C \\ 0 & I \end{pmatrix} \begin{pmatrix} A & 0 \\ 0 & I \end{pmatrix}.$$

Since $\begin{pmatrix} I & C \\ 0 & I \end{pmatrix}$ is invertible for every $C \in B(K, H)$, and since $\begin{pmatrix} I & 0 \\ 0 & B \end{pmatrix}$ and $\begin{pmatrix} A & 0 \\ 0 & I \end{pmatrix}$ are both upper semi-Fredholm, it follows that M_C is upper semi-Fredholm. But since $i(M_C) = i(A) + i(B) \leq 0$, hence $M_C \in \Phi_+^-(H \oplus K)$. The inclusions in (2.1.1) are evident from the first assertion. $\square$

Theorem 2.2. *Suppose a -Weyl's theorem holds for $A \in B(H)$ and $B \in B(K)$.*

(1) *If a -Weyl's theorem holds for $\begin{pmatrix} A & 0 \\ 0 & B \end{pmatrix}$, then*

$$(2.2.1) \quad \sigma_{ea} \begin{pmatrix} A & 0 \\ 0 & B \end{pmatrix} = \sigma_{ea}(A) \cup \sigma_{ea}(B).$$

(2) *If A and B are a -isoloid, then the converse of (1) is true.*

Proof. (1) By Lemma 2.1, $\sigma_{ea} \begin{pmatrix} A & 0 \\ 0 & B \end{pmatrix} \subset \sigma_{ea}(A) \cup \sigma_{ea}(B)$. Since a -Weyl's theorem holds for $\begin{pmatrix} A & 0 \\ 0 & B \end{pmatrix}$, a -Browder's theorem holds for $\begin{pmatrix} A & 0 \\ 0 & B \end{pmatrix}$. Therefore $\sigma_{ea}(A) \cup \sigma_{ea}(B) \subset \sigma_{ab}(A) \cup \sigma_{ab}(B) = \sigma_{ab} \begin{pmatrix} A & 0 \\ 0 & B \end{pmatrix} = \sigma_{ea} \begin{pmatrix} A & 0 \\ 0 & B \end{pmatrix}$, that is, $\sigma_{ea}(A) \cup \sigma_{ea}(B) \subset \sigma_{ea} \begin{pmatrix} A & 0 \\ 0 & B \end{pmatrix}$.

(2) For the statement (2) observe that if A and B are a -isoloid, then

$$(2.2.2) \quad \pi_{00}^a \begin{pmatrix} A & 0 \\ 0 & B \end{pmatrix} = (\pi_{00}^a(A) \setminus \sigma_a(B)) \cup (\pi_{00}^a(B) \setminus \sigma_a(A)) \cup (\pi_{00}^a(A) \cap \pi_{00}^a(B)).$$

If a -Weyl's theorem holds for A and B , then the right-hand side of (2.2.2) is the set $(\sigma_a(A) \cup \sigma_a(B)) \setminus (\sigma_{ea}(A) \cup \sigma_{ea}(B))$. Thus if (2.2.1) holds, then $\pi_{00}^a \begin{pmatrix} A & 0 \\ 0 & B \end{pmatrix} = (\sigma_a(A) \cup \sigma_a(B)) \setminus (\sigma_{ea}(A) \cup \sigma_{ea}(B)) = \sigma_a \begin{pmatrix} A & 0 \\ 0 & B \end{pmatrix} \setminus \sigma_{ea} \begin{pmatrix} A & 0 \\ 0 & B \end{pmatrix}$. Thus a -Weyl's holds for $\begin{pmatrix} A & 0 \\ 0 & B \end{pmatrix}$. $\square$

The assumption “ A and B are a -isoloid” is essential in the statement (2) of Theorem 2.2. For example if $A \in B(l_2)$ and $B \in B(l_2)$ are given by

$$A(x_1, x_2, x_3, x_4, \dots) = (x_1, 0, x_3, x_4, \dots)$$

and

$$B(x_1, x_2, x_3, \dots) = (0, x_1, \frac{1}{2}x_2, \frac{1}{3}x_3, \dots),$$

then we have that (a) a -Weyl's theorem holds for A and B ; (b) $\sigma_{ea}(A) = \{1\}$ and $\sigma_{ea}(B) = \{0\}$; (c) $\sigma_a\left(\begin{smallmatrix} A & 0 \\ 0 & B \end{smallmatrix}\right) = \sigma_{ea}\left(\begin{smallmatrix} A & 0 \\ 0 & B \end{smallmatrix}\right) = \{0, 1\}$; (d) $\pi_{00}^a\left(\begin{smallmatrix} A & 0 \\ 0 & B \end{smallmatrix}\right) = \{0\}$; (e) B is not a -isoloid.

Next we consider a -Weyl's theorem for 2×2 skew-diagonal operator matrix of the form $\begin{pmatrix} 0 & A \\ B & 0 \end{pmatrix}$. a -Weyl's theorem for the skew-diagonal matrices is more intricate in comparison with the diagonal matrices. We begin with:

Lemma 2.3. *If $A \in B(H, K)$ and $B \in B(K, H)$, then*

- (1) $\sigma_a(AB) \setminus \{0\} = \sigma_a(BA) \setminus \{0\}$.
- (2) $\sigma_{ea}(AB) \setminus \{0\} = \sigma_{ea}(BA) \setminus \{0\}$.

Proof. (1) Remember ([8]) that if $\lambda \neq 0$, then

$$(2.3.1) \quad \begin{pmatrix} AB - \lambda & 0 \\ 0 & I \end{pmatrix} = F(\lambda) \begin{pmatrix} BA - \lambda & 0 \\ 0 & I \end{pmatrix} E(\lambda),$$

where $E(\lambda)$ and $F(\lambda)$ are both invertible for each $\lambda \neq 0$. But if ST is bounded below and T is invertible, then S is bounded below, and so $\sigma_a(AB) \setminus \{0\} \subset \sigma_a(BA) \setminus \{0\}$. Similarly, $\sigma_a(BA) \setminus \{0\} \subset \sigma_a(AB) \setminus \{0\}$.

(2) It follows from (2.3.1) that if $\lambda \neq 0$,

$$N(AB - \lambda) \cong N(BA - \lambda) \text{ and } (R(AB - \lambda))^\perp \cong (R(BA - \lambda))^\perp,$$

which gives the result, where $\cong$ means that there exists an invertible operator between spaces. $\square$

Although $\sigma_a(AB) = \sigma_a(BA)$, we need not expect that $\sigma_{ea}(AB) = \sigma_{ea}(BA)$. Consider the following example: let $\dim \mathcal{H} < \infty$ and let $S, T \in B(l_2)$ be given by

$S(x_1, x_2, x_3, \dots) = (0, x_1, 0, x_2, 0, x_3, \dots)$ and $T(x_1, x_2, x_3, \dots) = (x_2, x_4, x_6, \dots)$, and let $A = \begin{pmatrix} 0_{\mathcal{H}} & 0 \\ 0 & S \end{pmatrix}$ and $B = \begin{pmatrix} 0_{\mathcal{H}} & 0 \\ 0 & T \end{pmatrix}$. Then $\sigma_a(AB) = \sigma_a(BA) = \{0, 1\}$, while $\sigma_{ea}(AB) = \{0, 1\} \neq \{1\} = \sigma_{ea}(BA)$.

Theorem 2.4. *If $A \in B(H, K)$ and $B \in B(K, H)$, then*

$$(2.4.1) \quad \sigma_{ea} \begin{pmatrix} AB & 0 \\ 0 & BA \end{pmatrix} = \sigma_{ea}(AB) \cup \sigma_{ea}(BA).$$

Hence, in particular, if AB and BA are a -isoloid and if a -Weyl's threorem holds for AB and BA , then a -Weyl's theorem holds for $\begin{pmatrix} AB & 0 \\ 0 & BA \end{pmatrix}$.

Proof. By Lemma 2.1, $\sigma_{ea}\left(\begin{smallmatrix} AB & 0 \\ 0 & BA \end{smallmatrix}\right) \subset \sigma_{ea}(AB) \cup \sigma_{ea}(BA)$. Conversely, suppose $\lambda \notin \sigma_{ea}\left(\begin{smallmatrix} AB & 0 \\ 0 & BA \end{smallmatrix}\right)$. Then $AB - \lambda$ and $BA - \lambda$ are upper semi-Fredholm and $i\left(\begin{smallmatrix} AB - \lambda & 0 \\ 0 & BA - \lambda \end{smallmatrix}\right) \leq 0$. Now we show that $i(AB - \lambda) = i(BA - \lambda)$. If $\lambda \neq 0$, then by Lemma 2.3 $i(AB - \lambda) = i(BA - \lambda)$. If $\lambda = 0$, then since AB and BA are upper semi-Fredholm, it follows from the continuity of the index that for sufficiently small $|\mu|$ with $\mu \neq 0$, $i(AB) = i(AB - \mu) = i(BA - \mu) = i(BA)$. This proves (2.4.1). The second assertion follows from Theorem 2.2. $\square$

Corollary 2.5. *Let $A \in B(H, K)$ and $B \in B(K, H)$. If AB and BA obeys a -Browder's theorem, then a -Browder's theorem holds for $\begin{pmatrix} AB & 0 \\ 0 & BA \end{pmatrix}$.*

Proof. Since $\sigma_{ea}\left(\begin{smallmatrix} AB & 0 \\ 0 & BA \end{smallmatrix}\right) = \sigma_{ea}(AB) \cup \sigma_{ea}(BA)$, the result follows from [7, Theorem 3.11]. $\square$

Example 2.6. (1) If $H = K$ in Theorem 2.4, one might expect to replace the condition “ a -Weyl’s theorem holds for AB and BA ” by the condition “ a -Weyl’s theorem holds for A and B ”. However, this is not true. Consider the following example as follows: let T_1 and T_2 be defined on l_2 by

$$\begin{aligned} T_1(x_1, x_2, x_3, \dots) &= (x_1, 0, \frac{1}{2}x_2, \frac{1}{2}x_3, \dots), \\ T_2(x_1, x_2, x_3, \dots) &= (0, \frac{1}{2}x_1, \frac{1}{3}x_2, \dots). \end{aligned}$$

Let $T = \begin{pmatrix} T_1 & 0 \\ 0 & T_2 - I \end{pmatrix}$ on $l_2 \oplus l_2$. Then

$$\begin{aligned} \sigma_a(T) &= \{-1\} \cup \{z \in \mathbb{C} : |z| = \frac{1}{2}\} \cup \{1\}, \\ \sigma_{ea}(T) &= \{-1\} \cup \{z \in \mathbb{C} : |z| = \frac{1}{2}\}, \\ \pi_{00}^a(T) &= \{1\}. \end{aligned}$$

Therefore a -Weyl’s theorem holds for T . However,

$$\begin{aligned} \sigma_a(T^2) &= \{z \in \mathbb{C} : |z| = \frac{1}{4}\} \cup \{1\}, \\ \sigma_{ea}(T^2) &= \{z \in \mathbb{C} : |z| = \frac{1}{4}\} \cup \{1\}, \\ \pi_{00}^a(T^2) &= \{1\}, \end{aligned}$$

and hence a -Weyl’s theorem doesn’t hold for T^2 .

(2) Since

$$\begin{pmatrix} 0 & A \\ B & 0 \end{pmatrix}^2 = \begin{pmatrix} AB & 0 \\ 0 & BA \end{pmatrix},$$

we might expect that “ a -Weyl’s theorem for $\begin{pmatrix} 0 & A \\ B & 0 \end{pmatrix}$ ” is inherited from “ a -Weyl’s theorem for $\begin{pmatrix} AB & 0 \\ 0 & BA \end{pmatrix}$ ”. However, in general, a -Weyl’s theorem need not be transmitted from T^2 to T . Consider the following example as follows: let T_1 and T_2 be defined on l_2 by

$$\begin{aligned} T_1(x_1, x_2, x_3, \dots) &= (0, x_1, x_2, x_3, x_4, \dots), \\ T_2(x_1, x_2, x_3, \dots) &= (\frac{1}{2}x_2, \frac{1}{3}x_3, \frac{1}{4}x_4, \dots). \end{aligned}$$

Let $T = \begin{pmatrix} T_1 + I & 0 \\ 0 & T_2 - I \end{pmatrix}$ on $l_2 \oplus l_2$. Then

$$\begin{aligned} \sigma_a(T) &= \{-1\} \cup \{z \in \mathbb{C} : |z - 1| = 1\}, \\ \sigma_{ea}(T) &= \{-1\} \cup \{z \in \mathbb{C} : |z| = 1\}, \\ \pi_{00}^a(T) &= \{-1\}. \end{aligned}$$

Therefore a -Weyl’s theorem does not hold for T . However,

$$\begin{aligned} \sigma_a(T^2) &= \{re^{i\theta} : r = 2(1 + \cos \theta)\}, \\ \sigma_{ea}(T^2) &= \{re^{i\theta} : r = 2(1 + \cos \theta)\}, \\ \pi_{00}^a(T^2) &= \phi, \end{aligned}$$

and hence a -Weyl’s theorem holds for T^2 .

(3) Generally, “ a -Weyl’s theorem holds for AB ” does not imply “ a -Weyl’s theorem holds for BA ”. For example suppose the operators $T_1, T_2, T_3 \in B(l_2)$ are given by

$$\begin{aligned} T_1(x_1, x_2, x_3, \dots) &= \left(\frac{1}{2}x_2, \frac{1}{3}x_3, \frac{1}{4}x_4, \dots\right), \\ T_2(x_1, x_2, x_3, \dots) &= (0, x_1, 0, x_2, 0, x_3, 0, \dots), \\ T_3(x_1, x_2, x_3, \dots) &= (x_2, x_4, x_6, x_8, \dots). \end{aligned}$$

Let $A = \begin{pmatrix} T_1 & 0 \\ 0 & T_2 \end{pmatrix}$ and $B = \begin{pmatrix} I & 0 \\ 0 & T_3 \end{pmatrix}$. Then

$$\begin{aligned} \sigma_a(AB) &= \sigma_{ea}(AB) = \sigma_a(BA) = \sigma_{ea}(BA) = \{0, 1\}, \\ \pi_{00}^a(AB) &= \phi, \\ \pi_{00}^a(BA) &= \{0\}. \end{aligned}$$

Therefore a -Weyl’s theorem holds for AB but fails for BA .

Theorem 2.7. *Let $A \in B(K, H)$ and $B \in B(H, K)$ be such that AB and BA are a -isoloid. If a -Weyl’s theorem holds for AB and BA , then the following statements are equivalent:*

- (1) a -Weyl’s theorem holds for $\begin{pmatrix} 0 & A \\ B & 0 \end{pmatrix}$;
- (2) $\sigma_{ea}\left(\begin{pmatrix} 0 & A \\ B & 0 \end{pmatrix}\right) = \sqrt{\sigma_{ea}(AB) \cup \sigma_{ea}(BA)}$;
- (3) a -Browder’s theorem holds for $\begin{pmatrix} 0 & A \\ B & 0 \end{pmatrix}$,

where $\sqrt{X}$ denotes the set of square roots of complex numbers in $X \subset \mathbb{C}$.

Proof. (1) $\Leftrightarrow$ (2) Suppose $\sigma_{ea}\left(\begin{pmatrix} 0 & A \\ B & 0 \end{pmatrix}\right) = \sqrt{\sigma_{ea}(AB) \cup \sigma_{ea}(BA)}$. Then it follows from a similar argument of ([13], (3.10); (4.3)) that

$$\begin{aligned} \sigma_a\left(\begin{pmatrix} 0 & A \\ B & 0 \end{pmatrix}\right) \setminus \sigma_{ea}\left(\begin{pmatrix} 0 & A \\ B & 0 \end{pmatrix}\right) &= \sqrt{\sigma_a(AB) \cup \sigma_a(BA)} \setminus \sqrt{\sigma_{ea}(AB) \cup \sigma_{ea}(BA)} \\ &= \sqrt{(\sigma_a(AB) \cup \sigma_a(BA)) \setminus (\sigma_{ea}(AB) \cup \sigma_{ea}(BA))} \\ &= \sqrt{\sigma_a\left(\begin{pmatrix} AB & 0 \\ 0 & BA \end{pmatrix}\right) \setminus \sigma_{ea}\left(\begin{pmatrix} AB & 0 \\ 0 & BA \end{pmatrix}\right)} \\ &= \sqrt{\pi_{00}^a\left(\begin{pmatrix} AB & 0 \\ 0 & BA \end{pmatrix}\right)}. \end{aligned}$$

Now we show that

$$(2.7.1) \quad \sqrt{\pi_{00}^a\left(\begin{pmatrix} AB & 0 \\ 0 & BA \end{pmatrix}\right)} = \pi_{00}^a\left(\begin{pmatrix} 0 & A \\ B & 0 \end{pmatrix}\right).$$

But since $\sigma_a\left(\begin{pmatrix} 0 & A \\ B & 0 \end{pmatrix}\right)$ is symmetric with respect to the origin, it follows from the approximate point spectral mapping theorem that

$$\sqrt{\text{iso } \sigma_a\left(\begin{pmatrix} AB & 0 \\ 0 & BA \end{pmatrix}\right)} = \sqrt{\text{iso } \left(\sigma_a\left(\begin{pmatrix} 0 & A \\ B & 0 \end{pmatrix}\right)\right)^2} = \text{iso } \sigma_a\left(\begin{pmatrix} 0 & A \\ B & 0 \end{pmatrix}\right),$$

where we write $\text{iso } X$ for the isolated points of $X \subset \mathbb{C}$. Thus for (2.7.1) it suffices to show that for any $\lambda \in \mathbb{C}$,

$$(2.7.2) \quad 0 < \alpha\left(\begin{pmatrix} AB - \lambda & 0 \\ 0 & BA - \lambda \end{pmatrix}\right) < \infty \iff 0 < \alpha\left(\begin{pmatrix} -\sqrt{\lambda} & A \\ B & -\sqrt{\lambda} \end{pmatrix}\right) < \infty.$$

If $\lambda = 0$, then (2.7.2) follows from the observation

$$(2.7.3) \quad 0 < \alpha(N(A) \oplus N(B)) < \infty \iff 0 < \alpha(N(AB) \oplus N(BA)) < \infty.$$

If $\lambda \neq 0$, then (2.7.2) follows from the observation

$$\begin{aligned} & \bigvee \left\{ \left(x, \frac{1}{\sqrt{\lambda}} Bx \right) : x \in N(AB - \lambda) \right\} \cup \bigvee \left\{ \left(\frac{1}{\sqrt{\lambda}} Ay, y \right) : y \in N(BA - \lambda) \right\} \\ & \subset N \begin{pmatrix} -\sqrt{\lambda} & A \\ B & -\sqrt{\lambda} \end{pmatrix} \subset N(AB - \lambda) \oplus N(BA - \lambda), \end{aligned}$$

where $\bigvee F$ denotes the closed linear span of F . This proves (2.7.1).

Suppose a -Weyl's theorem holds for $\begin{pmatrix} 0 & A \\ B & 0 \end{pmatrix}$.

Then $\sigma_{ea} \begin{pmatrix} 0 & A \\ B & 0 \end{pmatrix} = \sigma_a \begin{pmatrix} 0 & A \\ B & 0 \end{pmatrix} \setminus \pi_{00}^a \begin{pmatrix} 0 & A \\ B & 0 \end{pmatrix}$. It will be shown that

$$\sigma_a \begin{pmatrix} 0 & A \\ B & 0 \end{pmatrix} \setminus \pi_{00}^a \begin{pmatrix} 0 & A \\ B & 0 \end{pmatrix} = \sqrt{\sigma_{ea}(AB) \cup \sigma_{ea}(BA)}.$$

Indeed,

$$\begin{aligned} \sigma_a \begin{pmatrix} 0 & A \\ B & 0 \end{pmatrix} \setminus \pi_{00}^a \begin{pmatrix} 0 & A \\ B & 0 \end{pmatrix} &= \sqrt{\sigma_a(AB) \cup \sigma_a(BA)} \setminus \sqrt{\pi_{00}^a \begin{pmatrix} AB & 0 \\ 0 & BA \end{pmatrix}} \\ &= \sqrt{\sigma_a \begin{pmatrix} AB & 0 \\ 0 & BA \end{pmatrix} \setminus \pi_{00}^a \begin{pmatrix} AB & 0 \\ 0 & BA \end{pmatrix}} \\ &= \sqrt{\sigma_{ea} \begin{pmatrix} AB & 0 \\ 0 & BA \end{pmatrix}} \\ &= \sqrt{\sigma_{ea}(AB) \cup \sigma_{ea}(BA)}. \end{aligned}$$

Therefore $\sigma_{ea} \begin{pmatrix} 0 & A \\ B & 0 \end{pmatrix} = \sqrt{\sigma_{ea}(AB) \cup \sigma_{ea}(BA)}$.

(2) $\Leftrightarrow$ (3) Suppose $\sigma_{ea} \begin{pmatrix} 0 & A \\ B & 0 \end{pmatrix} = \sqrt{\sigma_{ea}(AB) \cup \sigma_{ea}(BA)}$. Then a -Weyl's theorem holds for $\begin{pmatrix} 0 & A \\ B & 0 \end{pmatrix}$, and so a -Browder's theorem holds for $\begin{pmatrix} 0 & A \\ B & 0 \end{pmatrix}$. Conversely, suppose a -Browder's theorem holds for $\begin{pmatrix} 0 & A \\ B & 0 \end{pmatrix}$. Since AB and BA obeys a -Weyl's theorem, $\sigma_{ea}(AB) = \sigma_{ab}(AB)$ and $\sigma_{ea}(BA) = \sigma_{ab}(BA)$, respectively.

But since $\sigma_{ab} \begin{pmatrix} 0 & A \\ B & 0 \end{pmatrix} = \sqrt{\sigma_{ab}(AB) \cup \sigma_{ab}(BA)}$, $\sigma_{ea} \begin{pmatrix} 0 & A \\ B & 0 \end{pmatrix} = \sigma_{ab} \begin{pmatrix} 0 & A \\ B & 0 \end{pmatrix} = \sqrt{\sigma_{ab}(AB) \cup \sigma_{ab}(BA)} = \sqrt{\sigma_{ea}(AB) \cup \sigma_{ea}(BA)}$ and completes the proof. $\square$

ACKNOWLEDGEMENT

The authors are grateful to the referee for helpful comments concerning this paper.

REFERENCES

1. S.K. Berberian, *An extension of Weyl's theorem to a class of not necessarily normal operators*, Michigan Math. J. **16** (1969), 273–279. MR **40**:3335
2. S.K. Berberian, *The Weyl spectrum of an operator*, Indiana Univ. Math. J. **20** (1970), 529–544. MR **43**:5344
3. B. Chevreau, *On the spectral picture of an operator*, J. Operator Theory **4** (1980), 119–132. MR **81k**:47002
4. L.A. Coburn, *Weyl's theorem for nonnormal operators*, Michigan Math. J. **13** (1966), 285–288. MR **34**:1846
5. S.V. Djordjević and D.S. Djordjević, *Weyl's theorems: continuity of the spectrum and quasi-hyponormal operators*, Acta Sci. Math. (Szeged) **64** (1998), 259–269. MR **2000c**:47009

6. S.V. Djordjević and B.P. Duggal, *Weyl's theorems and continuity of spectra in the class of p -hyponormal operators*, Studia Math **143** (2000), 23–32. CMP 2001:08
7. S.V. Djordjević and Y.M. Han, *Browder's theorems and spectral continuity*, Glasgow Math. J. **42** (2000), 479–486. CMP 2001:04
8. I. Gohberg, S. Goldberg and M. A. Kaashoek, *Classes of Linear Operators (vol I)*, Birkhäuser, Basel, 1990. MR **93d**:47002
9. J.K. Han, H.Y. Lee and W.Y. Lee, *Invertible Completions of 2×2 Upper Triangular Operator Matrices*, Proc. Amer. Math. Soc. **128** (2000), 119–123. MR **2000c**:47003
10. R.E. Harte, *Fredholm, Weyl and Browder theory*, Proc. Royal Irish Acad. **85A** (2) (1985), 151–176. MR **87h**:47029
11. R.E. Harte, *Invertibility and Singularity for Bounded Linear Operators*, Dekker, New York, 1988. MR **89d**:47001
12. R.E. Harte and W.Y. Lee, *Another note on Weyl's theorem*, Trans. Amer. Math. Soc. **349** (1997), 2115–2124. MR **98j**:47024
13. R.E. Harte, W.Y. Lee and L.L. Littlejohn, *On generalized Riesz points*, J. Operator Theory (to appear).
14. W.Y. Lee and S.H. Lee, *A spectral mapping theorem for the Weyl spectrum*, Glasgow Math. J. **38**(1) (1996), 61–64. MR **97c**:47023
15. W.Y. Lee, *Weyl's Theorem For Operator Matrices*, Integral Equations and Operator Theory **32** (1998), 319–331. MR **99g**:47023
16. W.Y. Lee, *Weyl Spectra of Operator Matrices*, Proc. Amer. Math. Soc. **129** (2001), 131–138. CMP 2001:01
17. K.K. Oberai, *On the Weyl spectrum (II)*, Illinois J. Math. **21** (1977), 84–90. MR **55**:1102
18. V. Rakočević, *On the essential approximate point spectrum II*, Mat. Vesnik **36** (1984), 89–97. MR **88h**:47019
18. V. Rakočević, *Approximate point spectrum and commuting compact perturbations*, Glasgow Math. J. **28** (1986), 193–198. MR **87k**:47006
19. H. Weyl, *Über beschränkte quadratische Formen, deren Differenz vollsteig ist*, Rend. Circ. Mat. Palermo **27** (1909), 373–392.

DEPARTMENT OF MATHEMATICS, SUNGKYUNKWAN UNIVERSITY, SUWON 440-746, KOREA

E-mail address: ymhan@math.skku.ac.kr

Current address: Department of Mathematics, 14 MacLean Hall, University of Iowa, Iowa City, Iowa 52242-1419

E-mail address: yhan@math.uiowa.edu

DEPARTMENT OF MATHEMATICS, FACULTY OF PHILOSOPHY, UNIVERSITY OF NIŠ, ĆIRILA AND METODIJA 2, 18000 NIŠ, YUGOSLAVIA

E-mail address: slavdj@archimed.filfak.ni.ac.yu