

DIRECT SUMS OF LOCAL TORSION-FREE ABELIAN GROUPS

DAVID M. ARNOLD

(Communicated by Stephen D. Smith)

ABSTRACT. The category of local torsion-free abelian groups of finite rank is known to have the cancellation and n -th root properties but not the Krull-Schmidt property. It is shown that 10 is the least rank of a local torsion-free abelian group with two non-equivalent direct sum decompositions into indecomposable summands. This answers a question posed by M.C.R. Butler in the 1960's.

1. INTRODUCTION

Let TF denote the category of local torsion-free abelian groups of finite rank, where an abelian group G is *local* if there is a fixed prime p with $qG = G$ for each prime $q \neq p$. Each M in TF has the *cancellation property* (if $M \oplus N$ is isomorphic to $M \oplus K$ in TF , then N is isomorphic to K), and the *n -th root property* (if the direct sum M^n of n copies of M is isomorphic to N^n for some N in TF , then M is isomorphic to N) [Lady 75]. An M in TF is a *Krull-Schmidt group* if any two direct sum decompositions of M into indecomposable summands are *equivalent*, i.e. unique up to isomorphism and order of summands.

M.C.R. Butler, in an unpublished note dating from the 1960's, constructed an example of a local torsion-free abelian group of rank 16 that is not a Krull-Schmidt group (see [Arnold 82]) and asked for the smallest such rank. An example of a rank-10 local torsion-free abelian group that is not a Krull-Schmidt group is given in [Arnold 01].

This paper is devoted to showing that 10 is the minimum such rank, i.e. if $M \in TF$ with $\text{rank } M \leq 9$, then M is a Krull-Schmidt group. Many arguments in this paper carry over directly to torsion-free modules of finite rank over valuation domains, keeping in mind that the existence and minimal rank of a non-Krull-Schmidt module depends on the structure of the valuation domain; see [Goldsmith May 99] and references.

The *quasi-isomorphism* category $TF_{\mathbb{Q}}$ of TF is an additive category with objects those of TF but with morphism sets $\mathbb{Q} \otimes \text{Hom}(M, N)$ for $M, N \in TF$ and \mathbb{Q} the rational numbers. The category $TF_{\mathbb{Q}}$ is a *Krull-Schmidt category* in that each object in $TF_{\mathbb{Q}}$ can be written uniquely, up to isomorphism in $TF_{\mathbb{Q}}$ and order, as a finite direct sum of indecomposable objects in $TF_{\mathbb{Q}}$; see [Walker 64]. This is because

Received by the editors October 4, 2000 and, in revised form, January 8, 2001.

2000 *Mathematics Subject Classification.* Primary 20K15, 20K25.

Key words and phrases. Krull-Schmidt groups, direct sum decompositions, local torsion-free abelian groups.

This research was supported, in part, by the Baylor University Summer Sabbatical Program.

an indecomposable object M in $TF_{\mathbb{Q}}$ has a local endomorphism ring $\mathbb{Q}\text{End} M$ in $TF_{\mathbb{Q}}$. Indecomposable objects in $TF_{\mathbb{Q}}$ are called *strongly indecomposable groups*, isomorphism in $TF_{\mathbb{Q}}$ is called *quasi-isomorphism*, and summands of groups in $TF_{\mathbb{Q}}$ are called *quasi-summands*.

Each $M \in TF$ is a torsion-free $\mathbb{Z}_{(p)}$ -module, where $\mathbb{Z}_{(p)}$ is the localization of the integers at the prime p . The rank of M as a group is equal to the rank of M as a $\mathbb{Z}_{(p)}$ -module, and $\text{Hom}_{\mathbb{Z}}(M, N) = \text{Hom}_{\mathbb{Z}_{(p)}}(M, N)$ for each $M, N \in TF$. Hence, $\text{rank } M = 1$ if and only if M is isomorphic to either $\mathbb{Z}_{(p)}$ or \mathbb{Q} . Define p -rank M to be the $\mathbb{Z}/p\mathbb{Z}$ -dimension of M/pM , a finite dimensional $\mathbb{Z}/p\mathbb{Z}$ -vector space. Notice that p -rank $M \leq \text{rank } M$, M is divisible if and only if p -rank $M = 0$, and M is isomorphic to a free $\mathbb{Z}_{(p)}$ -module if and only if p -rank $M = \text{rank } M$. Moreover, if N is a $\mathbb{Z}_{(p)}$ -submodule of M , then the p -rank of the pure submodule of M generated by N is less than or equal to the p -rank of N and if M is quasi-isomorphic to $N \oplus K$, then p -rank $M = p$ -rank $N + p$ -rank K [Arnold 72]. If M is *reduced* (no proper divisible subgroups), then M is isomorphic to a pure subgroup of M^* , the completion of M in the p -adic topology. Moreover, M^* is a free \mathbb{Z}^* -module with rank equal to the p -rank of M , where \mathbb{Z}^* is the p -adic completion of $\mathbb{Z}_{(p)}$. Each endomorphism of M lifts to a unique \mathbb{Z}^* -endomorphism of M^* , whence $\text{End } M$ is a pure subring of $\text{End}_{\mathbb{Z}^*} M^*$.

2. UNIQUENESS OF DIRECT SUMS

A group $M \in TF$ has the *one-sided UDS property* if whenever $M \oplus N$ is isomorphic to $K_1 \oplus \dots \oplus K_n \in TF$ with M quasi-isomorphic to each K_j , then M is isomorphic to some K_j . The group M has the *UDS property* if whenever $N_1 \oplus \dots \oplus N_m$ is isomorphic to $K_1 \oplus \dots \oplus K_n \in TF$ with M quasi-isomorphic to each N_i and K_j , then $m = n$ and there is a relabelling of indices with each N_i isomorphic to K_i .

Given a strongly indecomposable M in TF , there is a faithful $G_M \in TF$ quasi-isomorphic to M such that $\text{End } G_M / N\text{End } G_M$ is a maximal order in the division algebra $\mathbb{Q}\text{End } M / J\mathbb{Q}\text{End } M$, where $J\mathbb{Q}\text{End } M$ is the Jacobson radical of the finite dimensional \mathbb{Q} -algebra $\mathbb{Q}\text{End } M$, $N\text{End } G_M = \text{End } G_M \cap J\mathbb{Q}\text{End } M$ is a nilpotent ideal of $\text{End } M$, and G_M is *faithful* if $IG_M \neq G_M$ for each maximal right ideal I of $\text{End } G_M$ [Arnold 01]. A maximal right ideal J of $\text{End } G_M / p\text{End } G_M$ has the *unique maximal condition* if whenever I is a non-zero right ideal and J is a unique maximal right ideal of $\text{End } G_M / p\text{End } G_M$ containing I , then $I = J$.

The first lemma is the local version of [Arnold 01, Theorem 1.5].

Lemma 1. *The following statements are equivalent for a strongly indecomposable N in TF , where G_N is as defined above:*

- (i) *N has the UDS property.*
- (ii) *Each group in TF quasi-isomorphic to N has the one-sided UDS property.*
- (iii) *Either $\text{End } G_N$ is a local ring or else $\text{End } G_N$ has exactly two maximal right ideals M_1 and M_2 such that M_1 is a principal right ideal of $\text{End } G_N$, $G_N / M_1 G_N \cong \mathbb{Z}/p\mathbb{Z}$, and $M_1 / p\text{End } G_N$ has the unique maximal condition in $\text{End } G_N / p\text{End } G_N$.*

Following are non-trivial examples of groups in TF with the UDS property.

Example 1. If $N \in TF$ is strongly indecomposable with p -rank $N \leq 2$, then N has the (one-sided) UDS property.

Proof. It suffices to confirm the conditions of Lemma 1(iii). If p -rank $N \leq 1$, then p -rank $G_N \leq 1$, since G_N is quasi-isomorphic to N . Thus, either p -rank $G_N = 0$ and $G_N \cong \mathbb{Q}$ or else p -rank $G_N = 1$, $G_N^* \cong \mathbb{Z}^*$, and $\text{End } G_N$ is isomorphic to a pure subring of $\mathbb{Z}^* \cong \text{End}_{\mathbb{Z}^*} \mathbb{Z}^*$. In either case, $\text{End } G_N$ is a local ring, as desired.

Now assume that p -rank $N = p$ -rank $G_N = 2$ and $\text{End } G_N$ is not a local ring. Let M_1, \dots, M_n be distinct maximal right ideals of $\text{End } G_N$ with $n \geq 2$. Then $G_N/(M_1 \cap \dots \cap M_n)G_N \cong G_N/M_1G_N \oplus \dots \oplus G_N/M_nG_N$ and $p\text{End } G_N \subseteq J\text{End } G_N \subseteq M_1 \cap \dots \cap M_n$. Since p -rank $G_N = 2$ and G_N is faithful, it follows that $n = 2$, $p\text{End } G_N = M_1 \cap M_2 = J\text{End } G_N$, each $G_N/M_iG_N \cong \mathbb{Z}/p\mathbb{Z}$, and each $M_i/p\text{End } G_N$ has the unique maximal condition. Finally, each M_i is principal as an application of Nakayama's Lemma, because $p\text{End } G_N = J\text{End } G_N$ and $\text{End } G_N/p\text{End } G_N$ is finite. \square

The next lemma is used for an induction step in the proof of the main theorem.

Lemma 2. *Assume that $M = N \oplus N' = K_1 \oplus \dots \oplus K_n \in TF$. There are subgroups K'_i of K_i with $N \oplus N' = N \oplus K'_1 \oplus \dots \oplus K'_n$ if either*

- (a) [Warfield 72] *End N is a local ring or*
- (b) [Arnold Lady 75] *N and N' have no quasi-summands in common.*

In this case, N' is isomorphic to $K'_1 \oplus \dots \oplus K'_n$.

An indecomposable $M \in TF$ is *purely indecomposable* if p -rank $M = 1$. In this case, $\text{End } M$ is a local ring, being a pure subring of $\mathbb{Z}^* \cong \text{End}_{\mathbb{Z}^*} \mathbb{Z}^*$. Dually, M is *co-purely indecomposable* if M is indecomposable with $\text{rank } M = p\text{-rank } M + 1$. There is a contravariant duality F on $TF_{\mathbb{Q}}$ sending a purely indecomposable group M to a co-purely indecomposable group $F(M)$ [Arnold 72] (see [Lady 77] for an alternate definition of the duality). Hence, $\mathbb{Q}\text{End } F(M)$ is isomorphic to $\mathbb{Q}\text{End } M$, a subring of the p -adic rationals \mathbb{Q}^* .

Following are some elementary properties of purely indecomposable and co-purely indecomposable groups that are consequences of the definitions and the duality F .

Proposition 1 ([Arnold 72]). *Let $M \in TF$.*

- (a) *If M is purely indecomposable, then:*
 - (i) *End M is a pure subring of \mathbb{Z}^* ;*
 - (ii) *each pure subgroup of M is strongly indecomposable;*
 - (iii) *if $K \in TF$ is a homomorphic image of M with $\text{rank } K < \text{rank } M$, then K is divisible; and*
 - (iv) *two purely indecomposable groups M and N in TF are isomorphic if and only if $\text{rank } M = \text{rank } N$ and $\text{Hom}(M, N) \neq 0$; equivalently M and N are quasi-isomorphic.*
- (b) *If M is co-purely indecomposable, then:*
 - (i) *End M is isomorphic to a subring of \mathbb{Q}^* , hence an integral domain;*
 - (ii) *each torsion-free homomorphic image of M is strongly indecomposable;*
 - (iii) *if K is a pure subgroup of M with $\text{rank } K < \text{rank } M$, then M is a free $\mathbb{Z}_{(p)}$ -module; and*
 - (iv) *two co-purely indecomposable groups M and N are quasi-isomorphic if and only if $\text{rank } M = \text{rank } N$ and $\text{Hom}(M, N) \neq 0$.*
- (c) *If M is indecomposable with $\text{rank } \geq 2$ and N is co-purely indecomposable with $\text{rank } M < \text{rank } N$, then $\text{Hom}(M, N) = 0$.*

- (d) If M is purely indecomposable with rank ≥ 3 and N is co-purely indecomposable with rank $M = \text{rank } N$, then $\text{Hom}(M, N) = 0$.

Remark 1. There is a co-purely indecomposable $M \in TF$ with p -rank 3 and rank 4 that does not have either UDS property. In this case $\text{End } M$ has 3 maximal right ideals and M is the summand of a non-Krull-Schmidt group of rank 12 [Arnold 01, Remark]. In view of the following lemma, this group cannot be a summand of a non-Krull-Schmidt group of rank $8 = 2(\text{rank } M)$. On the other hand, if $M \in TF$ and $\text{End } M$ has at least 4 maximal right ideals, then M is a summand of a non-Krull-Schmidt group of rank equal to $2(\text{rank } M)$.

The next lemma is used in the proof of the main theorem. In view of Proposition 1(b)(i), the hypotheses are satisfied if N is co-purely indecomposable.

Lemma 3. Assume that $N \in TF$ with $\text{End } N$ an integral domain and $M = N \oplus N' = K_1 \oplus K_2 \in TF$ with each K_i indecomposable. If $p\text{-rank } N \leq 3$, then N is isomorphic to some K_i .

Proof. The proof is a variation on a proof given in [Arnold 01]. Let π be a projection of M onto N with kernel N' , and π_i a projection of M onto K_i for each i with $1_M = \pi_1 + \pi_2$. Then $1_N = \beta_1 + \beta_2$, where $\beta_i \in \text{End } N$ is the restriction of π_i to N and $\beta_i(N)$ is contained in a subgroup $\pi(K_i)$ of N . Since $\text{End } N$ is an integral domain, $\mathbb{Q}\text{End } N$ is a field and each β_i is a unit in $\mathbb{Q}\text{End } N$.

For each $1 \leq i \leq 2$, let $I_i = \beta_i \text{End } N$, a right ideal of $\text{End } N$. Then $\text{End } N = I_1 + I_2$, since $1_N = \beta_1 + \beta_2$. Each $(\text{End } N)/I_i$ is bounded by a power of p since β_i is a unit in $\mathbb{Q}\text{End } N$. Moreover, $I_i N$ is contained in $A_i = \pi(K_i)$ so that $[N : A_i]$ is finite. It now suffices to prove that $N \cong A_i$ for some i , in which case $N \cong K_i$.

If some $[N : A_i] = 1$, then $N = A_i$ and the proof is complete. The next step is to assume that each $[N : A_i] \neq 1$ and reduce to the case that each $[N : A_i] = p$. Suppose, by way of induction, that $[N : A_i] \neq p$. Choose $x \in N \setminus A_i$ such that $px \in A_i$. Then $A_i \subset A_i + \mathbb{Z}x$. If N and $A_i + \mathbb{Z}x$ are not isomorphic, then replace A_i by $A'_i = A_i + \mathbb{Z}x$. If $N \cong A_i + \mathbb{Z}x$, say $f \in \text{End } N$ with $f(N) = A_i + \mathbb{Z}x$, then replace A_i by $A'_i = f^{-1}(A_i)$. In either case, $[N : A'_i]$ is a proper divisor of $[N : A_i]$.

The substitution of A'_i for A_i doesn't change the hypothesis that $\text{End } N = I_1 + I_2$ for right ideals I_i of bounded index with $I_i N$ contained in A_i . In particular, $I'_i = f^{-1}I_i$ is an ideal of $\text{End } N$ (since $I_i N$ is a subgroup of $f(N)$), $I'_i N$ is contained in A'_i , and $I'_i + \Sigma\{I_j : j \neq i\} = \text{End } N$ (since $f\text{End } N = \Sigma fI_i$ is contained in $I_1 + \Sigma\{fI_j : j \neq i\}$). If $N \cong A'_i$, then, by the construction of A'_i , $N \cong A_i$. By induction, and the fact that $[N : A'_i]$ is a proper divisor of $[N : A_i]$, the A_i 's can be chosen with each $[N : A_i] = p$.

At this stage, $\text{End } N = I_1 + I_2$ for right ideals I_i of finite index in $\text{End } N$ with $I_i N$ contained in a subgroup A_i of N and $[N : A_i] = p$ for each i . Replace I_i by $I_i + p\text{End } N$, if necessary, to guarantee that $p\text{End } N$ is contained in I_i for each i . But $p\text{-rank } N \leq 3$, $p\text{End } N \subseteq J\text{End } N$, $\text{End } N$ is an integral domain, and $N/(M_1 \cap \dots \cap M_n)N \cong N/M_1N \oplus \dots \oplus N/M_nN$ for maximal ideals M_i of $\text{End } N$. Hence, $\text{End } N$ has at most 3 maximal right ideals M_1, M_2 , and M_3 and $p\text{End } N = M_1^{i_1} M_2^{i_2} M_3^{i_3}$ with $i_1 + i_2 + i_3 \leq 3$. Furthermore, $pN \subseteq (I_1 \cap I_2)N$, and $N/(I_1 \cap I_2)N \cong N/I_1N \oplus N/I_2N$. After relabelling subscripts, if necessary, $I_1 = M_1$, $N/I_1N = \mathbb{Z}/p\mathbb{Z}$, and $I_1N = A_1$. Finally, I_1 is principal by Nakayama's

Lemma, since $p\text{End } N \subseteq J\text{End } N$ and $I_1/p\text{End } N$ is principal. This shows that A_1 is isomorphic to N , as desired. \square

The point of the next lemma, as used in the proof of the main theorem, is that Lemma 2(a) applies to a group quasi-isomorphic to a direct sum of two purely indecomposable groups of the same rank.

Lemma 4. *If $N \in TF$ is indecomposable and quasi-isomorphic to $A \oplus B$ for purely indecomposable groups A and B in TF with $\text{rank } A = \text{rank } B$, then $\text{Hom}(A, B) = 0 = \text{Hom}(B, A)$ and $\text{End } N$ is a local ring.*

Proof. Choose purely indecomposable pure subgroups A and B of N and some least positive integer i with $p^i N \subset A \oplus B \subset N$. Since $p\text{-rank } N = 2$, $N/p^i N \cong \mathbb{Z}/p^i \mathbb{Z} \oplus \mathbb{Z}/p^i \mathbb{Z}$ for some $1 \leq j \leq i$. Because A and B are purely indecomposable pure subgroups of N , $N/(A \oplus B) \cong \mathbb{Z}/p^j \mathbb{Z}$, say $N = A \oplus B + \mathbb{Z}(a, b)(1/p^j)$ for some $a \in A \setminus pA$ and $b \in B \setminus pB$.

If $\text{Hom}(A, B) \neq 0$ or $\text{Hom}(B, A) \neq 0$, then A and B are isomorphic by Proposition 1(iv). Moreover, $C = N/A$ is purely indecomposable and quasi-isomorphic to B . Hence, $C \cong A \cong B$ and $\text{Hom}(C, N)C = N$. By Baer's Lemma [Arnold 82], A is a summand of N , a contradiction to the assumption that N is indecomposable.

Now assume that $\text{Hom}(A, B) = 0 = \text{Hom}(B, A)$. Then A and B are fully invariant subgroups of $N = A \oplus B + \mathbb{Z}(a, b)(1/p^j)$. Thus, $\text{End } N$ is the pullback of a homomorphism $A \rightarrow \mathbb{Z}/p^j \mathbb{Z}$ with kernel $p^j A$ and a homomorphism $B \rightarrow \mathbb{Z}/p^j \mathbb{Z}$ with kernel $p^j B$. It follows that $\text{End } N/p^j \text{End } N \cong \mathbb{Z}/p^j \mathbb{Z}$, whence $\text{End } N$ is a local ring. \square

3. THE MAIN THEOREM

Theorem 1. *If $M \in TF$ and $\text{rank } M \leq 9$, then M is a Krull-Schmidt group.*

Proof. Let N be an indecomposable summand of M of minimal rank and $M = N \oplus N_1 \oplus \dots \oplus N_m = K_1 \oplus \dots \oplus K_n$ with each N_i and K_j indecomposable. Then $\text{rank } N \leq 4$, $\text{rank } N \leq \text{rank } N_j$, and $\text{rank } N \leq \text{rank } K_i$ for each i and j , since $\text{rank } M \leq 9$ and N is an indecomposable summand of M of minimal rank.

If $p\text{-rank } N \leq 1$, then $\text{End } N$ is a local ring, as noted above. In this case, by Lemma 2(a), $N_1 \oplus \dots \oplus N_m$ is isomorphic to $K'_1 \oplus \dots \oplus K'_n$ for subgroups K'_i of K_i . It follows, by an induction on the rank of M , that M is a Krull-Schmidt group. In particular, if $p\text{-rank } N = \text{rank } N$, then N is free and cyclic, hence of $p\text{-rank } 1$.

If N and $N_1 \oplus \dots \oplus N_m$ have no quasi-summands in common, then, by Lemma 2(b), the proof is completed by an induction on the rank of M .

In view of the preceding remarks, it is now sufficient to assume that M is reduced, $2 \leq p\text{-rank } N < \text{rank } N \leq 4$ for each indecomposable summand N of minimal rank, and if $M = N \oplus N'$, then N and $N' = N_1 \oplus \dots \oplus N_m$ have a quasi-summand in common. Under these assumptions, M has no rank-1 quasi-summands. This is because the only rank-1 groups in TF are $\mathbb{Z}_{(p)}$ and \mathbb{Q} and, since M is reduced, any rank-1 quasi-summand must actually be a summand isomorphic to $\mathbb{Z}_{(p)}$. The strategy of the remainder of the proof is to show that N must be isomorphic to some K_i , in which case the cancellation property for $N \in TF$ and an induction on the rank of M shows that M is a Krull-Schmidt group.

First assume that $\text{rank } N = 4$, $p\text{-rank } N = 3$. Then N , being indecomposable, is co-purely indecomposable, hence strongly indecomposable by Proposition 1. Thus,

$N_1 \oplus \dots \oplus N_m$ is quasi-isomorphic to $N \oplus L$ for some L of rank ≤ 1 . To see this, recall that N and $N_1 \oplus \dots \oplus N_m$ have a quasi-summand in common, N is strongly indecomposable, $\text{rank } N \geq \text{rank } N_i$, and $\text{rank } N + \sum_i \text{rank } N_i = 4 + \sum_i \text{rank } N_i \leq 9$. Since M has no rank-1 quasi-summands, $L = 0$, $m = 1$, and $n = 2$. But $TF_{\mathbb{Q}}$ is a Krull-Schmidt category so that $M = N \oplus N_1 = K_1 \oplus K_2$ has rank 8 with N quasi-isomorphic to N_1 , K_1 , and K_2 . By Lemma 3, N is isomorphic to either K_1 or K_2 , as desired.

Next, consider the case that $\text{rank } N = 4$ and $p\text{-rank } N = 2$. If N is strongly indecomposable, then, as above, $M = N \oplus N_1 = K_1 \oplus K_2$ has rank 8 and N is quasi-isomorphic to N_1 , K_1 , and K_2 . By Example 1, N has the UDS property so that N is isomorphic to either K_1 or K_2 , as desired. If N is not strongly indecomposable, then N is quasi-isomorphic to $A \oplus B$, where A and B are purely indecomposable groups with $p\text{-rank } 1$ and rank 2. This is because M has no rank-1 quasi-summands. Now apply Lemmas 2 and 4 and induction on the rank of M to see that M is a Krull-Schmidt group.

The only remaining case is that $p\text{-rank } N = 2$, $\text{rank } N = 3$. In this case N is co-purely indecomposable, hence strongly indecomposable by Proposition 1. Since N and N_1 have a quasi-summand in common, N_1 is quasi-isomorphic to $N \oplus A$ for some pure subgroup A of N_1 with $1 \leq p\text{-rank } A < \text{rank } A \leq 3$. This is because M has no rank-1 quasi-summands and $\text{rank } M \leq 9$.

If A has $p\text{-rank } 1$, then $\text{Hom}(A, N) = 0$ by Proposition 1(c) and (d), since A is purely indecomposable with $2 \leq \text{rank } A \leq 3 = \text{rank } N$, and N is co-purely indecomposable. In this case, $\text{Hom}(A, M) = \text{Hom}(A, A)$. It follows that A is a pure fully invariant subgroup, hence equal to a subgroup of some K_i , say K_1 . Thus, $N \oplus (N_1/A)$ is isomorphic to $(K_1/A) \oplus K_2 \oplus K_3$ and induction on the rank of M completes the proof.

Finally, assume that A has $p\text{-rank } 2$. Then $\text{rank } A = 3 = \text{rank } N$ and A and N are both co-purely indecomposable. If $\text{Hom}(A, N) = 0$, then, as above, M is a Krull-Schmidt group. Finally, if $\text{Hom}(A, N) \neq 0$, then A is quasi-isomorphic to N , since A and N are both co-purely indecomposable modules with the same rank. Hence, $M = N \oplus N_1 = K_1 \oplus K_2 \oplus \dots \oplus K_n$ has rank 9 with $n \leq 3$. If $n = 3$, then N is quasi-isomorphic to K_1, K_2 and K_3 by the minimality of the rank of N . In this case, Example 1 yields N isomorphic to some K_i . If $n = 2$, then, by Lemma 3, N is isomorphic to some K_i , as desired. \square

Example 2 ([Arnold 01]). There is a rank-10 group in TF that is not a Krull-Schmidt group.

Proof. The argument is briefly outlined. There is $M \in TF$ of $p\text{-rank } 4$ and rank 5 such that $M \cong \text{End } M$, a subring of an algebraic number field with exactly four maximal ideals M_1, M_2, M_3 , and M_4 , and $pM = p\text{End } M = M_1 \cap M_2 \cap M_3 \cap M_4$. Furthermore, there are subgroups A_1 and A_2 of M not isomorphic to M with $(M_1 \cap M_2)M \subset A_1$ and $(M_3 \cap M_4)M \subset A_2$. It follows that there is $B \in TF$ with $M \oplus B = A_1 \oplus A_2$, a rank 10 group in TF that is not a Krull-Schmidt group. \square

REFERENCES

- [Arnold 72] Arnold, D. A duality for torsion-free modules of finite rank over a discrete valuation ring, *Proc. Lond. Math. Soc.* (3) 24 (1972), 204-216.
- [Arnold 82] Arnold, D. *Finite Rank Torsion-Free Abelian Groups and Rings*, Lect. Notes in Math. 931, Springer-Verlag, New York, 1982. MR **84d**:20002

- [Arnold 00] Arnold, D. *Abelian Groups and Representations of Finite Partially Ordered Sets*, CMS Books in Mathematics, Springer-Verlag, New York, 2000. MR **2001g**:16030
- [Arnold 01] Arnold, D. Direct sum decompositions of torsion-free abelian groups of finite rank, *Abelian Groups, Rings, and Modules* (Proc. of 2000 Perth Conf.), Cont. Math., AMS, Providence, Rhode Island, 2001, 65-74. CMP 2001:09
- [Arnold Dugas 00] Arnold, D. and Dugas, M. Co-purely indecomposable modules over a discrete valuation ring, *J. Pure and Appl. Alg.* 161 (2001), 1-12.
- [Arnold Lady 75] Arnold, D. and Lady, E.L. Endomorphism rings and direct sums of torsion-free abelian groups, *Trans. Amer. Math. Soc.* 211 (1975), 225-237. MR **54**:5370
- [Fuchs 73] Fuchs, L. *Infinite Abelian Groups*, Vol. II, Academic Press, New York, 1973. MR **50**:2362
- [Goldsmith May 99] Goldsmith, B. and May, W. The Krull-Schmidt problem for modules over valuation domains, *J. Pure and Appl. Alg.* 140 (1999), 57-63. MR **2000d**:13016
- [Lady 75] Lady, E.L. Nearly isomorphic torsion-free abelian groups, *J. Alg.* 35 (1975), 235-238. MR **51**:5801
- [Lady 77] Lady, E.L. Splitting fields for torsion-free modules over discrete valuation rings, *J. Alg.* 49 (1977), 261-275. MR **58**:22039
- [Walker 64] Walker, E.A. Quotient categories and quasi-isomorphisms of abelian groups, *Proc. Colloq. Abelian Groups*, Budapest, 1964, 147-162. MR **31**:2327
- [Warfield 72] Warfield, R.B. Jr. Exchange rings and decompositions of modules, *Math. Ann.* 199 (1972), 31-36. MR **48**:11218

DEPARTMENT OF MATHEMATICS, BAYLOR UNIVERSITY, WACO, TEXAS 76798-7328

E-mail address: David_Arnold@baylor.edu