ON THE PREDICTABILITY OF DISCRETE DYNAMICAL SYSTEMS

NILSON C. BERNARDES JR.

(Communicated by Alan Dow)

ABSTRACT. Let X be a metric space. A function $f : X \rightarrow X$ is said to be non-sensitive at a point $a \in X$ if for every $\epsilon > 0$ there is a $\delta > 0$ such that for any choice of points $a_0 \in B(a; \delta)$, $a_1 \in B(f(a_0); \delta)$, $a_2 \in B(f(a_1); \delta), \dots$, we have that $d(a_m, f^m(a)) < \epsilon$ for every $m \geq 0$. Let $H(X)$ be the set of all homeomorphisms from X onto X endowed with the topology of uniform convergence. The main goal of the present paper is to prove that for certain spaces X , “most” functions in $H(X)$ are non-sensitive at “most” points of X .

1. INTRODUCTION

Consider a discrete dynamical system (X, f) , where X is a metric space with metric d . If $a \in X$, the sequence $a, f(a), f^2(a), \dots$ can be thought of as the actual behaviour of the system (X, f) at a . However, in concrete situations, we are often unable to compute the initial condition a exactly. We just compute a value a_0 close to a . It may also be the case that we cannot compute $f(a_0)$ exactly, but just a value a_1 close to $f(a_0)$. Then we compute a value a_2 close to $f(a_1)$ and so on. In this way, we obtain a sequence $a_0, a_1, a_2, \dots$ that can be thought of as the predicted behaviour of the system (X, f) at a . It is natural to ask whether or not this predicted behaviour is close to the actual behaviour of the system. This leads to the following definition:

Definition 1. We say that f is non-sensitive at a if for every $\epsilon > 0$ there is a $\delta > 0$ such that for any choice of points

$$a_0 \in B(a; \delta), a_1 \in B(f(a_0); \delta), a_2 \in B(f(a_1); \delta), \dots,$$

we have that

$$d(a_m, f^m(a)) < \epsilon \quad \text{for every } m \geq 0.$$

If f is non-sensitive at a , then the discrete dynamical system (X, f) is “predictable at a ”, in the sense that we can predict the future evolution of a in the system forever as accurately as we want provided we can compute the initial condition and the values of f precisely enough.

The above definition may remind the reader of the well-known notion of “shadowing” which, in the case X is compact, may be defined as follows [1]: f is said

Received by the editors August 10, 1999 and, in revised form, January 16, 2001.

2000 *Mathematics Subject Classification.* Primary 37B25, 37B20, 54H20.

Key words and phrases. Homeomorphisms, predictability, recurrence, Baire category, Lebesgue measure.

to have the shadowing property (also called the pseudo-orbit tracing property) if for every $\epsilon > 0$ there is a $\delta > 0$ such that for every sequence $a_0, a_1, a_2, \dots$ in X satisfying $d(f(a_i), a_{i+1}) < \delta$ for all $i \geq 0$, there is a point $x \in X$ such that

$$d(a_m, f^m(x)) < \epsilon \quad \text{for all } m \geq 0.$$

There is an important difference between “shadowing” and “non-sensitivity” despite the obvious fact that the former is a global notion whereas the latter is a pointwise one. To understand this difference, let $a \in X$ and choose points $a_0 \in B(a; \delta)$, $a_1 \in B(f(a_0); \delta)$, $a_2 \in B(f(a_1); \delta)$, etc. The shadowing property gives us the existence of a point $x \in X$ (possibly different from a) such that $d(a_m, f^m(x)) < \epsilon$ for all $m \geq 0$. On the other hand, non-sensitivity at a guarantees that this holds with $x = a$. This is a major difference if we are interested in the problem of predictability: the shadowing property guarantees only that the predicted behaviour of the system (X, f) at a is close to some actual behaviour of the system (but this actual behaviour may be different from the actual behaviour at the initial condition a !). For results and further references on “shadowing”, see the book [1] by Aoki and Hiraide.

Let $H(X)$ be the set of all homeomorphisms from X onto X . If X is compact, we consider $H(X)$ endowed with the supremum metric: $d(f, g) = \sup_{x \in X} d(f(x), g(x))$; $H(X)$ is then a Baire space.

If M is a Baire space, we say that “most elements of M ” satisfy a certain property P if the set of all $x \in M$ that do not satisfy property P is of the first category in M . The terms “typical” and “generic” are often used instead of “most”.

A result related to the notion of predictability was obtained by the author in [2]. Let B^n denote the closed unit ball of $\mathbf{R}^n$. Then, Theorem 20 of [2] can be stated as follows:

Let $n \geq 2$. For most functions $f \in H(B^n)$, the family $\{f^m; m \geq 1\}$ is equicontinuous at most points of B^n .

The equicontinuity of $\{f^m; m \geq 1\}$ at a point a can be thought of as a kind of “predictability with respect to the initial condition a ”. This property is much weaker than the notion of non-sensitivity (which implies that the family $\{f^m; m \geq 1\}$ is uniformly equicontinuous on the orbit $\{a, f(a), f^2(a), \dots\}$ of a under f). Nevertheless, it is indeed true that we can assure the non-sensitivity of f at most points of B^n , for most $f \in H(B^n)$. In fact, this is true in a much more general setting:

Theorem 2. *Fix $n \geq 1$. Let X be a metrizable compact topological n -manifold with (or without) boundary [5] and fix a metric d compatible with the topology of X . Then, most functions in $H(X)$ are non-sensitive at most points of X .*

We shall prove Theorem 2 in Section 4.

In the case X is a metrizable compact smooth manifold without boundary, Odani [6] proved that most functions in $CLD(X)$ (the closure in $H(X)$ of the set of all diffeomorphisms of X) have the shadowing property. His arguments were based on tools from Differentiable Dynamics. This work was preceeded by Yano [9], who proved that most functions in $H(S^1)$ have the shadowing property (where S^1 is the unit circle). By Munkres [4] and Whitehead [8], $CLD(X) = H(X)$ whenever $\dim X \leq 3$; but Munkres [4] also showed that this is false if $\dim X > 3$. The problem of whether most functions in $H(X)$ have the shadowing property was recently solved in the affirmative for $\dim X$ arbitrary by Pilyugin and Plamenevskaya [7]. However, this problem seems to remain open if X is an arbitrary metrizable compact topological manifold with (or without) boundary.

An interesting problem is to change the notion of “most” on B^n from category sense to measure-theoretic (probability) sense, that is, to consider “most” as meaning “full measure”. In this direction we have the following result:

Theorem 3. *Fix $n \geq 2$. Most functions in $H(B^n)$ are non-sensitive at every point of a full (Lebesgue) measure subset of B^n .*

As an immediate consequence of this theorem we have the following:

Corollary 4. *Let $n \geq 2$. For most functions $f \in H(B^n)$, if f is sensitive to initial conditions on a subset Y of B^n , then $\bar{Y}$ has Lebesgue measure zero.*

Recall that f is sensitive to initial conditions on $Y \subset B^n$ if there is an $\epsilon > 0$ such that for any $y \in Y$ and any $\delta > 0$, there is an $x \in B^n$ with $\|x - y\| < \delta$ and there is an $m \geq 1$ so that $\|f^m(x) - f^m(y)\| \geq \epsilon$.

We shall prove Theorem 3 in Section 2. Its proof will establish the following result at the same time:

Theorem 5. *Let $n \geq 2$. For most functions $f \in H(B^n)$, the set Ω_f of all non-wandering points of f has Lebesgue measure zero.*

Theorem 5 improves Theorem 17 of [2], which asserts that for most functions $f \in H(B^n)$, the set P_f of all periodic points of f has Lebesgue measure zero. In this direction, it is interesting to observe the following fact:

Proposition 6. *Fix $n \geq 2$. For most functions $f \in H(B^n)$, most points of Ω_f are recurrent and non-periodic.*

We shall prove Proposition 6 in Section 3.

2. PROOF OF THEOREMS 3 AND 5

Consider $\mathbf{R}^n$ endowed with the metric given by its euclidean norm $\|\cdot\|$. Given $A \subset \mathbf{R}^n$, $\bar{A}$, $\text{Int}A$, $\text{Bd}A$ and $\text{diam}A$ denote the closure, the interior, the boundary and the diameter of A in $\mathbf{R}^n$, respectively, and we define

$$N_\delta(A) = \bigcup_{a \in A} \{x \in \mathbf{R}^n; \|x - a\| < \delta\} \quad \text{for } \delta > 0.$$

Throughout the present section, X denotes the closed unit ball of $\mathbf{R}^n$ and μ denotes Lebesgue measure on $\mathbf{R}^n$. By an open box we mean a set of the form

$$\{(x_1, \dots, x_n) \in \mathbf{R}^n; \alpha_i < x_i < \alpha_i + \delta \text{ for } 1 \leq i \leq n\},$$

where $\alpha_1, \dots, \alpha_n \in \mathbf{R}$ and $\delta > 0$. A closed box is just the closure of an open box. By a tree we mean a finite rooted tree [3]. If T is a tree, $V(T)$ denotes the set of all vertices of T . Moreover, if $v_1, v_2 \in V(T)$, we write “ $v_1 > v_2$ ” or “ $v_2 < v_1$ ” to mean that v_1 and v_2 are adjacent [3] and that the unique path connecting v_2 to the root of T passes through v_1 . A B -tree is a pair (T, φ) , where T is a tree and φ is a bijective correspondence between $V(T)$ and a collection of pairwise disjoint closed boxes contained in $\text{Int}X$. If (T, φ) is a B -tree, we usually omit the correspondence φ and speak just of the B -tree T ; moreover, we make no distinction between a vertex of T and its corresponding closed box.

For each $k \geq 1$, let A_k be the set of all functions $f \in H(X)$ for which there are finitely many B -trees $T_1, \dots, T_s$ so that the following properties are satisfied:

- (i) $C \cap F = \emptyset$ whenever $C \in V(T_i)$, $F \in V(T_j)$ and $i \neq j$.

- (ii) $\text{diam} C < 1/k$ whenever $C \in V(T_i)$ for some $1 \leq i \leq s$.
- (iii) If $C, F \in V(T_i)$ and $C > F$, then $f(F) \subset \text{Int} C$.
- (iv) For each i , there is a chain $C_{i,1} > C_{i,2} > \cdots > C_{i,t_i}$ of successive boxes in $V(T_i)$, beginning with the root $C_{i,1}$ of T_i , so that $f(C_{i,1}) \subset \text{Int} C_{i,t_i}$.
- (v) The set

$$\bigcup \{C; C \in V(T_i) - \{C_{i,1}, \dots, C_{i,t_i}\} \text{ for some } 1 \leq i \leq s\}$$

together with the set

$$\bigcup_{i=1}^s [(C_{i,t_i} - f(C_{i,1})) \cup (C_{i,t_i-1} - f^2(C_{i,1})) \cup \cdots \cup (C_{i,1} - f^{t_i}(C_{i,1}))]$$

form a set of Lebesgue measure $> \mu(X) - 1/k$.

It is immediate to check that each A_k is open in $H(X)$. Moreover, $\mu(\Omega_f) < 1/k$ for every $f \in A_k$, and so $\mu(\Omega_f) = 0$ whenever $f \in \bigcap_{k=1}^{\infty} A_k$. Now, suppose $f \in \bigcap_{k=1}^{\infty} A_k$. Then, for each $k \geq 1$, there are B -trees $T_{k,1}, \dots, T_{k,s_k}$ so that (i) through (v) hold with $T_{k,1}, \dots, T_{k,s_k}$ in place of $T_1, \dots, T_s$. Put

$$M_k = \bigcup \{C; C \in V(T_{k,j}) \text{ for some } 1 \leq j \leq s_k\} \quad (k \geq 1) \quad \text{and} \quad M = \bigcap_{r=1}^{\infty} \bigcup_{k=r}^{\infty} M_k.$$

Since $\mu(M_k) > \mu(X) - 1/k$ for every $k \geq 1$, it follows that $\mu(M) = \mu(X)$. Moreover, for each $k \geq 1$, there is a $0 < \delta_k < 1/k$ such that

$$f(N_{\delta_k}(F)) \subset \text{Int} C \quad \text{whenever } C > F,$$

where $C, F \in V(T_{k,j})$ for some $1 \leq j \leq s_k$, and

$$f(N_{\delta_k}(C_{k,j,1})) \subset \text{Int} C_{k,j,t_{k,j}},$$

where $C_{k,j,1} > \cdots > C_{k,j,t_{k,j}}$ is related to $T_{k,j}$ ($1 \leq j \leq s_k$) as $C_{i,1} > \cdots > C_{i,t_i}$ is related to T_i in property (iv). Hence, if $a \in M_k$ and we choose $a_0 \in B(a; \delta_k)$, $a_1 \in B(f(a_0); \delta_k)$, $a_2 \in B(f(a_1); \delta_k)$, $\dots$, then $\|a_m - f^m(a)\| < 2/k$ for every $m \geq 0$. This implies that f is non-sensitive at every point of M .

It remains to show that each A_k is dense in $H(X)$. Fix $k \geq 1$, $f \in H(X)$ and $\epsilon > 0$. Let $0 < \delta < 1/k$ be such that

$$\text{diam} f(Y) < \frac{\epsilon}{2} \quad \text{whenever } Y \subset X \text{ and } \text{diam} Y < \delta.$$

Let $\mathcal{C}$ be a finite collection of pairwise disjoint closed boxes contained in $\text{Int} X$ such that

$$\mu(X - \bigcup_{C \in \mathcal{C}} C) < \frac{1}{k} \quad \text{and} \quad \text{diam} C < \delta \text{ for every } C \in \mathcal{C}.$$

Let r be the cardinality of $\mathcal{C}$ and choose $\eta > 0$ such that

$$\text{diam} f(Y) < \frac{\epsilon}{2^{r+1}} \quad \text{whenever } Y \subset X \text{ and } \text{diam} Y < \eta.$$

Our next goal is to construct finitely many B -trees $T_1, \dots, T_s$ satisfying (i) and the following properties:

- (a) $\text{diam} C < \delta$ whenever $C \in V(T_i)$ for some $1 \leq i \leq s$.
- (b) $\mathcal{C} \subset V(T_1) \cup \cdots \cup V(T_s)$.
- (c) If $C, F \in V(T_i)$ and $C > F$, then we have two possibilities:
 - either $C \in \mathcal{C}$ and $f(F) \subset C$,
 - or $C \notin \mathcal{C}$ and $(\text{Int} C) \cap f(\text{Bd} F) \neq \emptyset$.

- (d) For each i , there is a chain $C_{i,1} > C_{i,2} > \cdots > C_{i,t_i}$ of successive boxes in $V(T_i)$, beginning with the root $C_{i,1}$ of T_i , with the following property: $f(C_{i,1}) \subset C_{i,t_i}$ and/or there is a $C_{i,t_i+1} \in V(T_i)$ with $C_{i,t_i} > C_{i,t_i+1}$ such that $C_{i,1}, \dots, C_{i,t_i+1} \notin \mathcal{C}$ and both $C_{i,1}$ and C_{i,t_i+1} are contained in an open box W_i of diameter $< \eta$.

In order to explain how to construct the trees $T_1, \dots, T_s$, we shall need a variable $\mathcal{B}$, which will denote the set of all closed boxes that have already been used in the construction up to the current step (at the beginning we have $\mathcal{B} = \emptyset$). It is also important to observe that during the construction the trees will be seen as variables. Even when the construction of a tree T_i is apparently done, it may be necessary to change it later on in the process.

We begin by choosing a closed box $C_1 \in \mathcal{C}$ and by putting it as a vertex of T_1 (note that now $\mathcal{B} = \{C_1\}$). Suppose that in a certain moment T_1 consists of the vertices $C_1 < C_2 < \cdots < C_j$. We look at the set $f(C_j)$. There are three possibilities:

Case 1. $f(C_j) \subset \mathcal{C}$ for some $C \in \mathcal{B}$.

We stop the construction of T_1 (for the time being). So, C_j is the root of T_1 .

Case 2. $f(C_j) \subset \mathcal{C}$ for some $C \in \mathcal{C} - \mathcal{B}$.

Let C_{j+1} be such a C . We put C_{j+1} as a vertex of T_1 adjacent to C_j and satisfying $C_j < C_{j+1}$.

Case 3. $f(C_j) \not\subset \bigcup\{C; C \in \mathcal{C} \text{ or } C \in \mathcal{B}\}$.

Then we also have that $f(\text{Bd}C_j) \not\subset \bigcup\{C; C \in \mathcal{C} \text{ or } C \in \mathcal{B}\}$. Hence, we can choose a closed box $C_{j+1} \subset \text{Int}X$ disjoint from each box in $\mathcal{C} \cup \mathcal{B}$ such that $\text{diam}C_{j+1} < \delta$ and $(\text{Int}C_{j+1}) \cap f(\text{Bd}C_j) \neq \emptyset$. We put C_{j+1} as a vertex of T_1 adjacent to C_j and satisfying $C_j < C_{j+1}$.

If Case 1 never happens, then the construction can go on forever. In this case we will stop the construction of T_1 as soon as we obtain a chain $C_{1,1} > C_{1,2} > \cdots > C_{1,t_1} > C_{1,t_1+1}$ beginning with the root $C_{1,1}$ of T_1 so that $C_{1,1}, \dots, C_{1,t_1+1} \notin \mathcal{C}$ and both $C_{1,1}$ and C_{1,t_1+1} are contained in an open box W_1 of diameter $< \eta$. Since $\mathcal{C}$ is finite and X is bounded, we will obtain such a chain in a finite number of steps.

Suppose that we have already constructed $T_1, \dots, T_{i-1}$. If $\mathcal{B} \supset \mathcal{C}$, we are done. If this is not the case, we choose a $C'_1 \in \mathcal{C} - \mathcal{B}$ and put it as a vertex of T_i . If in a certain moment T_i consists of the vertices $C'_1 < C'_2 < \cdots < C'_j$, we then look at $f(C'_j)$. Cases 2 and 3 are treated as before. However, Case 1 should be divided in two possibilities:

Case 1a. $f(C'_j) \subset \mathcal{C}$ for some $C \in V(T_i)$.

We stop the construction of T_i (for the time being). So, C'_j is the root of T_i .

Case 1b. $f(C'_j) \subset \mathcal{C}$ for some $C \in \mathcal{B} - V(T_i)$.

Let $\tilde{C}$ be such a C . Then $\tilde{C}$ is a vertex of a previous tree, say $\tilde{C} \in V(T_{i_0})$, where $1 \leq i_0 < i$. In this case we will have no tree T_i for the time being. We will just enlarge T_{i_0} by putting the chain $C'_1 < C'_2 < \cdots < C'_j$ as a new branch of it, satisfying the relation $C'_j < \tilde{C}$.

If Cases 1a and 1b never happen, we will stop the construction of T_i when we obtain a chain $C_{i,1} > C_{i,2} > \dots > C_{i,t_i} > C_{i,t_i+1}$ beginning with the root $C_{i,1}$ of T_i so that $C_{i,1}, \dots, C_{i,t_i+1} \notin \mathcal{C}$ and both $C_{i,1}$ and C_{i,t_i+1} are contained in an open box W_i of diameter $< \eta$.

By the way the trees $T_1, \dots, T_s$ were constructed, it is immediate to check that they have all the desired properties. Note also that $s \leq r$.

Let $I = \{i \in \{1, \dots, s\}; f(C_{i,1}) \subset C_{i,t_i}\}$ and $J = \{1, \dots, s\} - I$. For each $C \in V(T_1) \cup \dots \cup V(T_s)$, choose an open box V_C with

$$\text{diam} V_C < \delta \quad \text{and} \quad C \subset V_C \subset \overline{V_C} \subset \text{Int} X,$$

so that the family $\{\overline{V_C}\}_{C \in V(T_1) \cup \dots \cup V(T_s)}$ is pairwise disjoint. We may also assume that

$$\overline{V_{C_{i,1}}} \subset W_i \quad \text{and} \quad \overline{V_{C_{i,t_i+1}}} \subset W_i \quad \text{for every } i \in J.$$

Now, we are going to define a function $g_0 \in H(X)$ as follows: Suppose that $C, F \in V(T_i)$ for some i , and $C > F$. By (c), there is a closed box $B \subset \text{Int} F$ such that $f(B) \subset \text{Int} C$. If $F \notin \{C_{i,t_i+1}; i \in J\}$, choose a $\varphi \in H(\overline{V_F})$ such that $\varphi(F) \subset B$ and $\varphi(x) = x$ for all $x \in \text{Bd} V_F$, and define

$$g_0(x) = f(\varphi(x)) \quad \text{for all } x \in \overline{V_F}.$$

If $F \in \{C_{i,t_i+1}; i \in J\}$, then $C \notin \mathcal{C}$ and so $(\text{Int} C) \cap f(\text{Bd} F) \neq \emptyset$. Choose an open box Z_F with $F \subset Z_F \subset \overline{Z_F} \subset V_F$ such that there is a closed box $D_F \subset V_F - \overline{Z_F}$ with $f(D_F) \subset \text{Int} C$, choose a $\varphi \in H(\overline{Z_F})$ such that $\varphi(F) \subset B$ and $\varphi(x) = x$ for all $x \in \text{Bd} Z_F$, and define

$$g_0(x) = f(\varphi(x)) \quad \text{for all } x \in \overline{Z_F}.$$

We make this definition for all $C, F \in V(T_i)$ with $C > F$ ($1 \leq i \leq s$). For $i \in I$, there also exists a closed box $B \subset \text{Int} C_{i,1}$ such that $f(B) \subset \text{Int} C_{i,t_i}$. So, choose a $\varphi \in H(\overline{V_{C_{i,1}}})$ such that $\varphi(C_{i,1}) \subset B$ and $\varphi(x) = x$ for all $x \in \text{Bd} V_{C_{i,1}}$, and put

$$g_0(x) = f(\varphi(x)) \quad \text{for all } x \in \overline{V_{C_{i,1}}}.$$

Let K be the union of all these closed boxes where we have already defined g_0 . We finally put $g_0(x) = f(x)$ for all $x \in X - K$. Then $g_0 \in H(X)$, $d(g_0, f) < \epsilon/2$, and (iii) holds for every $i \in \{1, \dots, s\}$ and (iv) holds for every $i \in I$ with g_0 in place of f .

Now, we need to change g_0 a little bit in order to obtain (iv) also for $i \in J$. If $i \in J$ and $1 \leq j \leq t_i + 1$, we denote the open box $V_{C_{i,j}}$ simply by $V_{i,j}$. Moreover, for $i \in J$, we write Z_i and D_i in place of $Z_{C_{i,t_i+1}}$ and $D_{C_{i,t_i+1}}$, respectively. Recall that

$$D_i \subset V_{i,t_i+1} - \overline{Z_i} \quad \text{and} \quad f(D_i) \subset \text{Int} C_{i,t_i} \quad (i \in J).$$

Write $J = \{i_1, \dots, i_w\}$ and put

$$K_1 = K \cup \overline{V_{i_2,1}} \cup \dots \cup \overline{V_{i_w,1}} \cup D_{i_2} \cup \dots \cup D_{i_w}.$$

Choose

$$a_1 \in \text{Int} C_{i_1,1} \subset \overline{V_{i_1,1}} \subset W_{i_1} - K_1 \quad \text{and} \quad b_1 \in \text{Int} D_{i_1} \subset W_{i_1} - K_1.$$

Since K_1 is the union of a finite collection of pairwise disjoint closed boxes contained in $\text{Int} X$, $W_{i_1} - K_1$ is connected. So, there is a continuous path $\alpha : [0, 1] \rightarrow W_{i_1} - K_1$ from a_1 to b_1 . Moreover, we may assume that $\alpha([0, 1]) \subset \text{Int} X$. Cover

$\alpha([0, 1])$ by finitely many open balls $B_1, \dots, B_\ell$ whose closures are contained in $(W_{i_1} - K_1) \cap \text{Int}X$ so that

$$\overline{B_1} \subset \text{Int}C_{i_1,1}, \quad \overline{B_\ell} \subset \text{Int}D_{i_1} \quad \text{and} \quad B_i \cap B_{i+1} \neq \emptyset \quad \text{for every } 1 \leq i < \ell.$$

By working on $\overline{V_{i_1,1}} \cup \overline{B_1} \cup \dots \cup \overline{B_\ell}$, we see that it is possible to construct a $\varphi \in H(X)$ such that

$$\varphi(C_{i_1,1}) \subset \text{Int}D_{i_1} \quad \text{and} \quad \varphi(x) = x \quad \text{if } x \notin \overline{V_{i_1,1}} \cup \overline{B_1} \cup \dots \cup \overline{B_\ell}.$$

Put $g_1 = g_0 \circ \varphi$. Then $g_1 \in H(X)$, $g_1 = g_0$ on $X - (\overline{V_{i_1,1}} \cup \overline{B_1} \cup \dots \cup \overline{B_\ell})$ and

$$g_1(C_{i_1,1}) \subset g_0(D_{i_1}) = f(D_{i_1}) \subset \text{Int}C_{i_1,t_{i_1}}.$$

Moreover, since $\text{diam}(\overline{V_{i_1,1}} \cup \overline{B_1} \cup \dots \cup \overline{B_\ell}) < \eta$, we have $d(g_1, g_0) < \epsilon/2^{r+1}$ and

$$\text{diam}g_1(Y) < \frac{\epsilon}{2^r} \quad \text{whenever } Y \subset X - K \text{ and } \text{diam}Y < \eta.$$

Put

$$K_2 = K \cup \overline{V_{i_1,1}} \cup \overline{V_{i_3,1}} \cup \dots \cup \overline{V_{i_w,1}} \cup D_{i_3} \cup \dots \cup D_{i_w}.$$

Choose

$$a_2 \in \text{Int}C_{i_2,1} \subset \overline{V_{i_2,1}} \subset W_{i_2} - K_2 \quad \text{and} \quad b_2 \in \text{Int}D_{i_2} \subset W_{i_2} - K_2,$$

and argue as before. We then obtain a $g_2 \in H(X)$ with $g_2(C_{i_2,1}) \subset \text{Int}C_{i_2,t_{i_2}}$. Moreover, since g_2 differs from g_1 only on a set of diameter $< \eta$, we have $d(g_2, g_1) < \epsilon/2^r$ and

$$\text{diam}g_2(Y) < \frac{\epsilon}{2^{r-1}} \quad \text{if } Y \subset X - K \text{ and } \text{diam}Y < \eta.$$

By continuing this process, we will obtain a function $g_w \in H(X)$ such that

$$d(g_w, g_0) < \frac{\epsilon}{2^{r+1-(w-1)}} + \frac{\epsilon}{2^{r+1-(w-2)}} + \dots + \frac{\epsilon}{2^{r+1}} < \frac{\epsilon}{2},$$

and so $d(g_w, f) < \epsilon$. Moreover, properties (iii) and (iv) hold for every $i \in \{1, \dots, s\}$ if we replace f by g_w .

Finally, it remains to deal with property (v). Recall that $\mathcal{C} \subset V(T_1) \cup \dots \cup V(T_s)$ and that $\mu(\bigcup_{C \in \mathcal{C}} C) > \mu(X) - 1/k$. If we had every box of $\mathcal{C}$ appearing in the first union in property (v) we would be done. However, this may not be the case, since for $i \in I$ there may exist boxes of $\mathcal{C}$ in the chain $C_{i,1} > C_{i,2} > \dots > C_{i,t_i}$. For each $i \in I$, let U_i be an open box such that

$$C_{i,1} \subset U_i \subset \overline{U_i} \subset V_{C_{i,1}} \quad \text{and} \quad g_w(\overline{U_i}) \subset \text{Int}C_{i,t_i}.$$

Let $\varphi_i \in H(\overline{U_i})$ be such that $\varphi_i(x) = x$ for all $x \in \text{Bd}U_i$ and $\text{diam}\varphi_i(C_{i,1})$ is very small. Put $g = g_w \circ \varphi_i$ on $\overline{U_i}$ for each $i \in I$ and $g = g_w$ on $X - \bigcup_{i \in I} \overline{U_i}$. Then $g \in H(X)$ and $d(g, f) < \epsilon$, because $g_w(\overline{U_i}) \subset f(\overline{V_{C_{i,1}}})$ for each $i \in I$, by the way the g_j 's were constructed. Moreover, (iii) and (iv) still hold with g in place of f . By choosing φ_i so that $\text{diam}\varphi_i(C_{i,1})$ is small enough, we will have that the measure of

$$(C_{i,t_i} - g(C_{i,1})) \cup (C_{i,t_i-1} - g^2(C_{i,1})) \cup \dots \cup (C_{i,1} - g^{t_i}(C_{i,1}))$$

is so close to the measure of $C_{i,t_i} \cup C_{i,t_i-1} \cup \dots \cup C_{i,1}$ that (v) will also hold with g in place of f . This completes the proof.

3. PROOF OF PROPOSITION 6

Put $X = B^n$. We know that for most functions $f \in H(X)$, $\Omega_f = \overline{P_f}$ and the set of all periodic points of f with period m is dense in the set of all periodic points of f with period q whenever q divides m [2]. Fix an $f \in H(X)$ which has these two properties. Let R_f denote the set of all recurrent points of f . For each $m \geq 1$ and each $r \geq 1$, let $V_{m,r} = \{x \in X; \|x - f^t(x)\| < 1/m \text{ for some } t \geq r\}$. Then, each $V_{m,r}$ is open and

$$R_f - P_f = \bigcap_{m,r,k} (V_{m,r} - F_{f^k})$$

(where F_{f^k} denotes the set of all fixed points of f^k). Thus, it remains to show that each set $(V_{m,r} - F_{f^k}) \cap \Omega_f$ is dense in Ω_f . For this purpose, let U be an open set that meets Ω_f . Choose a $y \in P_f \cap U$. Let p be the period of y and choose an integer t of the form sp (for some $s \geq 1$) which is greater than k . Now, choose a periodic point z of f with period t which lies in U . Then, $z \in (V_{m,r} - F_{f^k}) \cap \Omega_f \cap U$, which completes the proof.

4. PROOF OF THEOREM 2

Let $i(X)$ denote the interior of the manifold X . If $A \subset X$, then $\overline{A}$ and $\text{Int}A$ denote the closure and the interior of A in X , respectively, and we define $N_\delta(A) = \bigcup_{a \in A} B(a; \delta)$ for $\delta > 0$. Moreover, if $f : X \rightarrow X$ is a mapping, we define $(f \circ N_\delta)^0(A) = A$, $(f \circ N_\delta)^1(A) = f(N_\delta(A))$, $(f \circ N_\delta)^2(A) = f(N_\delta(f(N_\delta(A))))$, and so on.

Fix a sequence $z_1, z_2, \dots$ in $i(X)$ which is dense in X . For each $r \geq 1$ and $k \geq 1$, let $\mathcal{O}_{r,k}$ be the set of all functions $f \in H(X)$ for which there is a closed set $V \subset i(X)$ and there are integers $q \geq 0$ and $m \geq 1$ so that $f^q(z_k) \in \text{Int}V$, $f^m(V) \subset \text{Int}V$ and $\text{diam}f^i(V) < 1/r$ for $0 \leq i \leq m-1$.

Clearly, each $\mathcal{O}_{r,k}$ is open. Let $f \in \bigcap_{r,k} \mathcal{O}_{r,k}$. By definition, for each $r \geq 1$ and each $k \geq 1$, there are a closed set $V_{r,k} \subset i(X)$ and integers $q_{r,k} \geq 0$ and $m_{r,k} \geq 1$ such that $f^{q_{r,k}}(z_k) \in \text{Int}V_{r,k}$, $f^{m_{r,k}}(V_{r,k}) \subset \text{Int}V_{r,k}$ and $\text{diam}f^i(V_{r,k}) < 1/r$ for $0 \leq i \leq m_{r,k}-1$. Let $W_{r,k}$ be an open ball centered at z_k such that

$$f^{q_{r,k}}(\overline{W_{r,k}}) \subset \text{Int}V_{r,k} \quad \text{and} \quad \text{diam}f^i(\overline{W_{r,k}}) < 1/r \quad \text{for } 0 \leq i \leq q_{r,k}-1.$$

Choose $0 < \delta_{r,k} < 1/r$ such that

$$(f \circ N_{\delta_{r,k}})^{q_{r,k}}(\overline{W_{r,k}}) \subset \text{Int}V_{r,k}, \quad (f \circ N_{\delta_{r,k}})^{m_{r,k}}(V_{r,k}) \subset \text{Int}V_{r,k},$$

$$\text{diam}(f \circ N_{\delta_{r,k}})^i(\overline{W_{r,k}}) < \frac{1}{r} - \delta_{r,k} \quad \text{for } 0 \leq i \leq q_{r,k}-1$$

and

$$\text{diam}(f \circ N_{\delta_{r,k}})^i(V_{r,k}) < \frac{1}{r} - \delta_{r,k} \quad \text{for } 0 \leq i \leq m_{r,k}-1.$$

Put $D_r = \bigcup_{k=1}^{\infty} W_{r,k}$. Then D_r is open and dense in X . Moreover, if $a \in D_r$, then $a \in W_{r,k}$ for some $k \geq 1$, and so for any choice of points $a_0 \in B(a; \delta_{r,k})$, $a_1 \in B(f(a_0); \delta_{r,k})$, $a_2 \in B(f(a_1); \delta_{r,k})$, $\dots$, we have that $d(a_m, f^m(a)) < 1/r$ for every $m \geq 0$. Therefore, $D = \bigcap_{r=1}^{\infty} D_r$ is a dense G_δ subset of X and f is non-sensitive at every point of D . Thus, it remains to show that each $\mathcal{O}_{r,k}$ is dense in $H(X)$. Fix $r \geq 1$, $k \geq 1$, $f \in H(X)$ and $\epsilon > 0$. Let a be a cluster point of the

sequence $(f^j(z_k))_{j \geq 0}$ and choose a neighborhood W of a in X for which there is a homeomorphism $\psi : W \rightarrow B^n$ with

$$\psi(i(X) \cap \text{Int}W) = \{x \in \mathbf{R}^n; \|x\| < 1\}.$$

Let $q \geq 0$ be the smallest integer such that $f^q(z_k) \in \text{Int}W$. Let $s \geq 1$ be such that $f^{q+s}(z_k) \in \text{Int}W$. We have two possibilities:

Case 1. $n \geq 2$.

We choose a point $b \in (i(X) \cap \text{Int}W) - \{f^j(z_k); j \in \mathbf{Z}\}$ so close to $f^q(z_k)$ that we have $f^s(b) \in \text{Int}W$, and we let $m \geq 1$ be the smallest integer such that $f^m(b) \in \text{Int}W$.

Case 2. $n = 1$.

We may think of W as being $[-1, 1]$. So, we may define the sets

$$L = \{x \in W; -1 < x < f^q(z_k)\} \quad \text{and} \quad R = \{x \in W; f^q(z_k) < x < 1\}.$$

If $f^{q+s}(z_k) = f^q(z_k)$, choose $b \in R - \{f^j(z_k); j \in \mathbf{Z}\}$ so close to $f^q(z_k)$ that we have $f^s(b)$ and $f^{2s}(b)$ in $\text{Int}W$; then $f^s(b)$ or $f^{2s}(b)$ belong to R . If $f^{q+s}(z_k) \neq f^q(z_k)$, then either $f^{q+s}(z_k) \in L$ or $f^{q+s}(z_k) \in R$; say $f^{q+s}(z_k) \in R$. In this case, choose $b \in R - \{f^j(z_k); j \in \mathbf{Z}\}$ so that $f^s(b) \in R$. Let $m \geq 1$ be the smallest integer such that $f^m(b) \in R \subset \text{Int}W$.

Let $\phi \in H(X)$ be such that

$$\phi(f^m(b)) = b \quad \text{and} \quad \phi(x) = x \quad \text{for all } x \in (X - \text{Int}W) \cup \{f^q(z_k)\}.$$

If $n = 1$, we also assume that $\phi(x) = x$ for all $x \in L$. Define $g = \phi \circ f \in H(X)$. Then

$$g(x) = f(x) \quad \text{whenever } f(x) \in (X - \text{Int}W) \cup \{f^q(z_k)\}$$

and b is a periodic point of g with period m . Moreover,

$$g^q(z_k) = f^q(z_k) \in \text{Int}W \quad \text{and} \quad g^j(z_k) = f^j(z_k) \notin \text{Int}W \quad \text{for } 0 \leq j \leq q-1.$$

Let Z and V be closed neighborhoods of b such that

$$g^q(z_k) \in \text{Int}V \subset V \subset \text{Int}Z \subset Z \subset \text{Int}W$$

and $\psi(Z)$ is a closed ball contained in $\{x \in \mathbf{R}^n; \|x\| < 1\}$. If $n = 1$, we also assume that

$$g(b), \dots, g^{m-1}(b) \notin Z.$$

Since $g(b), \dots, g^{m-1}(b) \notin Z$, there is a closed neighborhood V' of b such that $V' \subset \text{Int}Z$, $Z, g(V'), \dots, g^{m-1}(V')$ are pairwise disjoint, $g^m(V') \subset \text{Int}V$ and $\text{diam} g^i(V') < 1/r$ for $1 \leq i \leq m-1$. Now, let $\varphi \in H(X)$ be such that

$$\varphi(V) \subset V' \quad \text{and} \quad \varphi(x) = x \quad \text{for all } x \in (X - \text{Int}Z) \cup \{b\}.$$

Let $h = g \circ \varphi \in H(X)$. Then

$$\begin{aligned} h^q(z_k) &= g^q(z_k) \in \text{Int}V, \\ h^m(V) &= h^{m-1}(g(\varphi(V))) \subset g^m(V') \subset \text{Int}V \end{aligned}$$

and

$$\text{diam} h^i(V) \leq \text{diam} g^i(V') < \frac{1}{r} \quad \text{for } 1 \leq i \leq m-1.$$

Moreover, by choosing W small enough we can also guarantee that $\text{diam} V < 1/r$ (hence, $h \in \mathcal{O}_{r,k}$) and $d(h, f) < \epsilon$. This completes the proof.

REFERENCES

- [1] N. Aoki and K. Hiraide, *Topological Theory of Dynamical Systems: Recent Advances*, North-Holland, 1994. MR **95m**:58095
- [2] N. C. Bernardes Jr., *On the dynamics of homeomorphisms on the unit ball of $\mathbf{R}^n$* , Positivity **3** (1999), 149–172. MR **2000g**:37018
- [3] R. Diestel, *Graph Theory*, Springer-Verlag, 1997. CMP 97:12
- [4] J. R. Munkres, *Obstructions to the smoothing of piecewise-differentiable homeomorphisms*, Ann. of Math. **72** (1960), 521–554. MR **22**:12534
- [5] J. R. Munkres, *Elements of Algebraic Topology*, Addison-Wesley Publishing Company, 1984. MR **85m**:55001
- [6] K. Odani, *Generic homeomorphisms have the pseudo-orbit tracing property*, Proc. Amer. Math. Soc. **110** (1990), 281–284. MR **90m**:58111
- [7] S. Yu. Pilyugin and O. B. Plamenevskaya, *Shadowing is generic*, Topology Appl. **97** (1999), 253–266. MR **2000i**:37023
- [8] J. H. C. Whitehead, *Manifolds with transverse fields in Euclidean space*, Ann. of Math. **73** (1961), 154–212. MR **23a**:2225
- [9] K. Yano, *Generic homeomorphisms of S^1 have the pseudo orbit tracing property*, J. Fac. Sci. Univ. Tokyo Sect. IA Math. **34** (1987), 51–55. MR **88e**:58051

INSTITUTO DE MATEMÁTICA, UNIVERSIDADE FEDERAL FLUMINENSE, RUA MÁRIO SANTOS BRAGA
S/N, 24020-140, NITERÓI, RJ, BRASIL

E-mail address: `bernardes@mat.uff.br`