OPERATORS WHICH HAVE A CLOSED QUASI-NILPOTENT PART

PIETRO AIENA, MARIA LUISA COLASANTE, AND MANUEL GONZÁLEZ

(Communicated by Joseph A. Ball)

ABSTRACT. We find several conditions for the quasi-nilpotent part of a bounded operator acting on a Banach space to be closed. Most of these conditions are established for semi-Fredholm operators or, more generally, for operators which admit a generalized Kato decomposition. For these operators the property of having a closed quasi-nilpotent part is related to the so-called single valued extension property.

1. THE QUASI-NILPOTENT PART OF AN OPERATOR AND THE SVEP

The single valued extension property was first introduced by Dunford [5], [6] and has, successively, received a more systematic treatment in Dunford-Schwartz [7]. It also plays an important role in local spectral theory; see the monograph of Laursen and Neumann [14]. The following local version of this property has been studied in recent papers by Aiena and Monsalve [1], [2] and previously by Finch [8].

Definition 1.1. Let X be a complex Banach space and $T \in L(X)$. The operator T is said to have the *single valued extension property* at $\lambda_o \in \mathbb{C}$ (abbreviated SVEP at λ_o), if for every open disc $\mathbf{D}_{\lambda_o}$ centered at λ_o the only analytic function $f : \mathbf{D}_{\lambda_o} \rightarrow X$ which satisfies the equation $(\lambda I - T)f(\lambda) = 0$ for all $\lambda \in \mathbf{D}_{\lambda_o}$ is the function $f \equiv 0$.

An operator $T \in L(X)$ is said to have the SVEP if T has the SVEP at every point $\lambda \in \mathbb{C}$.

Let us consider the so-called *local resolvent set* $\rho_T(x)$ of T at the point $x \in X$, defined as the union of all open subsets $\mathcal{U}$ of $\mathbb{C}$ such that there exists an analytic function $f : \mathcal{U} \rightarrow X$ which satisfies $(\lambda I - T)f(\lambda) = x$ for all $\lambda \in \mathcal{U}$. The *local spectrum* $\sigma_T(x)$ of T at x is the set defined by $\sigma_T(x) := \mathbb{C} \setminus \rho_T(x)$. Obviously, $\sigma_T(x) \subseteq \sigma(T)$, where $\sigma(T)$ denotes the spectrum of T .

For every subset Ω of $\mathbb{C}$, the *analytic spectral subspace* of T associated with Ω is the set

$$X_T(\Omega) := \{x \in X : \sigma_T(x) \subseteq \Omega\}.$$

Received by the editors December 8, 2000 and, in revised form, April 20, 2001.

2000 *Mathematics Subject Classification.* Primary 47A10, 47A11; Secondary 47A53, 47A55.

Key words and phrases. Quasi-nilpotent part, single valued extension property, operators with a generalized Kato decomposition.

The research of the first two authors was supported by the International Cooperation Project between the University of Palermo (Italy) and Conicit-Venezuela.

The research of the third author was supported by DGICYT, Spain.

It is easy to see from the definition that $X_T(\Omega)$ is a T -hyperinvariant linear subspace of X [14].

The SVEP, as well as the SVEP at a point $\lambda_o \in \mathbb{C}$, may be characterized in a very simple way.

Theorem 1.2. *Let $T \in L(X)$, X a Banach space. Then:*

- (i) *T has the SVEP at λ_o if and only if $\ker(\lambda_o I - T) \cap X_T(\emptyset) = \{0\}$; see [1, Theorem 1.9].*
- (ii) *T has the SVEP if and only if $X_T(\emptyset) = \{0\}$, and this is the case if and only if $X_T(\emptyset)$ is closed; see [14, Proposition 1.2.16].*

Definition 1.3. Let X be a Banach space and $T \in L(X)$. The *analytical core* of T is the set $K(T)$ of all $x \in X$ such that there exists a sequence $(u_n)_{n \in \mathbb{N}} \subset X$ and $\delta > 0$ for which $x = u_o$, $Tu_{n+1} = u_n$ and $\|u_n\| \leq \delta^n \|x\|$, for every $n \in \mathbb{N} = \{0, 1, \dots\}$.

It is easy to check that $K(T)$ is a linear, generally not closed, subspace of X . Furthermore, $T(K(T)) = K(T)$ and if F is a closed subspace of X for which $T(F) = F$, then $F \subseteq K(T)$; see [19, Proposition 2]. Note that if T is quasi-nilpotent, then $K(T) = \{0\}$; see [16, Remarque 1.1].

Definition 1.4. Let $T \in L(X)$, X a Banach space. The *quasi-nilpotent part* of T is the set

$$H_o(T) := \{x \in X : \lim_{n \rightarrow \infty} \|T^n x\|^{\frac{1}{n}} = 0\}.$$

Also $H_o(T)$ is a linear subspace of X , generally not closed. Furthermore, $\ker(T^m) \subseteq H_o(T)$ for every $m \in \mathbb{N}$, and T is quasi-nilpotent if and only if $H_o(T) = X$; see [20, Theorem 1.5].

The next result may be found in Vrbová [20], or Mbekhta [16]; see also [14, Propositions 3.3.7 and 3.3.13].

Theorem 1.5. *For a bounded operator $T \in L(X)$, X a Banach space, we have:*

- (i) $K(\lambda_o I - T) = X_T(\mathbb{C} \setminus \{\lambda_o\})$.
- (ii) $H_o(\lambda_o I - T) \subseteq X_T(\{\lambda_o\})$ and the equality holds whenever T has the SVEP.

In the sequel by $M^\perp$ we shall denote the *annihilator* of the subset $M \subseteq X$, and by ${}^\perp N$ the *pre-annihilator* of the subset $N \subseteq X^*$.

Theorem 1.6. *For a bounded operator $T \in L(X)$, X a Banach space, the following implications hold:*

- (i) $H_o(\lambda_o I - T)$ closed $\Rightarrow H_o(\lambda_o I - T) \cap K(\lambda_o I - T) = \{0\} \Rightarrow T$ has the SVEP at λ_o .
- (ii) $X = H(\lambda_o I - T) + K(\lambda_o I - T) \Rightarrow T^*$ has the SVEP at λ_o .

Proof. Without loss of generality, we may consider $\lambda_o = 0$.

(i) Assume that $H_o(T)$ is closed. Let $\tilde{T}$ denote the restriction of T to the Banach space $H_o(T)$. Clearly, $H_o(T) = H_o(\tilde{T})$, thus $\tilde{T}$ is quasi-nilpotent. Therefore $K(\tilde{T}) = \{0\}$. It is easy to see that $H_o(T) \cap K(T) = K(\tilde{T})$. This shows the first implication.

The second implication of (i) is an immediate consequence of Theorem 1.5. Indeed, we have

$$\ker(\lambda_o I - T) \cap X_T(\emptyset) \subseteq H_o(\lambda_o I - T) \cap K(\lambda_o I - T),$$

so, if the last intersection is $\{0\}$, then T has the SVEP at λ_o , by Theorem 1.2.

(ii) From [15, Proposition 1.8] we know that $H_o(T) \subseteq^\perp K(T^*)$ and therefore $K(T^*) \subseteq H_o(T)^\perp$. We also have $H_o(T^*) \subseteq K(T)^\perp$. Indeed, let $\varphi \in H_o(T^*)$ and consider an arbitrary element $x \in K(T)$. According to the definition of $K(T)$, there is a sequence $(u_n)_{n \in \mathbb{N}} \subset X$, and a $\delta > 0$, such that $u_o = x$, $Tu_{n+1} = u_n$ and $\|u_n\| \leq \delta^n \|x\|$ for every $n \in \mathbb{N}$. Clearly, $T^n u_n = x$ for every $n \in \mathbb{N}$. Consequently,

$$|\varphi(x)| = |\varphi(T^n u_n)| = |(T^{*n} \varphi)(u_n)| \leq \|u_n\| \|T^{*n} \varphi\| \leq \delta^n \|T^{*n} \varphi\|,$$

and hence $|\varphi(x)|^{\frac{1}{n}} \leq \delta \|T^{*n} \varphi\|^{\frac{1}{n}}$ for every $n \in \mathbb{N}$. The last term converges to 0 as $n \rightarrow \infty$, since $\varphi \in H_o(T^*)$, and from this it follows that $\varphi(x) = 0$, i.e. $\varphi \in K(T)^\perp$. Finally, if $X = H_o(T) + K(T)$, then $\{0\} = H_o(T)^\perp \cap K(T)^\perp \supseteq H_o(T^*) \cap K(T^*)$. Thus, by part (i), T^* has the SVEP at 0. $\square$

The two implications of part (i) of Theorem 1.6 were observed in [16, Proposition 1.10]. Later we shall prove that in the case of semi-Fredholm operators, or more generally in the case that $\lambda_o I - T$ admits a generalized Kato decomposition, the implications of Theorem 1.6 are actually equivalences.

Theorem 1.6 suggests in a very natural way the following definition:

Definition 1.7. A bounded operator $T \in L(X)$, X a Banach space, is said to have *property (Q)* if $H_o(\lambda I - T)$ is closed for every $\lambda \in \mathbb{C}$.

Recall that a bounded operator $T \in L(X)$, X a Banach space, is said to have *Dunford's property (C)*, shortly *property (C)*, if the analytic subspace $X_T(\Omega)$ is closed for every closed subset $\Omega \subseteq \mathbb{C}$. From part (ii) of Theorem 1.2 it follows that if $T \in L(X)$ has *property (C)*, then T has the SVEP.

An obvious consequence of part (ii) of Theorem 1.5 is that if T has *property (C)*, then $H_o(\lambda I - T) = X_T(\{\lambda\})$ is closed for every $\lambda \in \mathbb{C}$, so that the following implications hold:

(1) T has *property (C)* $\Rightarrow T$ has *property (Q)* $\Rightarrow T$ has the SVEP.

Note that neither of the implications in (1) may be reversed in general. A first counterexample, which shows that the first implication is not reversed in general, may be found among the convolution operators of group algebras.

Recall that a Banach algebra A is said to be *semi-simple* if the radical $\text{rad } A = \{0\}$ [3]; A is said to be *semi-prime* if there is no non-zero two-sided ideal J for which $J^2 = \{0\}$. Note that A is semi-prime if and only if, for every $x \in A$, the identity $xAx = \{0\}$ implies that $x = 0$, and that a commutative algebra is semi-prime if and only if it contains no non-zero nilpotent elements. Clearly, any semi-simple Banach algebra is semi-prime. A map $T : A \rightarrow A$, A a Banach algebra, is said to be a *multiplier* if $(Tx)y = x(Ty)$ holds for all $x, y \in A$. Note that if T is a multiplier of a semi-prime commutative Banach algebra A , then $(Tx)y = x(Ty) = T(xy)$ for every $x, y \in A$; see the proof of [11, Theorem 1.1.1]. A very important example of a multiplier is given in the case that A is the semi-simple Banach algebra $L_1(G)$, the group algebra of a locally compact abelian group G with convolution as multiplication. Indeed, in this case to any complex Borel measure μ on G there corresponds a multiplier T_μ defined by

$$T_\mu(f) := \mu \star f \quad \text{for all } f \in L_1(G),$$

where

$$(\mu \star f)(t) := \int_G f(t-s) d\mu(s).$$

The classical Helson-Wendel Theorem shows that each multiplier is a convolution operator and the multiplier algebra of $A := L_1(G)$ may be identified with the measure algebra $M(G)$; see [11, Chapter 0].

Theorem 1.8. *Let T be a multiplier of a semi-simple Banach algebra A . Then*

$$H_o(T) = \ker T.$$

In particular, T has property (Q).

Proof. We know that $\ker T \subseteq H_o(T)$, so it remains to prove the inverse inclusion.

Suppose that $x \in H_o(T)$. By an easy inductive argument we have

$$(Ty)^n = (T^n y)y^{n-1} \text{ for every } y \in A \text{ and } n \in \mathbb{N}.$$

From this it follows that

$$\begin{aligned} \|(aTx)^n\| &= \|(T^na)x\| = \|T^n(ax)(ax)^{n-1}\| \\ &\leq \|a\| \|T^n x\| \|(ax)^{n-1}\| \end{aligned}$$

for every $a \in A$, so the spectral radius of the element aTx satisfies

$$r(aTx) = \lim_{n \rightarrow \infty} \|(aTx)^n\|^{\frac{1}{n}} = 0$$

for every $a \in A$. This implies that $Tx \in \text{rad } A$; see [3, §25, Proposition 1]. Since A is semi-simple then $Tx = 0$ and therefore $H_o(T) \subseteq \ker T$.

The last assertion is clear, because $\lambda I - T$ is a multiplier of A for every $\lambda \in \mathbb{C}$. $\square$

Clearly, if T is a quasi-nilpotent multiplier, then $\ker T = H_o(T) = A$, so $T = 0$ ([13]).

The following example shows that the assumption that A is semi-simple in Theorem 1.8 cannot be replaced by the weaker assumption that A is semi-prime.

Example 1.9. Let $\omega := (\omega_n)_{n \in \mathbb{N}}$ be a sequence with the property that $0 < \omega_{m+n} \leq \omega_m \omega_n$ for all $m, n \in \mathbb{N}$. Let $\ell^1(\omega)$ denote the space of all complex sequences $x := (x_n)_{n \in \mathbb{N}}$ for which $\|x\|_\omega := \sum_{n=0}^\infty \omega_n |x_n| < \infty$. The space $\ell^1(\omega)$ equipped with convolution

$$(x \star y)_n := \sum_{j=0}^n x_{n-j} y_j \text{ for all } n \in \mathbb{N}$$

is a commutative unital Banach algebra. Let A_ω denote the maximal ideal of $\ell^1(\omega)$ given by

$$A_\omega := \{(x_n)_{n \in \mathbb{N}} \in \ell^1(\omega) : x_0 = 0\}.$$

The Banach algebra A_ω is an integral domain and hence semi-prime. Moreover, if the weight sequence ω satisfies the condition $\rho_\omega := \lim_{n \rightarrow \infty} \omega_n^{\frac{1}{n}} = 0$, then A_ω is a radical algebra ([14, Example 4.1.9]), i.e. coincides with its radical. Now, fix $0 \neq a \in A_\omega$ and let $T_a(x) := a \star x$, $x \in A_\omega$, denote the multiplication operator by the element a . From the estimate

$$\|T^n x\|^{\frac{1}{n}} = \|a^n \star x\|^{\frac{1}{n}} \leq \|a^n\|^{\frac{1}{n}} \|x\|^{\frac{1}{n}}$$

we see that T_a is quasi-nilpotent, thus $H_o(T_a) = A_\omega$. On the other hand, A_ω is an integral domain so that $\ker T_a = \{0\}$.

Theorem 1.8 suggests the way of obtaining examples of operators which have property (Q) but not property (C). Indeed, there are convolution operators T_μ , $\mu \in M(G)$, on the group algebra $L_1(G)$ which do not enjoy property (C); see [14, Chapter 4].

The next example, which is obtained by a slight modification of Example 3.9 of [4], shows that also the second implication of (1) may be not reversed in general.

Example 1.10. Let $X := \ell_2 \oplus \ell_2 \cdots$ and define

$$T_n e_i := \begin{cases} e_{i+1} & \text{if } i = 1, \dots, n, \\ \frac{e_{i+1}}{i-n} & \text{if } i > n. \end{cases}$$

Clearly,

$$\|T_n^{n+k}\| = \frac{1}{k!} \quad \text{and} \quad \left(\frac{1}{k!}\right)^{\frac{1}{n+k}} \rightarrow 0 \text{ as } k \rightarrow \infty.$$

From this it follows that $\sigma(T_n) = \{0\}$. Moreover, T_n is injective and the point spectrum $\sigma_p(T_n)$ is empty. Now, let $T := T_1 \oplus \cdots \oplus T_n \oplus \cdots$. From $\|T_n\| = 1$, for every $n \in \mathbf{N}$, we obtain $\|T\| = 1$. From $\sigma_p(T_n) = \emptyset$ it also follows that $\sigma_p(T) = \emptyset$. Take $x = (x_n) \in X$ with $x_n := \frac{e_n}{n}$. We have

$$\|x\| = \left(\sum_{n=1}^{\infty} \frac{1}{n^2}\right)^{\frac{1}{2}} < \infty,$$

thus $x \in X$. Moreover,

$$\|T^n x\|^{\frac{1}{n}} \geq \|T_n^n \frac{e_1}{n}\|^{\frac{1}{n}} = \left(\frac{1}{n}\right)^{\frac{1}{n}}$$

and the last term does not converge to 0. Clearly, $x \notin H_o(T)$. Finally, $\ell_2 \oplus \ell_2 \cdots \oplus \ell_2 \oplus \{0\} \cdots \subset H_o(T)$, where the non-zero terms are n . Since $H_o(T)$ contains all sequences with only finitely many non-zero terms, it follows that $H_o(T)$ is dense in X . Since $H_o(T) \neq X$ then $H_o(T)$ is not closed, thus T has not property (Q). Note that the operator T has the SVEP, since $\sigma_p(T)$ is empty.

2. THE CASE OF SEMI-FREDHOLM OPERATORS

For every linear operator T on a vector space X , let us consider the increasing sequence of kernels $\ker T^n$ and the decreasing sequence of ranges $T^n(X)$.

Definition 2.1. Let T be a linear operator on a vector space X . The *generalized kernel* of T is defined by

$$\mathcal{N}(T) := \bigcup_{n \in \mathbf{N}} \ker T^n.$$

The *hyperrange* of T is defined by

$$T^\infty(X) := \bigcap_{n \in \mathbf{N}} T^n(X).$$

Clearly, $T^\infty(X)$ is a T -invariant subspace and it is easily seen that, if X is a Banach space, $K(T) \subseteq T^\infty(X)$. Moreover, for every $n \in \mathbf{N}$ we have

$$\ker (\lambda_o I - T)^n \subseteq \mathcal{N}(\lambda_o I - T) \subseteq H_o(\lambda_o I - T).$$

Recall that T is said to have *finite ascent* if $\mathcal{N}(T) = \ker T^k$ for some positive integer k . Clearly, in such a case there is a smallest positive integer $p = p(T)$ such that $\ker T^p = \ker T^{p+1}$. The positive integer p is called the *ascent* of T . Analogously, T is said to have *finite descent* if $T^\infty(X) = T^k(X)$ for some k . The smallest integer $q = q(T)$ such that $T^{q+1}(X) = T^q(X)$ is called the *descent* of T . It is possible to prove that if $p(T)$ and $q(T)$ are both finite, then $p(T) = q(T)$; see [9, Proposition 38.3].

Theorem 2.2 ([1]). *For a bounded operator T on a Banach space X the following implications hold:*

- i) $p(\lambda_o I - T) < \infty \Rightarrow T$ has SVEP at λ_o .
- ii) $q(\lambda_o I - T) < \infty \Rightarrow T^*$ has SVEP at λ_o .

Hence each one of the two conditions $p(\lambda_o I - T) < \infty$ or $H_o(\lambda_o I - T)$ closed implies the SVEP at λ_o . In general these two conditions are not related. The operator T of Example 1.10 has ascent $p(T) = 0$ and quasi-nilpotent part $H_o(T)$ not closed. In the following example we find an operator T which has a closed quasi-nilpotent part but ascent $p(T) = \infty$.

Example 2.3. Let $T : \ell_2 \rightarrow \ell_2$ be defined by

$$Tx := \left(\frac{x_2}{2}, \dots, \frac{x_n}{n}, \dots\right), \text{ where } x = (x_1, \dots, x_n, \dots).$$

It is easily seen that $\|T^k\| = \frac{1}{(k+1)!}$ and from this it follows that T is quasi-nilpotent and therefore $H_o(T) = \ell_2$. Obviously, $p(T) = \infty$.

Definition 2.4. An operator $T \in L(X)$, X a Banach space, is said to be *semi-regular* if $T(X)$ is closed and $\ker T \subseteq T^\infty(X)$.

An operator $T \in L(X)$ is said to admit a *generalized Kato decomposition*, abbreviated GKD, if there exists a pair of T -invariant closed subspaces (M, N) such that $X = M \oplus N$, the restriction $T|_M$ is semi-regular and $T|_N$ is quasi-nilpotent.

Remark 2.5. In the sequel we list some examples of operators which admit a GKD.

(i) Every semi-regular operator has the GKD $M = X$ and $N = \{0\}$. Note that if T is semi-regular, then $\overline{H_o(T)} = \overline{\mathcal{N}(T)}$; see [15, Proposition 2.10].

(ii) Every quasi-nilpotent operator has the GKD $M = \{0\}$ and $N = X$.

(iii) An important case is obtained if we assume in the definition above that $T|_N$ is nilpotent. In this case T is said to be of *Kato type*; see [15]. Obviously, any semi-regular operator is of Kato type. Note that if T is of Kato type, then $T^\infty(X) = K(T)$ and $K(T)$ is closed; see [1, Lemma 2.4] or [2, Theorem 2.3 and Theorem 2.4].

(iv) Let

$$\Phi_+(X) := \{T \in L(X) : \dim \ker T < \infty, T(X) \text{ closed}\}$$

denote the class of all *upper semi-Fredholm* operators, and let

$$\Phi_-(X) := \{T \in L(X) : \operatorname{codim} T(X) < \infty\}$$

denote the class of all *lower semi-Fredholm* operators. The class of all *semi-Fredholm operators* is defined as $\Phi_\pm(X) := \Phi_+(X) \cup \Phi_-(X)$ and the class of all *Fredholm operators* is defined as $\Phi(X) := \Phi_+(X) \cap \Phi_-(X)$. A well-known result of Kato [10, Theorem 4] establishes that every $T \in \Phi_\pm(X)$ is of Kato type. More precisely, T admits a GKD (M, N) with $T|_N$ nilpotent and $\dim N < \infty$.

Recall that for $T \in \Phi_{\pm}(X)$ the *index* of T is defined by $\text{ind } T := \dim \ker T - \text{codim } T(X)$. The index is an integer or $\pm\infty$.

Theorem 2.6. *Let $T \in L(X)$, X a Banach space, and assume that $\lambda_o I - T$ has a GKD (M, N) . Then the following properties are equivalent:*

- (i) T has the SVEP at λ_o .
- (ii) $H_o(\lambda_o I - T) \cap K(\lambda_o I - T) = \{0\}$.
- (iii) $H_o(\lambda_o I - T)$ is closed.
- (iv) $H_o(\lambda_o I - T) = N$.

In particular, if $\lambda_o I - T$ is semi-regular the conditions (i)-(iv) are equivalent to the following one.

- (v) $H_o(\lambda_o I - T) = \{0\}$.

Proof. Also here we only consider the case $\lambda_o = 0$. Clearly, (iv) $\Rightarrow$ (iii) and from Theorem 1.6 we know that (iii) $\Rightarrow$ (ii) $\Rightarrow$ (i).

(i) $\Rightarrow$ (iv). First note that if T admits a GKD (M, N) , then $H_o(T) = H_o(T|_M) + H_o(T|_N)$. The inclusion $H_o(T) \supseteq H_o(T|_M) + H_o(T|_N)$ is obvious. In order to show the opposite inclusion, consider an arbitrary element $x \in H_o(T)$ and let $x = u + v$, with $u \in M$ and $v \in N$. Since $T|_N$ is quasi-nilpotent then $N = H_o(T|_N) \subseteq H_o(T)$. Consequently, $u = x - v \in H_o(T) \cap M = H_o(T|_M)$ and therefore $H_o(T) \subseteq H_o(T|_M) + H_o(T|_N)$. Hence $H_o(T) = H_o(T|_M) + H_o(T|_N) = H_o(T|_M) + N$. Now, suppose that T has the SVEP 0. Clearly, the SVEP at a point is inherited by the restrictions to closed invariant subspaces, so $T|_M$ has the SVEP at 0 and from the semi-regularity of $T|_M$ it follows that $T|_M$ is injective; see [1, Theorem 2.14]. From this we obtain

$$H_o(T|_M) = \overline{\bigcup_{n=1}^{\infty} \ker (T|_M)^n} = \{0\}$$

and therefore $H_o(T) = N$.

The final assertion is clear. □

Corollary 2.7. *Let $T \in L(X)$, X a Banach space, and assume that $\lambda_o I - T$ is of Kato type. Then the conditions (i)-(iv) of Theorem 2.6 are equivalent to the following one:*

- (v') $p(\lambda_o I - T) < \infty$.

In this case, if $p := p(\lambda_o I - T)$, then

$$H_o(\lambda_o I - T) = \mathcal{N}(\lambda_o I - T) = \ker (\lambda_o I - T)^p.$$

Proof. Assume $\lambda_o = 0$. We know that the inclusions $H_o(T) \supseteq \mathcal{N}(T) \supseteq \ker T^n$ hold for every $T \in L(X)$ and for every $n \in \mathbb{N}$. Let (M, N) be a GKD for T such that $(T|_N)^k = 0$ for some $k \in \mathbb{N}$. Then $H_o(T) = N \subseteq \ker T^k$ and hence $H_o(T) = \mathcal{N}(T) = \ker T^k$. From this it follows that $p := p(T) \leq k$ and therefore $\ker T^k = \ker T^p$. □

The next result shows that, under the assumption that $\lambda_o I - T$ is semi-Fredholm, another equivalent condition can be added to those given in Corollary 2.7.

Theorem 2.8. *Suppose that $\lambda_o I - T \in L(X)$ is a semi-Fredholm operator. Then the following statements are equivalent:*

- (i) $H_o(\lambda_o I - T)$ is closed.
- (ii) $H_o(\lambda_o I - T)$ is finite dimensional.

Proof. We have only to prove the implication (i) $\Rightarrow$ (ii). This immediately follows from the mentioned Kato decomposition of a semi-Fredholm operator: if (M, N) is a GKD for $\lambda_o I - T$ such that $\lambda_o I - T|_N$ is nilpotent and $\dim N < \infty$, then $H_o(\lambda_o I - T) = N$, by Theorem 2.6. $\square$

The preceding result was obtained in [12, Theorem 2] under the assumption that $\lambda_o I - T$ is a Fredholm operator.

Theorem 2.9. *Suppose that $\lambda_o I - T \in L(X)$ is of Kato type. Then the following statements are equivalent:*

- (i) T^* has the SVEP at λ_o .
- (ii) $q := q(\lambda_o I - T) < \infty$.
- (iii) $X = H_o(\lambda_o I - T) + K(\lambda_o I - T)$.

Moreover, if any of the equivalent conditions (i)-(iii) holds, then

$$(\lambda_o I - T)^\infty(X) = K(\lambda_o I - T) = (\lambda_o I - T)^q(X).$$

Proof. Assume that $\lambda_o = 0$. The equivalence (i) $\Leftrightarrow$ (ii) has been proved in [2, Theorem 2.6]. The implication (iii) $\Rightarrow$ (i) has been proved in Theorem 1.6.

(ii) $\Rightarrow$ (iii). Assume that $q := q(T) < \infty$. Since T is of Kato type then $K(T) = T^\infty(X) = T^q(X)$. Moreover, $X = \ker T^q + T^n(X)$ for every $n \in \mathbb{N}$ (see [9, Proposition 38.2]), and therefore $X = H_o(T) + T^\infty(X)$. $\square$

Theorem 2.10. *Suppose that $\lambda_o I - T \in L(X)$ is a semi-Fredholm operator. Then the following statements are equivalent:*

- (i) T^* has the SVEP at λ_o .
- (ii) $K(\lambda_o I - T)$ is finite codimensional.

Proof. Also here we assume that $\lambda_o = 0$.

(i) $\Rightarrow$ (ii). From Fredholm theory we know that T^* is also a semi-Fredholm operator and $\text{ind } T^* = -\text{ind } T$. Now, if T^* has the SVEP at 0, then $\text{ind } T^* \leq 0$ (see [1, Corollary 2.7]) and therefore $\text{ind } T \geq 0$. From this it follows that T is a lower semi-Fredholm and consequently also T^q is lower semi-Fredholm, i.e. $T^q(X) = T^\infty(X) = K(T)$ is finite codimensional.

(ii) $\Rightarrow$ (i). Since $K(T) = T^\infty(X)$, condition (ii) means that $T^\infty(X)$ is of finite codimension. But from this it is immediate that $q(T) < \infty$, so that T^* has the SVEP at 0, by Theorem 2.9. $\square$

It should be noted that if T is semi-Fredholm, then $T^\infty(X)$ coincides with the so-called algebraic core of T , i.e. the greatest subspace M of X for which $T(M) = M$; see for instance [1, Theorem 2.3].

Corollary 2.11. *Assume that $\lambda_o I - T \in L(X)$ is a semi-Fredholm operator. Then the following statements are equivalent:*

- (i) T and T^* have the SVEP at λ_o .
- (ii) $X = H_o(\lambda_o I - T) \oplus K(\lambda_o I - T)$.
- (iii) $H_o(\lambda_o I - T)$ is closed and $K(\lambda_o I - T)$ is finite-codimensional.
- (iv) λ_o is a pole of $(\lambda I - T)^{-1}$, or equivalently $p(\lambda_o I - T) = q(\lambda_o I - T) < \infty$.
- (v) The spectrum does not cluster at λ_o .

In particular, if any of the equivalent conditions (i)-(v) holds and $p := p(\lambda_o I - T) = q(\lambda_o I - T)$, then

$$H_o(\lambda_o I - T) = \mathcal{N}(\lambda_o I - T) = \ker(\lambda_o I - T)^p$$

and

$$K(\lambda_o I - T) = (\lambda_o I - T)^\infty(X) = (\lambda_o I - T)^p(X).$$

Proof. The equivalences of (i), (ii), (iii), and (iv) are obtained by combining all the results established in this section. The implication (iv) $\Rightarrow$ (v) is obvious. The implication (v) $\Rightarrow$ (i) is an immediate consequence of the fact that both T and T^* have the SVEP at every point of the resolvent, as well as at every isolated point of the spectrum. $\square$

Note that rather similar results to those of Corollary 2.11 have been established by Mbekhta [15, Théorème 1.6] and Schmoege [19], in the case that λ_o is an isolated point of the spectrum.

Remark 2.12. Recall that for every $T \in L(X)$ the *semi-Fredholm region* is defined to be

$$\Sigma(T) := \{\lambda \in \mathbb{C} : \lambda I - T \text{ is semi-Fredholm}\}.$$

It is well-known that $\Sigma(T)$ is an open set and hence it may be decomposed in connected disjoint open nonempty components. Suppose that T has SVEP at some $\lambda_o \in \Omega$, Ω a component of $\Sigma(T)$. Then, by Theorem 2.6 and Corollary 2.7

$$\begin{aligned} \{0\} &= H_o(\lambda_o I - T) \cap K(\lambda_o I - T) = \mathcal{N}(\lambda_o I - T) \cap (\lambda_o I - T)^\infty(X) \\ &= \overline{\mathcal{N}(\lambda_o I - T)} \cap (\lambda_o I - T)^\infty(X), \end{aligned}$$

and from the constancy of the map $\lambda \in \Omega \rightarrow \overline{\mathcal{N}(\lambda I - T)} \cap (\lambda I - T)^\infty(X)$ (see [18, Theorem 4.2]) we conclude that $\mathcal{N}(\lambda I - T) \cap (\lambda I - T)^\infty(X) = \{0\}$ for every point $\lambda \in \Omega$. From [2, Theorem 1.10] it follows that T has the SVEP at every point $\lambda \in \Omega$. Moreover, the set

$$\Gamma := \{\lambda \in \Omega : \text{jump } (\lambda I - T) \neq 0\}$$

is countable ([18]) and is equal to the set of all $\lambda \in \Omega$ such that $\lambda I - T$ is not semi-regular; see [21, Proposition 2.2]. From Theorem 2.6 then $H_o(\lambda I - T) = \{0\}$ for every $\lambda \in \Omega \setminus \Gamma$, while the remaining points $\lambda \in \Gamma$ are eigenvalues with ascent $p := p(\lambda I - T) < \infty$, $H_o(\lambda I - T) = \ker (\lambda I - T)^p$ and $0 < \dim H_o(\lambda I - T) < \infty$, by Corollary 2.7 and Theorem 2.8. In particular this situation occurs for every component of the semi-Fredholm region of an operator which has the SVEP.

REFERENCES

- [1] P. Aiena, O. Monsalve *Operators which do not have the single valued extension property*. J. Math. Anal. Appl. **250** (2000), 435-448. MR **2001g**:47005
- [2] P. Aiena, O. Monsalve *The single valued extension property and the generalized Kato decomposition property*. Acta Sci. Math.(Szeged) **67** (2001), 461-477.
- [3] F. F. Bonsall, J. Duncan *Complete normed algebras*. Springer-Verlag, Berlin, 1973. MR **54**:11013
- [4] I. Colojoară, C. Foiaş *Theory of generalized spectral operators*. Gordon and Breach, New York, 1968. MR **52**:15085
- [5] N. Dunford *Spectral theory II. Resolution of the identity*. Pacific J. Math. **2** (1952), 559-614. MR **14**:479a
- [6] N. Dunford *Spectral operators*. Pacific J. Math. **4** (1954), 321-354. MR **16**:142d
- [7] N. Dunford, J. T. Schwartz *Linear operators, Part III*. (1971), Wiley, New York. MR **54**:1009
- [8] J. K. Finch *The single valued extension property on a Banach space*. Pacific. J. Math. **58** (1975), 61-69. MR **51**:11181
- [9] H. Heuser *Functional Analysis*. (1982), Wiley, New York. MR **83m**:46001

- [10] T. Kato *Perturbation theory for nullity, deficiency and other quantities of linear operators*. J. Anal. Math. **6** (1958), 261-322. MR **21**:6541
- [11] R. Larsen *An introduction to the theory of multipliers*. Springer-Verlag, New York, 1979. MR **55**:8695
- [12] K. B. Laursen, *Essential spectra through local spectral theory*. Proc. Amer. Math. Soc. **125** (1997), 1425-1434. MR **97g**:46066
- [13] K. B. Laursen, M. Mbekhta *Closed range multipliers and generalized inverses*. Studia Math. **107** (1993), 127-135. MR **94i**:47052
- [14] K. B. Laursen, M. M. Neumann *An introduction to local spectral theory*. Clarendon Press, Oxford, 2000. MR **2001k**:47002
- [15] M. Mbekhta *Généralisation de la décomposition de Kato aux opérateurs paranormaux et spectraux*. Glasgow Math. J. **29** (1987), 159-175. MR **88i**:47010
- [16] M. Mbekhta *Sur la théorie spectrale locale et limite des nilpotents*. Proc. Amer. Math. Soc. **110** (1990), 621-631. MR **91b**:47004
- [17] M. Mbekhta, A. Ouahab *Opérateur s -régulier dans un espace de Banach et théorie spectrale*. Acta Sci. Math. (Szeged) **59** (1994), 525-543. MR **96a**:47018
- [18] M. Ó Searcóid, T. T. West *Continuity of the generalized kernel and range for semi-Fredholm operators*. Math. Proc. Camb. Phil. Soc. **105** (1989), 513-522. MR **90d**:47017
- [19] C. Schmoege *On isolated points of the spectrum of a bounded operator*. Proc. Amer. Math. Soc. **117** (1993), 715-719. MR **93d**:47007
- [20] P. Vrbová *On local spectral properties of operators in Banach spaces*. Czechoslovak Math. J. **23** (98) (1973), 483-92. MR **48**:898
- [21] T. T. West *A Riesz-Schauder theorem for semi-Fredholm operators*. Proc. Roy. Irish. Acad. **87 A** (1987), 137-146. MR **89i**:47020

DIPARTIMENTO DI MATEMATICA ED APPLICAZIONI, FACOLTÀ DI INGEGNERIA, UNIVERSITÀ DI PALERMO, VIALE DELLE SCIENZE, I-90128 PALERMO, ITALY

E-mail address: `paiena@mbbox.unipa.it`

DEPARTAMENTO DE MATEMÁTICAS, FACULTAD DE CIENCIAS, UNIVERSIDAD DE LOS ANDES, MERIDA, VENEZUELA

E-mail address: `marucola@ciens.ula.ve`

DEPARTAMENTO DE MATEMÁTICAS, FACULTAD DE CIENCIAS, UNIVERSIDAD DE CANTABRIA, SANTANDER, SPAIN

E-mail address: `gonzalem@ccaix3.unican.es`