TOPOLOGICALLY TRANSVERSAL REVERSIBLE HOMOCLINIC SETS

MICHAL FEČKAN

(Communicated by Carmen Chicone)

ABSTRACT. An R -reversible diffeomorphism on $\mathbb{R}^{2N}$ is studied possessing a hyperbolic fixed point. If the stable manifold of the hyperbolic fixed point and the fixed point set $\text{Fix } R$ of R have a nontrivial local topological crossing, then an infinite number of R -symmetric periodic orbits of the diffeomorphism is shown. A perturbed problem is also studied by showing the relationship between a corresponding Melnikov function and the nontriviality of a local topological crossing of the set $\text{Fix } R$ and the stable manifold for the perturbed diffeomorphism.

1. INTRODUCTION

Let $R : \mathbb{R}^{2N} \rightarrow \mathbb{R}^{2N}$ be a linear involution, i.e. $R^2 = I$, such that $\dim \text{Fix } R = N$, where $\text{Fix } R = \{x \in \mathbb{R}^{2N} \mid Rx = x\}$. Consider a C^1 -smooth diffeomorphism $f : \mathbb{R}^{2N} \rightarrow \mathbb{R}^{2N}$ which is R -reversible, i.e. $Rf(x) = f^{-1}(Rx)$, $\forall x \in \mathbb{R}^{2N}$, and possessing an R -symmetric hyperbolic fixed point $p \in \text{Fix } R$. Any subset of $\mathbb{R}^{2N}$ invariant under the action of R is called R -symmetric. Reversible diffeomorphisms naturally come from mechanics [5] as the time flow mappings of second order gradient differential equations. Let W_p^s, W_p^u be the global stable and unstable manifolds of p , respectively. Let $\widetilde{W}_p^s$ be an open subset of W_p^s which is a submanifold of $\mathbb{R}^{2N}$, i.e. the immersed and induced topologies on $\widetilde{W}_p^s$ coincide, and such that $\widetilde{W}_p^s \setminus \{p\} \cap \text{Fix } R \neq \emptyset$, i.e. there is an R -symmetric point q homoclinic to p [10]. Since $RW_p^s = W_p^u$, we put $\widetilde{W}_p^u = R\widetilde{W}_p^s$. We also suppose the existence of a compact component $K \ni q$ of the set $\widetilde{W}_p^s \cap \text{Fix } R$, that is a compact subset $K \subset \widetilde{W}_p^s \setminus \{p\} \cap \text{Fix } R$ such that $q \in K$ and there exists an open bounded subset $U \subset \widetilde{U} \subset \overline{\mathbb{R}^{2N} \setminus \{p\}}$ satisfying $U \cap \widetilde{W}_p^s \cap \text{Fix } R = K$. By shrinking U , we can assume that $\widetilde{W}_p^s \cap U = \widetilde{W}_p^s \cap \widetilde{U}$. We note that $\widetilde{W}_p^s \cap U$ is an orientable submanifold of $\mathbb{R}^{2N}$. Then we can define the local intersection number $\#(\widetilde{W}_p^s \cap U, \text{Fix } R \cap U)$ of the stable manifold W_p^s and the plain $\text{Fix } R$ in $U \subset \mathbb{R}^{2N}$ [7]. The main purpose of this note is to prove the following result.

Theorem 1.1. *If $\#(\widetilde{W}_p^s \cap U, \text{Fix } R \cap U) \neq 0$, then there is an $\omega_0 \in \mathbb{N}$ such that for any $\mathbb{N} \ni \omega \geq \omega_0$ the diffeomorphism f possesses a 2ω -periodic orbit $\{x_n^\omega\}_{n \in \mathbb{Z}}$ such*

Received by the editors April 25, 2001 and, in revised form, June 29, 2001.

2000 *Mathematics Subject Classification.* Primary 37C25, 37C29, 57R50.

The author was partially supported by Grant GA-MS 1/6179/00.

that $Rx_n^\omega = x_{-n}^\omega$, $n \in \mathbb{Z}$. Moreover, $x_0^\omega \in \text{Fix } R$ is near the set K , while $x_\omega^\omega \in \text{Fix } R$ is near the point p .

When q is a transversal intersection of W_p^s and $\text{Fix } R$, then Theorem 1.1 is proved in [3], [4], [5], [9] and [10]. Then clearly $\#(\widetilde{W}_p^s \cap U, \text{Fix } R \cap U) \neq 0$ for a small open neighbourhood U of q . Furthermore, we study the case where W_p^s and $\text{Fix } R$ intersect on a compact manifold. Then we consider a C^2 -smooth R -reversible perturbation of f . Associated to such a perturbation there is a Melnikov function. We show that if the Brouwer degree [6] of this Melnikov function is not zero, and the perturbation is small, then the perturbed stable manifold $W_{p,per}^s$ and the plain $\text{Fix } R$ satisfy $\#(\widetilde{W}_{p,per}^s \cap U, \text{Fix } R \cap U) \neq 0$ with the corresponding infinitely many R -symmetric periodic orbits of the perturbed diffeomorphism. Finally, we note that any accumulation point of the set $\{x_0^\omega\}_{\omega \geq \omega_0} \subset \text{Fix } R$ from Theorem 1.1 is a starting point of an R -symmetric homoclinic orbit of f to p .

If p is a hyperbolic fixed point of f but not R -symmetric, then Rp is also a hyperbolic fixed point of f . If $q \in W_p^s \cap \text{Fix } R$, then $q \in W_p^s \cap W_{Rp}^u$ [5], hence q lies on an R -symmetric heteroclinic orbit connecting p and Rp . Consequently, as for Theorem 1.1, we can prove the following result.

Theorem 1.2. *Suppose f has a non- R -symmetric hyperbolic fixed point p . If W_p^s and W_p^u meet $\text{Fix } R$ locally topologically transversally, then f has an infinite number of R -symmetric periodic orbits with periods tending to infinity.*

We note that for the case $N = 1$ it is elementary to show that if p is a hyperbolic fixed point of f and W_p^s (or W_p^u) meets $\text{Fix } R$, then a local intersection number of W_p^s (or W_p^u) with $\text{Fix } R$ is nonzero. Then Theorems 1.1 and 1.2 can be applied. Indeed, let $q \in W_p^s \cap \text{Fix } R$ be the first intersection starting on W_p^s from p . Then the points $f^{-1}(q), f(q) \in W_p^s$ lie on the opposite half-planes separated by $\text{Fix } R$. Hence an open bounded connected part $\widetilde{W}_p^s$ of W_p^s such that $f^{-1}(q), f(q) \in \widetilde{W}_p^s$ topologically nontrivially crosses $\text{Fix } R$. Similarly for W_p^u .

The paper is finished with an example of a perturbed second order differential equation in $\mathbb{R}^2$ with a topologically transversal, but non- C^1 -transversal, intersection of the stable manifold and $\text{Fix } R$.

2. PRELIMINARY RESULTS

Let $(\cdot, \cdot)$ be an inner product on $\mathbb{R}^{2N}$. By following [10], we set

$$\langle x, y \rangle = \frac{1}{2}((x, y) + (Rx, Ry)).$$

Then $\langle Rx, Ry \rangle = \langle x, y \rangle$, $x, y \in \mathbb{R}^{2N}$, and so $\|R\| = \|R^{-1}\| = 1$. Since $RK = K$, we can assume that $RU = U$.

For any $\xi \in \widetilde{W}_p^s \cap \bar{U}$ we set $\xi_n = f^n(\xi)$, $n \in \mathbb{Z}_+$, $\eta_n = f^n(\eta)$, $n \in \mathbb{Z}_-$, $\eta = R\xi$, where $\mathbb{Z}_+ = \{0, 1, 2, \dots\}$ and $\mathbb{Z}_- = \{\dots, -2, -1, 0\}$. We note that $\eta_{-n} = R\xi_n$, $n \in \mathbb{Z}_+$. Then the linear system

$$(1) \quad v_{n+1} = Df(\xi_n)v_n, \quad n \in \mathbb{Z}_+,$$

has an exponential dichotomy on $\mathbb{Z}_+$ [8], i.e. there are positive constants $L, \delta \in (0, 1)$ and the orthogonal projection $P_\xi : \mathbb{R}^{2N} \rightarrow T_\xi \widetilde{W}_p^s$ such that the fundamental

solution $V_\xi(n)$ of (1) satisfies the following:

$$\|V_\xi(n)P_\xi V_\xi(m)^{-1}\| \leq L\delta^{n-m}, \quad m \leq n, \quad m, n \in \mathbb{Z}_+,$$

$$\|V_\xi(n)(I - P_\xi)V_\xi(m)^{-1}\| \leq L\delta^{m-n}, \quad n \leq m, \quad m, n \in \mathbb{Z}_+.$$

We note that L and δ can be chosen to be independent of $\xi \in \widetilde{W}_p^s \cap \bar{U}$ [1].

By defining $Rv_n = w_{-n}$ in (1), the reversibility of f implies

$$(2) \quad w_{n+1} = Df(\eta_n)w_n, \quad n \in \mathbb{Z}_-, \quad n \neq 0.$$

Hence, the fundamental solution $W_\xi(n)$ of (2) is given by $W_\xi(n) = RV_\xi(-n)R^{-1}$, $n \in \mathbb{Z}_-$, and since $\|R\| = \|R^{-1}\| = 1$, then (2) has an exponential dichotomy on $\mathbb{Z}_-$ with the constants L , δ and the orthogonal projection $I - Q_\eta$, where $Q_\eta = RP_\xi R^{-1}$. We note $\eta = R\xi$.

Now we fix $\omega \in \mathbb{N}$ large and put

$$J_\omega = \{-\omega, -\omega + 1, \dots, \omega - 1, \omega\},$$

$$J_\omega^- = \{-\omega, -\omega + 1, \dots, -1, 0\}, \quad I_\omega^- = \{-\omega, -\omega + 1, \dots, -1\},$$

$$J_\omega^+ = \{0, 1, \dots, \omega - 1, \omega\}, \quad I_\omega^+ = \{0, 1, \dots, \omega - 2, \omega - 1\}.$$

We note that the family $\{P_\xi \mid \xi \in \widetilde{W}_p^s \cap U\}$ is continuous on $\widetilde{W}_p^s \cap U$. In this paper, RL and NL denote, respectively, the range and the kernel of a linear operator L .

Theorem 2.1 ([1]). *There exist $\omega_0 \in \mathbb{N}$ and a constant $c > 0$ such that, for any $\omega \in \mathbb{N}$, $\omega \geq \omega_0$, and $\xi \in \widetilde{W}_p^s \cap U$, there exist unique $\{x_n^+(\omega, \xi)\}_{n \in J_\omega^+}$ and $\{x_n^-(\omega, \xi)\}_{n \in J_\omega^-}$ which satisfy $x_{n+1} = f(x_n)$ separately on I_ω^+ and I_ω^- such that*

$$P_\xi x_0^+(\omega, \xi) = P_\xi \xi, \quad Q_{R\xi} x_0^-(\omega, \xi) = Q_{R\xi} R\xi,$$

$$x_\omega^+(\omega, \xi) = x_{-\omega}^-(\omega, \xi),$$

together with

$$\max_{n \in J_\omega^+} |x_n^+(\omega, \xi) - \xi_n| \leq c\delta^\omega,$$

$$\max_{n \in J_\omega^-} |x_n^-(\omega, \xi) - \eta_n| \leq c\delta^\omega.$$

Moreover, $x_n^\pm(\omega, \xi)$ are continuous with respect to ξ .

Proof. We study the nonlinear system

$$(3) \quad x_{n+1} = f(x_n)$$

near $\{\xi_n\}_{n \in J_\omega^+}$ and $\{\eta_n\}_{n \in J_\omega^-}$. By putting $x_n^+ = \xi_n + v_n$, $n \in J_\omega^+$ and $x_n^- = \eta_n + w_n$, $n \in J_\omega^-$, we get the systems

$$(4) \quad \begin{aligned} v_{n+1} &= Df(\xi_n)v_n + f(\xi_n + v_n) - f(\xi_n) - Df(\xi_n)v_n \\ &= Df(\xi_n)v_n + o(|v_n|), \quad n \in I_\omega^+, \end{aligned}$$

and

$$(5) \quad \begin{aligned} w_{n+1} &= Df(\eta_n)w_n + f(\eta_n + w_n) - f(\eta_n) - Df(\eta_n)w_n \\ &= Df(\eta_n)w_n + o(|w_n|), \quad n \in I_\omega^-. \end{aligned}$$

Since we are looking for solutions of equation (3) such that $x_\omega^+ = x_\omega^-$, we add the boundary value conditions

$$(6) \quad v_\omega - w_{-\omega} = \eta_{-\omega} - \xi_\omega = O(\delta^\omega), \quad P_\xi v_0 = 0, \quad Q_{R\xi} w_0 = 0.$$

Let $v = (v_0, \dots, v_\omega) \in \mathbb{R}^{N(\omega+1)}$, $w = (w_{-\omega}, \dots, w_0) \in \mathbb{R}^{N(\omega+1)}$. To solve equations (4)–(6), we take the mapping $\Gamma_\omega : \widetilde{W}_p^s \cap \widetilde{U} \times \mathbb{R}^{2N(\omega+1)} \rightarrow \mathbb{R}^{2N(\omega+1)}$ defined by

$$\Gamma_\omega(\xi, v, w) = \begin{pmatrix} (v_{n+1} - f(\xi_n + v_n) + f(\xi_n))_{n \in I_\omega^+} \\ (w_{n+1} - f(\eta_n + w_n) + f(\eta_n))_{n \in I_\omega^-} \\ v_\omega - w_{-\omega} - (\eta_{-\omega} - \xi_\omega) \\ P_\xi v_0 \\ Q_{R\xi} w_0 \end{pmatrix}.$$

where $\begin{pmatrix} P_\xi v_0 \\ Q_{R\xi} w_0 \end{pmatrix}$ has to be meant as a vector in $\mathbb{R}^N = \mathcal{R}P_\xi \times \mathcal{R}Q_{R\xi}$. We already know that P_ξ and $Q_{R\xi}$ are continuous. Thus, for any fixed $\omega \geq \omega_0$, Γ_ω is continuous in (ξ, v, w) as well as its derivatives with respect to (v, w) when we take on $\mathbb{R}^{2N(\omega+1)}$ the maximum norm $\max_i \{|v_i|, |w_i|\}$. We have $\Gamma_\omega(\xi, 0, 0) = O(\delta^\omega)$ uniformly with respect to ξ and the linearized map $D_{(v,w)}\Gamma_\omega(\xi, 0, 0)$ has the form

$$D_{(v,w)}\Gamma_\omega(\xi, 0, 0) \begin{pmatrix} v \\ w \end{pmatrix} = \begin{pmatrix} (v_{n+1} - Df(\xi_n)v_n)_{n \in I_\omega^+} \\ (w_{n+1} - Df(\eta_n)w_n)_{n \in I_\omega^-} \\ v_\omega - w_{-\omega} \\ P_\xi v_0 \\ Q_{R\xi} w_0 \end{pmatrix}.$$

Arguing as in Lemma 2.1 of [1] or Lemma 2 of [2], the map $D_{(v,w)}\Gamma_\omega(\xi, 0, 0)$ is invertible and that its inverse is bounded uniformly with respect to ξ . Hence from the implicit function theorem we get that $c > 0$ and $\omega_0 \gg 1$ exist such that for $\omega \geq \omega_0$, the equation $\Gamma_\omega(\xi, v, w) = 0$ can be solved uniquely for (v, w) in a neighborhood of $(0, 0)$ in terms of (ξ, ω) . Moreover $\max_i \{|v_i|, |w_i|\} < c\delta^\omega$, and the solution is continuous in ξ , for any fixed $\omega \geq \omega_0$. The proof is finished.

Since $Q_{R\xi} = RP_\xi R^{-1}$, $\eta_{-n} = R\xi_n$, $n \in \mathbb{Z}_+$, we see that the sequences given by $y_n^-(\omega, \xi) = Rx_{-n}^+(\omega, \xi)$, $n \in J_\omega^-$, $y_n^+(\omega, \xi) = Rx_{-n}^-(\omega, \xi)$, $n \in J_\omega^+$ also satisfy the statement of Theorem 2.1. The uniqueness of such orbits implies that

$$\begin{aligned} x_n^+(\omega, \xi) &= y_n^+(\omega, \xi) = Rx_{-n}^-(\omega, \xi), & n \in J_\omega^+, \\ x_n^-(\omega, \xi) &= y_n^-(\omega, \xi) = Rx_{-n}^+(\omega, \xi), & n \in J_\omega^-. \end{aligned}$$

Hence $Rx_n^\pm(\omega, \xi) = x_{-n}^\mp(\omega, \xi)$, $n \in J_\omega^\pm$. So the orbit of f in Theorem 2.1 is R -symmetric.

3. R -SYMMETRIC PERIODIC ORBITS

In this section we prove Theorem 1.1. If $x_0^+(\omega, \xi) \in \text{Fix } R$, then

$$x_0^+(\omega, \xi) = Rx_0^+(\omega, \xi) = x_0^-(\omega, \xi).$$

Consequently, the orbit of Theorem 2.1 becomes an R -symmetric periodic orbit of f . Hence we have to solve the equation

$$(7) \quad (I - R)x_0^+(\omega, \xi) = 0, \quad \xi \in \widetilde{W}_p^s \cap U.$$

Let V be an open subset such that $K \subset V \subset \bar{V} \subset U$ and let ω_0 be as in Theorem 2.1. Note that the solution $x_0^+(\omega, \xi)$ is defined for $\xi \in \widetilde{W}_p^s \cap \bar{V}$ and

$$\#(\widetilde{W}_p^s \cap V, \text{Fix } R \cap V) = \#(\widetilde{W}_p^s \cap U, \text{Fix } R \cap U) \neq 0.$$

To solve (7), we put $F_\omega(\xi) = (I - R)x_0^+(\omega, \xi)$, $F_\omega : \widetilde{W}_p^s \cap \bar{V} \rightarrow R_- = \mathcal{R}(I - R)$. We note that $\dim R_- = \dim \widetilde{W}_p^s$.

Now we put F_ω into the homotopy

$$\begin{aligned} H_\omega : \widetilde{W}_p^s \cap \bar{V} \times [0, 1] &\rightarrow R_-, \\ H_\omega(\xi, \lambda) &= \lambda F_\omega(\xi) + (1 - \lambda)(I - R)\xi. \end{aligned}$$

Theorem 2.1 gives

$$|H_\omega(\xi, \lambda) - (I - R)\xi| = |\lambda(F_\omega(\xi) - (I - R)\xi)| \leq c\delta^\omega.$$

Consequently, $H_\omega(\cdot, \lambda) \neq 0$ on the boundary $\partial(\widetilde{W}_p^s \cap V)$ for any $0 \leq \lambda \leq 1$. This gives for the Brouwer degree [6],

$$\deg(F_\omega, \widetilde{W}_p^s \cap V, 0) = \pm \#(\widetilde{W}_p^s \cap V, \text{Fix } R \cap V) \neq 0.$$

Summarizing, we see that $F_\omega(\xi) = 0$ has a solution $\xi \in \widetilde{W}_p^s \cap V$ for any $\omega \geq \omega_0$, where ω_0 is sufficiently large. This proves Theorem 1.1.

4. PERTURBATION THEORY

In this section, we consider a C^2 -smooth perturbation $f(x, \varepsilon)$ of f , i.e. we suppose that $f(x, 0) = f(x)$ and $Rf(x, \varepsilon) = f^{-1}(Rx, \varepsilon)$, $\forall x \in \mathbb{R}^{2N}$, ε small. Then Theorem 2.1 gives a C^1 -mapping $x_0^+(\omega, \xi, \varepsilon)$ and we are led to the equation

$$(I - R)x_0^+(\omega, \xi, \varepsilon) = 0, \quad \xi \in \widetilde{W}_p^s \cap U.$$

By taking $\omega \rightarrow \infty$ in the above equation, we get $F(\xi, \varepsilon) = (I - R)x_0^+(\infty, \xi, \varepsilon) = 0$ which is precisely the equation of R -symmetric homoclinic solutions to the hyperbolic symmetric fixed point p_ε of $f(x, \varepsilon)$ near p [10]. We assume that this equation has a compact nondegenerate solution manifold, i.e.

(H1) There is an embedded compact C^2 -smooth submanifold $\mathcal{M} \subset \widetilde{W}_p^s \setminus \{p\} \cap \text{Fix } R$ for an open subset $\widetilde{W}_p^s$ of W_p^s which is a submanifold of $\mathbb{R}^{2N}$ and such that $\dim \mathcal{N}(I - R)D_\xi x_0^+(\infty, \xi, 0) = \dim \mathcal{M}$ for any $\xi \in \mathcal{M}$. Furthermore, let $\mathcal{O} \subset \widetilde{W}_p^s \setminus \{p\}$ be an open bounded subset such that $\mathcal{M} \subset \mathcal{O}$. Then $\mathcal{O}$ can be oriented. We suppose that $\mathcal{M}$ is orientable embedded into $\mathcal{O}$.

We note that always $T_\xi \mathcal{M} \subset \mathcal{N}(I - R)D_\xi x_0^+(\infty, \xi, 0)$, $\forall \xi \in \mathcal{M}$ and $\dim \mathcal{M} = \dim T_\xi \mathcal{M}$, hence (H1) implies $T_\xi \mathcal{M} = \mathcal{N}(I - R)D_\xi x_0^+(\infty, \xi, 0)$, $\forall \xi \in \mathcal{M}$. Since $x_0^+(\infty, \xi, 0) = \xi$, we have $D_\xi x_0^+(\infty, \xi, 0)v = v$, $v \in T_\xi \widetilde{W}_p^s$. Hence

$$\mathcal{N}(I - R)D_\xi x_0^+(\infty, \xi, 0) = \text{Fix } R \cap T_\xi \widetilde{W}_p^s.$$

Now we take a tubular neighbourhood $\mathcal{V}$ of $\mathcal{M}$ in $\widetilde{W}_p^s$, i.e. any $\xi \in \mathcal{V}$ can be uniquely expressed as a pair (τ, v) , where $\tau \in \mathcal{M}$ and $v \in T_\tau \widetilde{W}_p^s / T_\tau \mathcal{M} = T_\tau \widetilde{W}_p^s / (\text{Fix } R \cap T_\tau \widetilde{W}_p^s) = N_\tau$ —the fiber of the normal vector bundle of $\mathcal{M}$ in $\widetilde{W}_p^s$, and $|v| < \Delta$ for some $\Delta > 0$. Hence we identify $\mathcal{V}$ with an open neighbourhood of the zero section of the normal vector bundle of $\mathcal{M}$ in $\widetilde{W}_p^s$. Let $S_\tau : \text{Fix}(-R) \rightarrow \mathcal{R}D_v F(\tau, 0, 0)$ be the orthogonal projection. We note that the assumption (H1)

implies the invertibility of the linear mapping $D_v F(\tau, 0, 0) : N_\tau \rightarrow \mathcal{R}D_v F(\tau, 0, 0)$, since $D_v F(\tau, 0, 0)w = (I - R)w$, $w \in N_\tau$.

From $F(\tau, 0, 0) = 0$, we get $F(\tau, v, \varepsilon) = D_v F(\tau, 0, 0)v + \varepsilon D_\varepsilon F(\tau, 0, 0) + o(|v|) + o(\varepsilon)$. We consider the homotopy

$$\begin{aligned} H(\tau, v, \varepsilon, \lambda) &= S_\tau(\lambda F(\tau, v, \varepsilon) + (1 - \lambda)D_v F(\tau, 0, 0)v) \\ &\quad + (I - S_\tau)(\lambda F(\tau, v, \varepsilon) + (1 - \lambda)\varepsilon D_\varepsilon F(\tau, 0, 0)). \end{aligned}$$

Then $H(\tau, v, \varepsilon, 0) = S_\tau D_v F(\tau, 0, 0)v + \varepsilon(I - S_\tau)D_\varepsilon F(\tau, 0, 0)$ and $H(\tau, v, \varepsilon, 1) = F(\tau, v, \varepsilon)$.

Now we suppose

(H2) There is an open connected subset $\Omega \subset \mathcal{M}$ such that $B(\tau) \neq 0$, $\forall \tau \in \partial\Omega$, where $B(\tau) = (I - S_\tau)D_\varepsilon F(\tau, 0, 0)$.

Since $\mathcal{M}$ is orientable embedded into $\mathcal{O}$ and $\mathcal{O}$ is orientable, the tangent vector bundle $T\mathcal{M}$ and the normal vector bundle $\bigcup_{\tau \in \mathcal{M}} N_\tau$ are both orientable. Hence the vector bundle $\bigcup_{\tau \in \mathcal{M}} (I - R)N_\tau = \bigcup_{\tau \in \mathcal{M}} \mathcal{R}D_v F(\tau, 0, 0)$ is also orientable, because $D_v F(\tau, 0, 0) : N_\tau \rightarrow \mathcal{R}D_v F(\tau, 0, 0)$ is invertible. Since $S_\tau : \text{Fix}(-R) \rightarrow \mathcal{R}D_v F(\tau, 0, 0)$ is the orthogonal projection and the vector bundle $\bigcup_{\tau \in \mathcal{M}} \text{Fix}(-R) = \mathcal{M} \times \text{Fix}(-R)$ is orientable, we get the orientability of the vector bundle $\bigcup_{\tau \in \mathcal{M}} \mathcal{R}(I - S_\tau)\text{Fix}(-R)$. Since $B(\tau)$ is a section of this vector bundle, the following assumption makes sense [7].

(H3) $\deg(B(\tau), \Omega, 0) \neq 0$.

According to (H2), there is an open connected bounded neighbourhood $U_1 \subset \mathcal{M}$ of $\bar{\Omega}$ such that $B(\lambda) \neq 0$, $\forall \tau \in U_1 \setminus \Omega$. Now we take an open subset $V_\varepsilon = \{(\tau, v) \in V \mid \tau \in U_1 \text{ and } |v| < |\varepsilon|r_1\}$ for a positive constant r_1 and $0 < |\varepsilon| < \Delta/r_1$. Since

$$S_\tau(\lambda F(\tau, v, \varepsilon) + (1 - \lambda)D_v F(\tau, 0, 0)v) = S_\tau D_v F(\tau, 0, 0)v + o(|v|) + O(\varepsilon)$$

and $S_\tau D_v F(\tau, 0, 0) : N_\tau \rightarrow \mathcal{R}D_v F(\tau, 0, 0)$ is invertible, we get that

$$S_\tau(\lambda F(\tau, v, \varepsilon) + (1 - \lambda)D_v F(\tau, 0, 0)v) \neq 0,$$

$\forall (\tau, v) \in V_\varepsilon$, $|v| = r_1|\varepsilon|$ for a r_1 sufficiently large and fixed. Furthermore, we have

$$(I - S_\tau)(\lambda F(\tau, v, \varepsilon) + (1 - \lambda)\varepsilon D_\varepsilon F(\tau, 0, 0)) = \varepsilon(I - S_\tau)D_\varepsilon F(\tau, 0, 0) + o(|v|) + o(\varepsilon)$$

since $(I - S_\tau)D_v F(\tau, 0, 0) = 0$. Hence $(I - S_\tau)(\lambda F(\tau, v, \varepsilon) + (1 - \lambda)\varepsilon D_\varepsilon F(\tau, 0, 0)) \neq 0$ for $(\tau, v) \in V_\varepsilon$, $|v| \leq r_1|\varepsilon|$ and $\tau \in U_1 \setminus \Omega$.

Summarizing, we see that $H(\tau, v, \varepsilon, \lambda) \neq 0$ for any $(\tau, v) \in \partial V_\varepsilon$, $\lambda \in [0, 1]$ and $\varepsilon \neq 0$ sufficiently small. Consequently [6],

$$\deg(F, V_\varepsilon, 0) = \deg(H(\cdot, \varepsilon, 0), V_\varepsilon, 0),$$

where $H(\tau, v, \varepsilon, 0) = S_\tau D_v F(\tau, v(\tau, \varepsilon), \varepsilon)v + \varepsilon(I - S_\tau)D_\varepsilon F(\tau, 0, 0)$. Since the linear map $S_\tau D_v F(\tau, v(\tau, \varepsilon), \varepsilon) : N_\tau \rightarrow \mathcal{R}D_v F(\tau, 0, 0)$ is invertible and U_1 is connected, we get

$$\deg(H(\cdot, \varepsilon, 0), V_\varepsilon, 0) = \pm \deg(B(\tau), \Omega, 0) \neq 0.$$

This implies $\#(\widetilde{W}_{p_\varepsilon}^s \cap V_\varepsilon, \text{Fix } R \cap V_\varepsilon) \neq 0$. Summarizing we get the following result.

Theorem 4.1. *Assume (H1), (H2) and (H3). Then there exists $\varepsilon_0 > 0$ such that for $0 < |\varepsilon| \leq \varepsilon_0$, it is nonzero the local intersection number of the plain $\text{Fix } R$ and the stable manifold of the hyperbolic fixed point of the map $x_{n+1} = f(x_n, \varepsilon)$ which is located near the fixed point p of the map $x_{n+1} = f(x_n)$.*

Hence Theorem 1.1 and the assumptions of Theorem 4.1 imply an infinite number of R -symmetric periodic orbits of $f(x, \varepsilon)$ accumulating on R -symmetric homoclinic orbits of $f(x, \varepsilon)$ for any $0 < |\varepsilon| \leq \varepsilon_0$.

By taking $\Omega = \mathcal{M}$, we see [7] that assumption (H3) holds if the Euler characteristic $\chi(\bigcup_{\tau \in \mathcal{M}} \mathcal{R}(I - S_\tau) \text{Fix}(-R))$ is nonzero. Then Theorem 4.1 holds under assumption (H1) for any R -reversible C^2 -smooth perturbation $f(x, \varepsilon)$.

If $f(x, \varepsilon)$ is C^3 -smooth, then F is C^2 -smooth. To solve $F(\tau, v, \varepsilon) = 0$, we follow the standard way [1] by splitting it as $F(\tau, v, \varepsilon) = S_\tau F(\tau, v, \varepsilon) + (I - S_\tau)F(\tau, v, \varepsilon)$. By using the implicit function theorem, we can solve the equation $S_\tau F(\tau, v, \varepsilon) = 0$ in v for ε small and $\tau \in \mathcal{M}$ to get the C^2 -smooth solution $v = v(\tau, \varepsilon) = O(\varepsilon)$. Then we consider the bifurcation equation $C(\tau, \varepsilon) = (I - S_\tau)F(\tau, v(\tau, \varepsilon), \varepsilon) = 0$. Clearly $C(\tau, \varepsilon)/\varepsilon \rightarrow B(\tau)$ in the C^1 -topology on $\mathcal{M}$ as $\varepsilon \rightarrow 0$. Consequently, a simple zero τ_0 of $B(\tau)$, i.e. $B(\tau_0) = 0$ and $DB(\tau_0)$ is nonsingular, implies the solvability of $C(\tau, \varepsilon) = 0$ in τ near τ_0 for $\varepsilon \neq 0$ small. Summarizing we see that B is the Melnikov function for this problem, since its simple zero τ_0 ensures the bifurcation of an R -symmetric homoclinic orbit of $f(x, \varepsilon)$ to p_ε for $\varepsilon \neq 0$ small bifurcating from the R -symmetric homoclinic orbit of $f(x, 0) = f(x)$ which starts from $\tau_0 \in \mathcal{M}$.

5. AN EXAMPLE

In this section, we present an example, but first we simplify the formula of $B(\tau)$. If $a \in \mathcal{R}(I - S_\tau)$, then $a \in \text{Fix}(-R)$ and $a \perp (I - R)T_\tau \widetilde{W}_p^s$. Hence for any $w \in T_\tau \widetilde{W}_p^s$ we have

$$0 = \langle a, (I - R)w \rangle = \langle a, w \rangle - \langle a, Rw \rangle = \langle a - Ra, w \rangle = 2\langle a, w \rangle.$$

Furthermore, since $RT_\tau \widetilde{W}_p^s = T_{R\tau} \widetilde{W}_p^u = T_\tau \widetilde{W}_p^u$ and for any $w \in T_\tau \widetilde{W}_p^s$, we have

$$\langle a, Rw \rangle = -\langle Ra, w \rangle = -\langle a, w \rangle = 0,$$

we see that $a \in \mathcal{R}(I - S_\tau)$ if and only if $a \perp (T_\tau \widetilde{W}_p^s + T_\tau \widetilde{W}_p^u + \text{Fix } R)$. We note that $\text{Fix}(-R) = (\text{Fix } R)^\perp$, and $\frac{1}{2}(I - R) : \mathbb{R}^{2N} \rightarrow \text{Fix}(-R)$ and $\frac{1}{2}(I + R) : \mathbb{R}^{2N} \rightarrow \text{Fix } R$ are the orthogonal projections. Consequently, if $a_i(\tau), i = 1, 2, \dots, \dim \mathcal{M}$, is a continuous orthonormal vector field such that $a_i(\tau) \perp (T_\tau \widetilde{W}_p^s + T_\tau \widetilde{W}_p^u + \text{Fix } R)$, then the components of $B(\tau)$ are given by

$$B_i(\tau) = \langle a_i(\tau), (I - R)D_\varepsilon x_0^+(\infty, \tau, 0) \rangle = 2\langle a_i(\tau), D_\varepsilon x_0^+(\infty, \tau, 0) \rangle.$$

Now we consider a perturbed second order differential equation

$$(8) \quad \ddot{z} = g(z) + \varepsilon h(z), \quad z \in \mathbb{R}^N,$$

where $g, h \in C^3(\mathbb{R}^N, \mathbb{R}^N)$, $g(0) = h(0) = 0$. (8) has the form

$$(9) \quad \dot{z}_1 = z_2, \quad \dot{z}_2 = g(z_1) + \varepsilon h(z_1).$$

Let $\phi(t, z_1, z_2, \varepsilon)$ be the flow of (9), then $f(x, \varepsilon) = \phi(T, x, \varepsilon)$, $x = (z_1, z_2)$ for a $T > 0$. Here $R(z_1, z_2) = (z_1, -z_2)$ and $\text{Fix } R = \{(z_1, 0) \mid z_1 \in \mathbb{R}^N\}$, $\text{Fix}(-R) = \{(0, z_2) \mid z_2 \in \mathbb{R}^N\}$. The inner product $\langle \cdot, \cdot \rangle$ is given by $\langle (z_1^1, z_2^1), (z_1^2, z_2^2) \rangle = (z_1^1, z_2^1) + (z_1^2, z_2^2)$, where $(\cdot, \cdot)$ is the usual inner product on $\mathbb{R}^N$. We assume that $p_\varepsilon = (0, 0)$ is a hyperbolic equilibrium of (9). Let $\tau \in \text{Fix } R \cap \widetilde{W}_p^s$. Then $\phi(t, \tau, 0) = (z_1^T(t), z_2^T(t))$ with $z_1^T(t)$ even and $z_2^T(t)$ odd. $\phi(t, \tau, 0)$ is a homoclinic solution of (9) with $\varepsilon = 0$. The linearization of (9) for $\varepsilon = 0$ along $\phi(t, \tau, 0)$ has the form

$$(10) \quad \dot{v} = w, \quad \dot{w} = Dg(z_1^T(t))v.$$

It is clear that $T_\tau \widetilde{W}_p^{s(u)} = \{(v(0), w(0)) \mid v(t), w(t) \text{ are bounded solutions of (10) on } \mathbb{R}_+(-)\}$, respectively. According to [2], the condition $a \perp (T_\tau \widetilde{W}_p^s + T_\tau \widetilde{W}_p^u + \text{Fix } R)$ is now equivalent to $a = (0, a_2)$ and $v_1(0) = 0$, $w_1(0) = a_2$, where $v_1(t)$ and $w_1(t)$ are the bounded solutions on $\mathbb{R}$ of the adjoint system of (10), i.e. w_1 is the even bounded solution on $\mathbb{R}$ of $\ddot{w}_1 = Dg(z_1^\tau(t))^* w_1$, $w_1(0) = a_2$. Furthermore, $D_\varepsilon x_0^+(\infty, \tau, 0) = (v_\tau(0), w_\tau(0)) \perp T_\tau \widetilde{W}_p^s$, where $v_\tau(t)$ and $w_\tau(t)$ are the bounded solutions on $\mathbb{R}_+$ of the equations

$$(11) \quad \dot{v}_\tau = w_\tau, \quad \dot{w}_\tau = Dg(z_1^\tau(t))v_\tau + h(z_1^\tau(t)).$$

Consequently, the corresponding component of $B(\tau)$ to a derived above is given by

$$2\langle a, D_\varepsilon x_0^+(\infty, \tau, 0) \rangle = 2\langle (0, a_2), (v_\tau(0), w_\tau(0)) \rangle = 2(a_2, w_\tau(0)) = 2(w_1(0), w_\tau(0)).$$

On the other hand, since (11) holds along with $\lim_{t \rightarrow +\infty} w_1(t) = 0$, we have

$$(12) \quad \int_0^\infty (h(z_1^\tau(t)), w_1(t)) dt = (w_1(0), w_\tau(0)).$$

To be more concrete, we consider the system

$$(13) \quad \ddot{x} = x - 2x(x^2 + y^2), \quad \ddot{y} = y - 2y(x^2 + y^2) + \varepsilon x^4, \quad x, y \in \mathbb{R}.$$

(13) has for $\varepsilon = 0$ a homoclinic manifold $x_\tau(t) = \sin \tau r(t)$, $y_\tau(t) = \cos \tau r(t)$, $r(t) = \text{sech } t$, which intersects $\text{Fix } R$ in the circle $\mathcal{M} = (\sin \tau, \cos \tau, 0, 0)$. It is not difficult to observe that now assumption (H1) holds and $w_1(t) = (y_\tau(t), -x_\tau(t))$. By (12), the function $B(\tau)$ now has the form

$$B(\tau) = -2 \int_0^\infty \sin^5 \tau r^5(t) dt = -\frac{3}{8} \pi \sin^5 \tau.$$

We note that $x_0(t)$, $y_0(t)$ are the even solutions of (13). The bifurcation equation $C(\tau, \varepsilon) = 0$ from Section 4 is now analytical. Hence $\tau = 0$ is its isolated solution for $\varepsilon \neq 0$ small and fixed. The Brouwer degree of $B(\tau)$ at $\tau = 0$ is -1 , so Theorem 4.1 implies the following result.

Theorem 5.1. *The point $(0, 1, 0, 0)$ is an isolated topologically transversal intersection of $W_{p_\varepsilon}^s$ and $\text{Fix } R$ for (13) with $\varepsilon \neq 0$ small. But this point is not a C^1 -transversal intersection.*

Proof. To prove the non- C^1 -transversal intersection, we consider a C^3 -perturbation of (13) given by

$$(14) \quad \ddot{x} = x - 2x(x^2 + y^2), \quad \ddot{y} = y - 2y(x^2 + y^2) + \varepsilon \phi_\delta(x), \quad x, y \in \mathbb{R},$$

where $\delta > 0$ and $\phi_\delta(x) = 0$ if $|x| \leq \delta$, $\phi_\delta(x) = (x - \delta \text{sgn } x)^4$ if $|x| \geq \delta$. We see that (14) has even homoclinics $x_\tau(t)$, $y_\tau(t)$ for $|\sin \tau| < \delta$ and any ε . Hence $(0, 1, 0, 0)$ is not an isolated reversible homoclinic point for the C^3 -perturbation (14) with $\varepsilon \neq 0$, $\delta > 0$ small of the system (13). The proof is finished.

REFERENCES

- [1] Battelli, F., Fečkan, M., *Chaos arising near a topologically transversal homoclinic set*, preprint.
- [2] Battelli, F., Fečkan, M., *Subharmonic solutions in singular systems*, J. Differential Equations, **132** (1996), pp. 21-45. MR **98k**:34068
- [3] Devaney, R., *Blue sky catastrophes in reversible and Hamiltonian systems*, Indiana Univ. Math. J., **26** (1977), pp. 247-263. MR **55**:4275

- [4] Devaney, R., *Reversible diffeomorphisms and flows*, Tran. Amer. Math. Soc., **218** (1976), pp. 89-113. MR **53**:6629
- [5] Devaney, R., *Homoclinic bifurcations and the area-conserving Hénon mapping*, J. Differential Equations, **51** (1984), pp. 254-266. MR **85k**:58054
- [6] Fonseca, I., Gangbo, W., Degree Theory in Analysis and Applications, Oxford Lec. Ser. Math. Appl. **2**, Clarendon Press, Oxford (1995). MR **96k**:47100
- [7] Hirsch, M.W., Differential Topology, Springer-Verlag, New York (1976). MR **56**:6669
- [8] Palmer, K. J., *Exponential dichotomies, the shadowing lemma and transversal homoclinic points*, Dynamics Report Ser. Dynam. Systems Appl. **1** (1988), pp. 265-306. MR **89j**:58060
- [9] Vanderbauwhede, A., *Heteroclinic cycles and periodic orbits in reversible systems*, in J. Wiener & J.K. Hale (Ed.) "Ordinary and Delay Differential Equations", Pitman Research Notes in Mathematics Series, No. **272** Pitman 1992, pp. 250-253. CMP 97:08
- [10] Vanderbauwhede, A., Fiedler, B., *Homoclinic period blow-up in reversible and conservative systems*, Z. Angew. Math. Phys. (ZAMP), **43** (1992), 292-318. MR **93f**:58209

DEPARTMENT OF MATHEMATICAL ANALYSIS, COMENIUS UNIVERSITY, MLYNSKÁ DOLINA, 842 48
BRATISLAVA, SLOVAKIA

E-mail address: Michal.Feckan@fmph.uniba.sk