FRAME WAVELETS IN SUBSPACES OF $L^2(\mathbb{R}^d)$

X. DAI, Y. DIAO, Q. GU, AND D. HAN

(Communicated by David R. Larson)

ABSTRACT. Let A be a $d \times d$ real expansive matrix. We characterize the reducing subspaces of $L^2(\mathbb{R}^d)$ for A -dilation and the regular translation operators acting on $L^2(\mathbb{R}^d)$. We also characterize the Lebesgue measurable subsets E of $\mathbb{R}^d$ such that the function defined by inverse Fourier transform of $[1/(2\pi)^{d/2}]\chi_E$ generates through the same A -dilation and the regular translation operators a normalized tight frame for a given reducing subspace. We prove that in each reducing subspace, the set of all such functions is nonempty and is also path connected in the regular $L^2(\mathbb{R}^d)$ -norm.

1. INTRODUCTION

A sequence $\{x_n\}$ in a Hilbert space $\mathcal{H}$ is called a *frame* for $\mathcal{H}$ if there exist constants $C_1, C_2 > 0$ such that

$$C_1\|x\|^2 \leq \sum_{n \in \mathbb{N}} |\langle x, x_n \rangle|^2 \leq C_2\|x\|^2, \quad \forall x \in \mathcal{H}.$$

If $C_1 = C_2 = 1$, it is called a *normalized tight* frame. It is known ([6]) that $\{x_n\}$ is a normalized tight frame for $\mathcal{H}$ if and only if $x = \sum_{n \in \mathbb{N}} \langle x, x_n \rangle x_n$ for all $x \in \mathcal{H}$, where the convergence is unconditional in norm.

In this article we will investigate a class of normalized tight frames for either $L^2(\mathbb{R}^d)$ or certain subspaces of $L^2(\mathbb{R}^d)$ which are called reducing subspaces. The normalized tight frames we will deal with are obtained by applying certain A -dilation and regular translation operators to a single Lebesgue integrable function. Let us first define the above mentioned operators.

Let A be a $d \times d$ real *invertible* matrix. It induces a unitary operator D_A acting on $L^2(\mathbb{R}^d)$ defined by

$$(1) \quad (D_A f)(t) = |\det A|^{\frac{1}{2}} f(At), \quad \forall f \in L^2(\mathbb{R}^d), t \in \mathbb{R}^d.$$

The matrix A is called *expansive* if all its eigenvalues have modulus greater than one. The operator D_A corresponding to a real expansive matrix A is called an *A -dilation* operator. In an analogous fashion, a vector s in $\mathbb{R}^d$ induces a unitary

Received by the editors January 5, 2001 and, in revised form, February 26, 2001 and June 6, 2001.

2000 *Mathematics Subject Classification.* Primary 42-XX, 47-XX.

Key words and phrases. Normalized tight frame wavelet set, reducing subspace, connectivity.

translation operator T_s defined by

$$(T_s f)(t) = f(t - s), \quad \forall f \in L^2(\mathbb{R}^d), t \in \mathbb{R}^d.$$

In this article we will only deal with expansive real matrices and translation operators T_ℓ with $\ell \in \mathbb{Z}^d$.

Throughout this article, we will use $\mathcal{F}$ to denote the Fourier-Plancherel transform on $L^2(\mathbb{R}^d)$. This is a unitary operator. If $f \in L^2(\mathbb{R}^d) \cap L^1(\mathbb{R}^d)$, then

$$(2) \quad (\mathcal{F}f)(s) = \frac{1}{(2\pi)^{d/2}} \int_{\mathbb{R}^d} e^{-i(s \circ t)} f(t) dm,$$

where $s \circ t$ denotes the real inner product. We also write $\widehat{f}$ for $\mathcal{F}f$. For a subset X of $L^2(\mathbb{R}^d)$, $\widehat{X}$ is the set of the Fourier-Plancherel transforms of all elements in X . For a bounded linear operator S on $L^2(\mathbb{R}^d)$, we will denote $\mathcal{F}S\mathcal{F}^{-1}$ by $\widehat{S}$. It is left to the reader to verify that we have $\widehat{D}_A = D_{(A')^{-1}} = D_{A'}^{-1} = D_{A'}^*$ for any $d \times d$ real invertible matrix A (A' is the transpose of A) and $\widehat{T}_\lambda f = e^{-i(\lambda \circ s)} \cdot f$ for any $\lambda \in \mathbb{R}^d$.

A function ψ is called an *A-dilation orthonormal wavelet* if the family of functions $\{D_A^n T_\ell \psi : n \in \mathbb{Z}, \ell \in \mathbb{Z}^d\}$ forms an orthonormal basis for $L^2(\mathbb{R}^d)$. When the above family forms a *normalized tight frame* for $L^2(\mathbb{R}^d)$, the function $\psi \in L^2(\mathbb{R}^d)$ is called an *A-dilation normalized tight frame wavelet*. The existence of *A-dilation orthonormal wavelets* was proved in [3] for any expansive matrix A . In fact, it was proved in [3] that there exist *A-dilation orthonormal wavelets* whose Fourier transforms are of the form $\frac{1}{(2\pi)^{d/2}} \chi_E$ for some measurable subset E of $\mathbb{R}^d$. Such *A-dilation orthonormal wavelets* are called *s-elementary wavelets* [2]. A function ψ defined by $\widehat{\psi} = \frac{1}{(2\pi)^{d/2}} \chi_E$ for some measurable set E of $\mathbb{R}^d$ is called an *s-frame wavelet* if $\{D_A^n T_\ell \psi : n \in \mathbb{Z}, \ell \in \mathbb{Z}^d\}$ forms a normalized tight frame for $L^2(\mathbb{R}^d)$. The set E in this case will be called an *s-frame wavelet set* for short. It is proved in [7] that for any given real expansive matrix A , the set of all *s-elementary wavelets* is path-connected in the L^2 -norm. In this paper, we are interested in the *s-frame wavelets* defined on a *reducing subspace* X of $L^2(\mathbb{R}^d)$ (X is a reducing subspace if $D_A X = X$ and $T_\ell X = X$ for each $\ell \in \mathbb{Z}^d$). We characterize the *s-frame wavelet sets* and show that the set of all *s-frame wavelets* (on any given reducing subspace) is also path-connected in the L^2 -norm.

2. SUBSPACE *s*-FRAME WAVELETS

In this section, we will characterize all reducing subspaces and all *s-frame wavelet sets* for a given reducing subspace. We will begin with the following definition.

Definition 1. A function ψ is an *s-frame wavelet* on X if $\widehat{\psi} = \frac{1}{(2\pi)^{d/2}} \chi_E$ for some measurable subset E of $\mathbb{R}^d$, $\{D_A^n T_\ell \psi : n \in \mathbb{Z}, \ell \in \mathbb{Z}^d\} \subset X$ and

$$f = \sum_{n \in \mathbb{Z}, \ell \in \mathbb{Z}^d} \langle f, D_A^n T_\ell \psi \rangle D_A^n T_\ell \psi, \quad \forall f \in X.$$

We also need the following lemmas in the proof of our theorems.

Lemma 1. Let A be a $d \times d$ real expansive matrix. Then $\lim_{k \rightarrow +\infty} \|A^{-k}\| = 0$ and $\lim_{k \rightarrow +\infty} \|A^k t\| = \infty$ for every nonzero element t in $\mathbb{R}^d$.

Proof. Since A is an expansive matrix, it follows that A is invertible and for any $\lambda \in \sigma(A^{-1})$, $|\lambda| < 1$ and thus $\lim_{k \rightarrow +\infty} \lambda^{-k} = 0$. This implies that ([5], p. 559, Theorem 9) $\lim_{k \rightarrow +\infty} \|A^{-k}\| = 0$. Now let $t \in \mathbb{R}^d$ with $t \neq 0$. The result then follows from the fact that $\|A^{-k}\| \|A^k t\| \geq \|t\|$. $\square$

Lemma 2. *Let $\{e_n : n \in \mathbb{N}\}$ be an orthonormal basis for a separable Hilbert space $\mathcal{H}$ and let P be an orthogonal projection onto a subspace $\mathcal{H}_0$. Then $\{Pe_n\}$ is a normalized tight frame for $\mathcal{H}_0$.*

Lemma 3. *Let $\{\mathcal{H}_n : n \in \mathbb{N}\}$ be a family of mutually orthogonal subspaces that sum to $\mathcal{H}$. Suppose that for each $n \in \mathbb{N}$, $\{x_{nm} : m \in \mathbb{N}\}$ is a normalized tight frame for $\mathcal{H}_n$. Then $\{x_{nm} : n, m \in \mathbb{N}\}$ is a normalized tight frame for $\mathcal{H}$.*

The proofs of Lemmas 2 and 3 are left to the reader.

Lemma 4. *Let A be a $d \times d$ real expansive matrix. Then $\{A^n \mathbb{Z}^d : n \in \mathbb{Z}\}$ is dense in $\mathbb{R}^d$.*

Proof. Let $\varepsilon > 0$. By Lemma 1 we can choose $k \in \mathbb{N}$ such that $\|A^{-k}\| < \frac{\varepsilon}{d}$. Let $\{e_j : 1 \leq j \leq d\}$ be the standard orthonormal basis for $\mathbb{R}^d$. Since A is expansive hence invertible, the set $\{A^{-k}e_j : 1 \leq j \leq d\}$ is also a basis for $\mathbb{R}^d$. Let $x \in \mathbb{R}^d$, then $x = \sum_{j=1}^d x^{(j)} A^{-k}e_j$ with some $x^{(j)} \in \mathbb{R}$ for $j \in \{1, 2, \dots, d\}$. Let $\ell = \sum_{j=1}^d [x^{(j)}]e_j \in \mathbb{Z}^d$, where $[x^{(j)}]$ denotes the largest integer among those that are no greater than $x^{(j)}$, then $\|A^{-k}\ell - x\| = \|\sum_{j=1}^d (x^{(j)} - [x^{(j)}])A^{-k}e_j\| \leq \sum_{j=1}^d \|A^{-k}\| \cdot \|e_j\| < \varepsilon$. $\square$

We now characterize the reducing subspaces in the following theorem.

Theorem 1. *Let A be a $d \times d$ real expansive matrix. A closed subspace X of $L^2(\mathbb{R}^d)$ is a reducing subspace if and only if $\hat{X} = L^2(\mathbb{R}^d) \cdot \chi_\Omega$ for some Lebesgue measurable subset Ω of $\mathbb{R}^d$ with the property that $\Omega = A'\Omega$.*

Proof. Let X be a reducing subspace and P be the orthogonal projection from $L^2(\mathbb{R}^d)$ onto X . Then P commutes with both D_A^n and T_ℓ for all $n \in \mathbb{Z}$ and $\ell \in \mathbb{Z}^d$. Note that $D_A^{-n} T_\ell D_A^n = T_{A^n \ell}$. It follows that P commutes with $T_{A^n \ell}$ for all $n \in \mathbb{Z}$ and $\ell \in \mathbb{Z}^d$. By Lemma 4, the set $\{A^n \mathbb{Z}^d : n \in \mathbb{Z}\}$ is dense in $\mathbb{R}^d$. So P commutes with T_t for all $t \in \mathbb{R}^d$. Hence $\hat{P}$ commutes with $\hat{T}_t$ for all $t \in \mathbb{R}^d$. If we use M_f to denote the multiplicative operator by $f(s)$, then $\hat{T}_t = M_{e^{-t \circ s}}$. $\{M_{e^{-t \circ s}} : t \in \mathbb{R}^d\}$ generates a maximal abelian von Neumann algebra $\mathcal{A} = \{M_f : f \in L^\infty(\mathbb{R}^d)\}$. Thus $\hat{P}$ must be in $\mathcal{A}$ and $\hat{P} = M_f$ for some $f \in L^\infty(\mathbb{R}^d)$. The relation $P^2 = P$ implies that $f^2 = f$. So f is χ_Ω for some measurable subset Ω of $\mathbb{R}^d$. Thus $\hat{X} = L^2(\mathbb{R}^d) \cdot \chi_\Omega$. Since $PD_A = D_A P$, we have $\hat{P} \hat{D}_A = \hat{D}_A \hat{P}$. Equivalently $M_{\chi_\Omega} D_{A'}^{-1} = D_{A'}^{-1} M_{\chi_\Omega}$, hence $M_{\chi_\Omega} D_{A'} = D_{A'} M_{\chi_\Omega}$. Let $f \in L^2(\mathbb{R}^d)$. Observe that

$$\begin{aligned} M_{\chi_\Omega} D_{A'} f(t) &= |A'|^{\frac{1}{2}} f(A't) \cdot \chi_\Omega(t), \\ D_{A'} M_{\chi_\Omega} f(t) &= |A'|^{\frac{1}{2}} f(A't) \cdot \chi_\Omega(A't). \end{aligned}$$

This implies that $\chi_\Omega(A't) = \chi_\Omega(t)$, therefore $A'\Omega = \Omega$. The proof of the other direction is simple and is omitted. $\square$

Remark. The above proof shows that the commutant of $\{\hat{D}_A^n, \hat{T}_\ell, \ell \in \mathbb{Z}^d\}$ is $\{M_f : f \in L^\infty(\mathbb{R}^d), f(A'x) = f(x)\}$.

Let E be a subset of $\mathbb{R}^d$. Two points $x, y \in E$ are said to be 2π -translation equivalent if $x - y = 2\pi\ell$ for some $\ell \in \mathbb{Z}^d$. This is an equivalence relation on E . The 2π -translation redundancy index of a point x in E is the cardinality of its equivalence class. We use $E(\tau, k)$ to denote the set of all points in E with 2π -translation redundancy index k . For $k \neq m$, $E(\tau, k) \cap E(\tau, m) = \emptyset$, so we have $E = E(\tau, \infty) \cup (\bigcup_{n \in \mathbb{N}} E(\tau, n))$. Similarly, two nonzero points $x, y \in E$ are said to be A -dilation equivalent if $y = A^k x$ for some $k \in \mathbb{Z}$. This is also an equivalence relation on E . The A -dilation redundancy index of a point x in E is the cardinality in its equivalence class. The set of all points in E with A -dilation redundancy index k is denoted by $E(\delta_A, k)$. For $k \neq m$, $E(\delta_A, k) \cap E(\delta_A, m) = \emptyset$. So $E = E(\delta_A, \infty) \cup (\bigcup_{n \in \mathbb{N}} E(\delta_A, n))$. For a Lebesgue measurable set E of $\mathbb{R}^d$, $E(\tau, k)$ and $E(\delta_A, k)$ are both measurable for any $k \in \mathbb{N} \cup \{\infty\}$. The proof of case $d = 1$ can be found in [1] and the general cases can be treated similarly. The details are left to the reader. Furthermore, $E(\delta_A, k)$ can be decomposed as a union of k disjoint measurable subsets $E^{(j)}(\delta_A, k)$ ($1 \leq j \leq k$) such that each $E^{(j)}(\delta_A, k)$ contains only points of A -dilation redundancy index 1. A set E is said to be the A -dilation generator of another set Ω if $E = E(\delta_A, 1)$ and $\bigcup_{k \in \mathbb{Z}} A^k E = \Omega$. Two sets are said to be A -dilation equivalent if they are both A -dilation generators of the same set.

Before we state and prove the next theorem, we need to point out that all the statements in the theorems of this paper about a measurable set are to be modulus a null set.

Theorem 2. *Let A be a $d \times d$ real expansive matrix and X be a reducing subspace. Then a measurable subset $E \subset \mathbb{R}^d$ is an s -frame wavelet set for X if and only if $E = E(\tau, 1) = E(\delta_{A'}, 1)$ and $\bigcup_{k \in \mathbb{Z}} (A')^k E = \Omega$ satisfies the conditions in Theorem 1.*

Proof. “ $\Leftarrow$ ” Since E is 2π -translation equivalent to a subset F of $[0, 2\pi)^d$, the set $G = E \cup ([0, 2\pi)^d \setminus F)$ is 2π -translation equivalent to $[0, 2\pi)^d$ and $\{\widehat{T}_\ell \widehat{\psi}_G : \ell \in \mathbb{Z}^d\}$ is an orthogonal basis for $L^2(G)$. Let P be the orthogonal projection from $L^2(G)$ onto $L^2(E)$. By Lemma 2, $\{P\widehat{T}_\ell \widehat{\psi}_G : \ell \in \mathbb{Z}^d\} = \{\widehat{T}_\ell \widehat{\psi}_E : \ell \in \mathbb{Z}^d\}$ is a normalized tight frame for $L^2(E)$. Therefore for each $k \in \mathbb{Z}$, $\{\widehat{D}_A^k \widehat{T}_\ell \widehat{\psi}_E : \ell \in \mathbb{Z}^d\}$ is a normalized tight frame for $L^2((A')^k E)$. Since Ω is the disjoint union of $\{(A')^k E : k \in \mathbb{Z}\}$, it follows that $L^2(\Omega) = L^2(\mathbb{R}^d) \chi_\Omega$ is the (orthogonal) direct sum of $L^2((A')^k E)$. Thus, by Lemma 3, $\{\widehat{D}_A^k \widehat{T}_\ell \widehat{\psi}_E : \ell \in \mathbb{Z}^d, k \in \mathbb{Z}\}$ is a normalized tight frame for $L^2(\Omega)$, i.e., E is an s -frame wavelet set for X .

“ $\Rightarrow$ ” Assume that E is an s -frame wavelet set for X . By Theorem 1, $\widehat{X} = L^2(\mathbb{R}^d) \cdot \chi_\Omega$ for some measurable set Ω with $\Omega = A'\Omega$. Note that $\widehat{X}$ contains functions $\widehat{D}_A^n \widehat{\psi}_E$, hence it contains $\chi_{(A')^n E}$ for any $n \in \mathbb{Z}$. This implies that $\Omega \supset \bigcup_{n \in \mathbb{Z}} (A')^n E$. On the other hand, any function in $\widehat{X}$ is in the span of $\{\widehat{D}_A^n \widehat{T}_\ell \widehat{\psi}_E : n \in \mathbb{Z}, \ell \in \mathbb{Z}^d\}$, hence it is supported on $\bigcup_{n \in \mathbb{Z}} (A')^n E$. Therefore $\Omega \subset \bigcup_{n \in \mathbb{Z}} (A')^n E$. In order to prove that $E = E(\delta_{A'}, 1)$, we need to prove that (modulo null sets) $\mu((A')^k E \cap (A')^j E) = 0$ for any distinct integers k, j , where μ is the Lebesgue measure in $\mathbb{R}^d$. If this is not true, then there exists an integer $j_0 > 0$ such that $\mu(E \cap (A')^{j_0} E) > 0$. We can then find a subset F of $E \cap (A')^{j_0} E$ with positive measure such that elements in the set $\{(A')^k F : k \in \mathbb{Z}\}$ are mutually disjoint. We can choose F in such a way that it is contained in the cube $\prod_{j=1}^d [2m_j\pi, 2(m_j+1)\pi)$ for some integers m_j with $j \in \{1, 2, \dots, d\}$. Now define f by $\widehat{f} = \chi_F$. $f \in X$

since $F \subset E \subset \Omega$. By assumption, ψ_E is an s -frame wavelet for X . Thus $f = \sum_{n \in \mathbb{Z}, \ell \in \mathbb{Z}^d} \langle f, D_A^n T_\ell \psi_E \rangle D_A^n T_\ell \psi_E$. It follows that

$$\begin{aligned} \|f\|^2 &= \sum_{n \in \mathbb{Z}, \ell \in \mathbb{Z}^d} |\langle f, D_A^n T_\ell \psi_E \rangle|^2 \\ &\geq \sum_{\ell \in \mathbb{Z}^d} |\langle \chi_F, \widehat{T}_\ell \widehat{\psi}_E \rangle|^2 + |\langle \chi_F, \widehat{D}_A^{j_0} \widehat{\psi}_E \rangle|^2 \\ &= \|f\|^2 + \frac{|\det A|^{2j_0}}{(2\pi)^d} \|f\|^2 > \|f\|^2. \end{aligned}$$

This is the contradiction we need.

To prove that $E = E(\tau, 1)$, it suffices to show that $E \cap (E + 2\pi\ell)$ is a null set for any $\ell \in \mathbb{Z}^d \setminus \{O\}$. If there exists $\ell_0 \neq \vec{O}$ in $\mathbb{Z}^d$ such that $E \cap (E + 2\pi\ell_0)$ has positive measure, we can choose a subset J of $E \cap (E + 2\pi\ell_0)$ with positive measure in $\prod_{j=1}^d [2\pi m_j, 2(m_j + 1)\pi)$ for some integers m_j with $j \in \{1, \dots, d\}$. Define g by $\widehat{g} = \chi_J - \chi_{J-2\pi\ell_0}$. $g \in X$ since the support of $\widehat{g}$ is contained in E . We have $\langle \widehat{g}, \widehat{D}_A^n \widehat{T}_\ell \widehat{\psi}_E \rangle = 0$ if $n \neq 0$ (since $E = E(\delta_{A'}, 1)$). Also, for any $\ell \in \mathbb{Z}^d$, $\langle \widehat{g}, \widehat{T}_\ell \widehat{\psi}_E \rangle = \langle \chi_J - \chi_{J-2\pi\ell_0}, \widehat{T}_\ell \widehat{\psi}_E \rangle = \langle \chi_J, \widehat{T}_\ell \widehat{\psi}_E \rangle - \langle \chi_{J-2\pi\ell_0}, \widehat{T}_\ell \widehat{\psi}_E \rangle = 0$, since $\widehat{T}_\ell \widehat{\psi}_E$ is a 2π -periodic function. So we have

$$\widehat{g} = \sum_{n \in \mathbb{Z}, \ell \in \mathbb{Z}^d} \langle \widehat{g}, \widehat{D}_A^n \widehat{T}_\ell \widehat{\psi}_E \rangle \widehat{D}_A^n \widehat{T}_\ell \widehat{\psi}_E = 0.$$

This is again a contradiction. $\square$

3. PATH CONNECTIVITY

In this section we will prove that for a fixed $d \times d$ real expansive matrix A , in a reducing subspace, the set of all s -frame wavelets is not empty and is path-connected in the L^2 -norm.

Theorem 3. *Let A be a $d \times d$ real expansive matrix A and let X be a reducing subspace. Then for any given positive ε , there exists ε_1 , $0 < \varepsilon_1 < \varepsilon$, such that the ring $\mathcal{B}(\varepsilon) \setminus \mathcal{B}(\varepsilon_1)$ contains an s -frame wavelet set for X , where $\mathcal{B}(r)$ stands for the ball centered at the original point with radius r .*

Proof. Without loss of generality, we can assume $\varepsilon < 1$. Let $C = (A'\mathcal{B}(1)) \setminus \mathcal{B}(1)$. Since A is expansive, A' is also expansive. Thus for any $x \in \mathbb{R}^d \setminus \{O\}$, by Lemma 1, we have $\lim_{k \rightarrow +\infty} \|(A')^{-k}x\| = 0$ and $\lim_{k \rightarrow +\infty} \|(A')^kx\| = \infty$. So there is an integer n such that $(A')^n x \notin \mathcal{B}(1)$ and $(A')^{n-1}x \in \mathcal{B}(1)$. It follows that $(A')^n x \in C$, hence $x \in (A')^{-n}C$. Therefore, we have

$$\mathbb{R}^d \setminus \{O\} = \bigcup_{n \in \mathbb{Z}} (A')^n C.$$

Now for any ε with $0 < \varepsilon < 1$, since C is a bounded set and $\lim_{k \rightarrow \infty} \|(A')^{-k}\| = 0$, there is an integer k_0 such that $(A')^{k_0}C \subset \mathcal{B}(\varepsilon)$. Note that 0 is an exterior point of C , hence it is also an exterior point of $(A')^{k_0}C$. So there is an $\varepsilon_1 > 0$ ($\varepsilon_1 < \varepsilon$) such that

$$E = (A')^{k_0}C \subset \mathcal{B}(\varepsilon) \setminus \mathcal{B}(\varepsilon_1) \subset [-\pi, \pi)^d.$$

It is clear that $E = E(\tau, 1)$. Now define $F = \bigcup_{k \geq 1} E^{(1)}(\delta_{A'}, k)$. We see that $F = F(\tau, 1)$, $F = F(\delta_{A'}, 1)$ and $\mathbb{R}^d \setminus \{O\} = \bigcup_{n \in \mathbb{Z}} (A')^n F$.

Since X is a reducing subspace, by Theorem 1, there is a measurable subset Ω of $\mathbb{R}^d$ such that $A'\Omega = \Omega$, and $\widehat{X} = L^2(\mathbb{R}^d) \cdot \chi_\Omega$. If we define $W = F \cap \Omega$, then W is an s -frame wavelet set for X . Indeed, it is clear that $W = W(\tau, 1) = W(\delta_{A'}, 1)$ and that $\{(A')^n W : n \in \mathbb{Z}\}$ is a disjoint family of subsets in Ω . Also from the definition of W and the facts that $\mathbb{R}^d \setminus \{O\} = \bigcup_{n \in \mathbb{Z}} (A')^n F$ and that $A'\Omega = \Omega$, it follows that $\Omega \setminus \{O\} = \bigcup_{n \in \mathbb{Z}} (A')^n W$. $\square$

Theorem 4. *Let A be a $d \times d$ real expansive matrix and let X be a reducing subspace. Then the set of all s -frame wavelet sets for X is path-connected in the L^2 -norm.*

Proof. Since $\lim_{k \rightarrow \infty} \|(A')^{-k}\| = 0$ (Lemma 1), the sequence $\{\|(A')^{-k}\|\}_{k \geq 0}$ is bounded by some $M > 0$. Let $\eta_0 > 0$ be a number less than $\frac{1}{M}$. If we have $\|x\| \geq 1$ and $\|(A')^k x\| < \eta_0$, then we must have $k < 0$ since otherwise we would have $\|(A')^k x\| \geq \frac{\|x\|}{\|(A')^{-k}\|} > \eta_0$. By Theorem 1, there exists a measurable subset Ω of $\mathbb{R}^d$ such that $\widehat{X} = L^2(\mathbb{R}^d) \cdot \chi_\Omega$ and $\Omega = A'\Omega$. By the same argument used in the proof of Theorem 3, there exist a η_1 such that $0 < \eta_1 < \eta_0$ and an s -frame wavelet set G for X such that $G \subset \mathcal{B}(\eta_0) \setminus \mathcal{B}(\eta_1)$. Let E be any s -frame wavelet set for X (so that $E = E(\delta_{A'}, 1)$, $\Omega = \bigcup_{k \in \mathbb{Z}} (A')^k E$ and $E = E(\tau, 1)$ by Theorem 2). It suffices to show that χ_E is path-connected to χ_G in L^2 -norm. For simplicity, we will assume that E and G are disjoint in the following proof. The general case can be obtained by applying the following proof to $E_1 = E \setminus (E \cap G)$ and $G_1 = G \setminus (E \cap G)$ while keeping $E \cap G$ intact.

Let $F_n = (A')^{-n}(G) \cap E$. F_n 's are mutually disjoint and $\bigcup_{n \in \mathbb{Z}} F_n = E$ as one can easily check. $(A')^n(F_n) \subset G$ for each integer n , $(A')^n(F_n)$'s are also disjoint and $\bigcup_{n \in \mathbb{Z}} (A')^n(F_n) = G$. For each integer n , F_n is bounded and the original point is an exterior point of it. Hence there exist real numbers r_n and q_n with $0 < r_n < q_n$, such that $F_n \subset \mathcal{B}(q_n) \setminus \mathcal{B}(r_n)$.

Define

$$\begin{aligned} F_n^s &= \mathcal{B}(r_n + (q_n - r_n)s) \cap F_n, \\ I_0^s &= \bigcup_{n \in \mathbb{Z}} F_n^s. \end{aligned}$$

We have $I_0^0 = \emptyset$ and $I_0^1 = E$. For subsets S , P and Q of $\mathbb{R}^d$, define

$$\begin{aligned} g(S) &= \bigcup_{n \in \mathbb{Z}} (A')^n ((A')^{-n}(G) \cap S) = G \cap \left(\bigcup_{n \in \mathbb{Z}} (A')^n S \right), \\ \tau(P) &= \bigcup_{\ell \in \mathbb{Z}^d} (P - 2\pi\ell), \\ h(P, Q) &= \tau(P) \cap Q. \end{aligned}$$

The following elementary properties of g, h are needed in the proof later. We will prove (v) and (vi) and leave the rest to the reader.

- (i) If $S \subset E$, then $g(S) \subset G$. Moreover, S and $g(S)$ are A' -dilation equivalent.
- (ii) If $\{S_n\}$ is a family of disjoint subsets of E , then $\{g(S_n)\}$ is a family of disjoint subsets of G . Furthermore, $(\bigcup g(S_n)) \cup (E \setminus \bigcup S_n)$ is an A' -dilation generator for Ω .

- (iii) For subsets $P \subset G$ and $Q \subset E$ the set $h(P, Q)$ is a subset of Q which does not intersect $\mathcal{B}(1)$. Furthermore, $\mu(h(P, Q)) \leq \mu(P)$ and $\mu(g(h(P, Q))) \leq \frac{\mu(h(P, Q))}{|\det(A)|} \leq \frac{\mu(P)}{|\det(A)|}$.
- (iv) If $\{P_n\}$ is a family of disjoint subsets of G and Q is a subset of E , then $\{h(P_n, Q)\}$ is a family of disjoint subsets of E .
- (v) The symmetric difference of two sets P and Q , denoted by $P\Delta Q$ is $(P \setminus Q) \cup (Q \setminus P)$. For any $\varepsilon > 0$, there exists $\delta > 0$, such that for any subsets P, Q of E , if $\mu(P\Delta Q) < \delta$, then $\mu(g(P)\Delta g(Q)) < \varepsilon$.
- (vi) For any $\varepsilon > 0$, there exists $\delta > 0$, such that for any subsets C, D of G , and any subsets P, Q of E , if $\mu(C\Delta D) < \delta$ and $\mu(P\Delta Q) < \delta$, then $\mu(h(C, P)\Delta h(D, Q)) < \varepsilon$.

Proof of property (v). Since $g(P)$ and $g(Q)$ are subsets of G and $\mu(G) < \infty$, there exists a natural number N so that

$$\mu\left(\bigcup_{|n| \geq N} (G \cap (A')^n P)\right) < \varepsilon/4 \quad \text{and} \quad \mu\left(\bigcup_{|n| \geq N} (G \cap (A')^n Q)\right) < \varepsilon/4.$$

On the other hand, for each n such that $1 \leq n \leq N$, it is apparent that

$$\mu(G \cap ((A')^n P \Delta (A')^n Q)) < \frac{\varepsilon}{2N}$$

if $\mu(P\Delta Q)$ is small enough, say less than $\delta_n > 0$. Let $\delta = \min\{\delta_1, \delta_2, \dots, \delta_N\}$, the result then follows by the definition of g and the triangle inequality. $\square$

Proof of property (vi). Using the fact that $(A \cap B)\Delta(A' \cap B') \subset (A\Delta A') \cup (B\Delta B')$ for any subsets A, B, A' and B' of $\mathbb{R}^d$, we have

$$\begin{aligned} h(C, P)\Delta h(D, Q) &= ((\tau(C) \cap E) \cap P)\Delta((\tau(D) \cap E) \cap Q) \\ &\subset ((\tau(C)\Delta\tau(D)) \cap E) \cup (P\Delta Q) \\ &= \left(\bigcup_{\ell \in \mathbb{Z}} (C - 2\pi\ell)\Delta\bigcup_{\ell' \in \mathbb{Z}} (D - 2\pi\ell')\right) \cap E \cup (P\Delta Q) \\ &\subset \left(\bigcup_{\ell \in \mathbb{Z}} (C\Delta D - 2\pi\ell) \cap E\right) \cup (P\Delta Q) \\ &= \left(\bigcup_{\ell \in \mathbb{Z}} E_\ell\right) \cup (P\Delta Q), \end{aligned}$$

where $E_\ell = (C\Delta D - 2\pi\ell) \cap E$. Since $\{E_\ell\}$ are disjoint subsets of E and $\mu(E_\ell) < \mu(C\Delta D)$ for each ℓ , it follows that $\mu(h(C, P)\Delta h(D, Q)) < \varepsilon$ if δ is small enough, using an argument similar to the one used in the proof of (v) above. $\square$

Finally, we define

$$\begin{aligned} J_0^s &= g(I_0^s), I_1^s = h(J_0^s, E \setminus I_0^s), \\ &\dots \\ J_n^s &= g(I_n^s), I_{n+1}^s = h(J_n^s, E \setminus \bigcup_{k=0}^n I_k^s), \\ &\dots \\ E^s &= \left(\bigcup_{k=0}^{\infty} J_k^s\right) \cup \left(E \setminus \bigcup_{k=0}^{\infty} I_k^s\right). \end{aligned}$$

It is clear that $E^0 = E$ and $E^1 = G$. $\{I_n^s\}$ is a family of disjoint subsets of E by the construction. It follows that E^s is an A -dilation generator for Ω by property

(ii) above. To see that $E^s = E^s(\tau, 1)$, assume the contrary. Since $G = G(\tau, 1)$ and $E = E(\tau, 1)$, there must exist some $k_0 \geq 0$ so that $J_{k_0}^s$ contains a measurable subset J' of positive measure with the property $J' + 2\pi\ell_0 \subset E \setminus \bigcup_{k=0}^{\infty} I_k^s$ for some $\ell_0 \in \mathbb{Z}^d$. This leads to $J' \subset I_{k_0+1}^s = h(J_{k_0}^s, E \setminus \bigcup_{k=0}^{k_0} I_k^s)$, which is a contradiction. This proves that E^s is an s -frame wavelet set for X for each $s \in [0, 1]$.

Finally, we will show that χ_{E^s} is continuous. Given $\varepsilon > 0$ and $0 \leq s \leq 1$, we need to find $\delta > 0$, such that for each $0 \leq t \leq 1$, if $|t - s| \leq \delta$, then $\|\chi_{E^t} - \chi_{E^s}\|^2 < \varepsilon$. First, by property (iii) above, we have $\mu(I_{k+1}^s) \leq \mu(J_k^s) \leq \frac{\mu(I_k^s)}{|\det(A)|}$ for any s and $k \geq 0$. It follows that $\mu(I_{k+1}^s) \leq \mu(J_k^s) \leq \frac{\mu(E)}{|\det(A)|^{k+1}}$. Since A is expansive, we have $|\det(A)| > 1$. So if k_1 is large enough, we will have

$$\sum_{k=k_1}^{\infty} (\|\chi_{J_k^t} - \chi_{J_k^s}\|^2 + \|\chi_{I_k^t} - \chi_{I_k^s}\|^2) < \frac{\varepsilon}{2}.$$

On the other hand, it is clear that $\chi_{I_0^s}$ is continuous. Property (v) then ensures that $\chi_{J_0^s}$ is continuous and property (vi) in turn guarantees the continuity of $\chi_{I_1^s}$. It follows that each $\chi_{J_k^s}$ and $\chi_{I_k^s}$ are continuous by induction. Hence, there exists $\delta > 0$, such that

$$\sum_{k=0}^{k_1-1} (\|\chi_{J_k^t} - \chi_{J_k^s}\|^2 + \|\chi_{I_k^t} - \chi_{I_k^s}\|^2) < \frac{\varepsilon}{2}$$

if $|s - t| < \delta$. We then have

$$\begin{aligned} \|\chi_{E^t} - \chi_{E^s}\|^2 &\leq \|\chi_{\bigcup_{k=0}^{\infty} J_k^t} - \chi_{\bigcup_{k=0}^{\infty} J_k^s}\|^2 + \|\chi_{E \setminus \bigcup_{k=0}^{\infty} I_k^t} - \chi_{E \setminus \bigcup_{k=0}^{\infty} I_k^s}\|^2 \\ &\leq \sum_{k=0}^{\infty} \|\chi_{J_k^t} - \chi_{J_k^s}\|^2 + \sum_{k=0}^{\infty} \|\chi_{I_k^t} - \chi_{I_k^s}\|^2 < \varepsilon. \end{aligned}$$

□

Remark. In any given reducing subspace X , an s -frame wavelet set E for X is an s -elementary wavelet set if and only if E has measure $(2\pi)^d$. Since we cannot control the measures of the s -frame wavelet sets in our proof, we are not able to recapture D. Speegle's connectivity result in the case when X is $L^2(\mathbb{R}^d)$ itself. On the other hand, our approach does provide a more elementary way of proving the path connectivity property for a larger class of basis functions.

ACKNOWLEDGEMENT

The authors thank the referee for pointing out an error in the proof of Theorem 4 and providing a correction for that.

REFERENCES

- [1] X. Dai, Y. Diao and Q. Gu, *Normalized Tight Frame Wavelet Sets*, Proc. Amer. Math. Soc., to appear.
- [2] X. Dai and D. Larson, *Wandering vectors for unitary systems and orthogonal wavelets*, Memoirs Amer. Math. Soc., 134(1998), no. 640. MR **98m**:47067
- [3] X. Dai, D. Larson and D. Speegle, *Wavelet sets in $\mathbb{R}^n$* , J. Fourier Anal. Appl., 3(1997), 451-456. MR **98m**:42048
- [4] X. Dai and S. Lu, *Wavelets in subspaces*, Mich. J. Math., 43(1996), 81-98. MR **97m**:42021
- [5] N. Dunford and J. Schwartz, *Linear Operators Part I*, Wiley-Interscience, 1958. MR **22**:8302

- [6] D. Han and D. Larson, *Bases, Frames and Group representations*, Memoirs. Amer. Math. Soc., 147(2000). MR **2001a**:47013
- [7] D. Speegle, *The s -elementary wavelets are path-connected*, Proc. Amer. Math. Soc., 127(1999), 223-233. MR **99h**:42045

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF NORTH CAROLINA AT CHARLOTTE, CHARLOTTE, NORTH CAROLINA 28223

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF NORTH CAROLINA AT CHARLOTTE, CHARLOTTE, NORTH CAROLINA 28223

DEPARTMENT OF MATHEMATICS, BEIJING UNIVERSITY, BEIJING, PEOPLE'S REPUBLIC OF CHINA
Current address: Department of Mathematics, East China Normal University, Shanghai, People's Republic of China

DEPARTMENT OF MATHEMATICS AND STATISTICS, MCMASTER UNIVERSITY, HAMILTON, ONTARIO, CANADA L8S 4K1

Current address: Department of Mathematics, University of Central Florida, Orlando, Florida 32816