BERNSTEIN–WALSH INEQUALITIES AND THE EXPONENTIAL CURVE IN $\mathbb{C}^2$

DAN COMAN AND EVGENY A. POLETSKY

(Communicated by Juha M. Heinonen)

ABSTRACT. It is shown that for the pluripolar set $K = \{(z, e^z) : |z| \leq 1\}$ in $\mathbb{C}^2$ there is a global Bernstein–Walsh inequality: If P is a polynomial of degree n on $\mathbb{C}^2$ and $|P| \leq 1$ on K , this inequality gives an upper bound for $|P(z, w)|$ which grows like $\exp(\frac{1}{2}n^2 \log n)$. The result is used to obtain sharp estimates for $|P(z, e^z)|$.

1. INTRODUCTION

If X is a non-pluripolar compact set in $\mathbb{C}^k$ and P is a polynomial of degree n on $\mathbb{C}^k$, the Bernstein–Walsh inequality is (see [K])

$$(1) \quad |P(z)| \leq \|P\|_X e^{nV_X(z)},$$

where $\|P\|_X$ is the uniform norm of P on X and $V_X(z)$ is the extremal function of X . For example, if $z = (z_1, \dots, z_k)$ and $X = \Delta^k = \{z \in \mathbb{C}^k : |z_j| \leq 1, 1 \leq j \leq k\}$ is the unit polydisk, then

$$V_X(z) = L(z) = \max\{\log^+ |z_1|, \dots, \log^+ |z_k|\}.$$

If X is pluripolar, then, in general, such estimates are impossible. For example, if X is any piece of an algebraic curve $\Gamma = \{(z, w) \in \mathbb{C}^2 : P(z, w) = 0\}$, where P is a polynomial, then $\|cP + 1\|_X = 1$ for every $c > 0$ and there are no upper bounds on $cP + 1$.

We consider the case when $\Gamma = \{(z, w) \in \mathbb{C}^2 : w = f(z)\}$ and the compact set

$$K = \{(z, f(z)) \in \mathbb{C}^2 : |z| \leq 1\},$$

where f is an entire transcendental function. Then any non-trivial polynomial is not identically equal to 0 on K . Therefore a compactness argument shows that, for every n , there is a number $c_n > 0$ such that for any polynomial $P(z, w)$ of degree at most n the norm $\|P\|_{\Delta^2} \leq c_n \|P\|_K$. Hence for every $(z, w) \in \mathbb{C}^2$

$$(2) \quad |P(z, w)| \leq \|P\|_K E_n(f) e^{nL(z, w)},$$

where $E_n(f)$ is the least value of c_n . (See also Section 2.)

Inequality (2) can be viewed as a transcendental global version of the Bernstein–Walsh inequality (1), provided that one can obtain good estimates for $E_n(f)$. Moreover, the numbers $E_n(f)$ can serve as a measure of transcendency of f : A “less

Received by the editors June 8, 2001 and, in revised form, October 18, 2001.

2000 *Mathematics Subject Classification*. Primary 41A17; Secondary 30D15, 30D20.

The second author was partially supported by NSF Grant DMS-9804755.

transcendental" function f has larger numbers $E_n(f)$. Note that if f was algebraic, hence a polynomial of degree l , then $E_n(f) = +\infty$ for every $n \geq l$.

In this paper we study the classical case of $f(z) = e^z$ and we let $E_n = E_n(e^z)$. For this function we prove the following global Bernstein–Walsh inequality:

Theorem 1.1. *If $f(z) = e^z$, then there exists a constant $C_1 > 0$ so that*

$$\exp\left(\frac{n^2 \log n}{2} - C_1 n^2\right) \leq E_n \leq \exp\left(\frac{n^2 \log n}{2} + C_1 n^2\right),$$

for all $n \geq 1$. If P is a polynomial of degree n on $\mathbb{C}^2$, then

$$(3) \quad |P(z, w)| \leq \|P\|_K \exp\left(\frac{n^2 \log n}{2} + C_1 n^2 + nL(z, w)\right).$$

Thus, despite the pluripolarity of K , there is an upper estimate for the absolute value of polynomials, which grows asymptotically as $\exp(n^2 \log n)$. This is not much worse than $\exp(n)$ in the classical Bernstein–Walsh inequality (1). Moreover, this estimate is asymptotically sharp.

Inequality (3) improves when $(z, w) \in \Gamma$. In [T] (see also [B]) it was proved that

$$|P(z, e^z)| \leq \|P\|_K e^{n^2 \log^+ |z| + 6n|z|}.$$

This inequality was used to prove deep theorems concerning the algebraic independence of values of e^z .

For a general transcendental function f we introduce the function

$$m_n(r) = \sup\{\log |P(z, f(z))| : \deg P \leq n, \|P\|_K \leq 1, |z| \leq r\}.$$

The numbers $m_n(r)$ can also serve as a measure of transcendency of f . Let $(|z| - 1)^+ = \max\{|z| - 1, 0\}$. As a consequence of Theorem 1.1, we prove the following restricted Bernstein–Walsh inequality for $f(z) = e^z$:

Theorem 1.2. *There exists an absolute constant $C_2 > 0$ such that for every polynomial P of degree $n \geq 1$ on $\mathbb{C}^2$ and every $z \in \mathbb{C}$ we have*

$$|P(z, e^z)| \leq \|P\|_K \exp\left[\frac{n^2}{2} \left(\log^+ |z| + C_2 \frac{(|z| - 1)^+}{1 + \log n}\right)\right].$$

Moreover $\lim_{n \rightarrow \infty} m_n(r)/n^2 = \frac{1}{2} \log r$, locally uniformly for $r \geq 1$.

This theorem provides the exact asymptotic behavior of the functions $m_n(r)$. It also improves Tijdeman's estimate if one fixes z and lets $n \rightarrow \infty$. On the other hand, if one fixes n and lets $z \rightarrow \infty$, then Tijdeman's estimate is better (at least if n is large).

There is a fundamental difference between classical and transcendental Bernstein–Walsh inequalities. In the classical case (1) the extremal function V_X is given by (see [K])

$$V_X(z, w) = \limsup_{n \rightarrow \infty} \frac{1}{n} \sup\{\log^+ |P(z, w)| : \deg P = n, \|P\|_X \leq 1\}.$$

In the transcendental case (3) it follows from Theorem 1.1 and the Hartogs lemma that

$$\limsup_{n \rightarrow \infty} \frac{2}{n^2 \log n} \sup\{\log^+ |P(z, w)| : \deg P = n, \|P\|_K \leq 1\} = 1$$

everywhere on $\mathbb{C}^2$ except a pluripolar set. Moreover, by Theorem 1.2,

$$\limsup_{n \rightarrow \infty} \frac{2}{n^2 \log n} \sup\{\log^+ |P(z, e^z)| : \deg P = n, \|P\|_K \leq 1\} = 0.$$

The next proposition holds for all entire transcendental functions.

Proposition 1.3. *If f is an entire transcendental function, then*

$$(4) \quad m_n(r) \geq \frac{n^2 + 3n}{2} \log r$$

for every $r \geq 1$. Moreover for $r \geq 1$

$$(5) \quad E_n(f) \geq \exp\left(\frac{n^2 + 3n}{2} \log r - nL(r, M_f(r))\right).$$

If f is of finite order of growth $< \rho$, or of finite order ρ and finite type, then $E_n(f) \geq \exp\left(\frac{n^2 \log n}{2\rho} - Cn^2\right)$ for all $n \geq 1$, where $C = C(f) > 0$.

Proposition 1.3 and the previous theorem imply that the function e^z provides asymptotically the smallest possible functions $m_n(r)$.

We are grateful to Norm Levenberg for the introduction to the problem and discussions.

2. PRELIMINARIES

We use the following notation. If g is an entire holomorphic function we let

$$M_g(r) = \max\{|g(z)| : |z| = r\}.$$

For $n \geq 0$ we denote by $\mathcal{P}_n$ the space of polynomials $P \in \mathbb{C}[z, w]$ of degree at most n . Then $\dim \mathcal{P}_n = (n+1)(n+2)/2 = N+1$, where $N = (n^2 + 3n)/2$.

Let f be an entire transcendental function. For any polynomial $P \in \mathcal{P}_n$ we denote by $P_\star$ the entire function

$$P_\star(z) = P(z, f(z)), \quad z \in \mathbb{C},$$

so $\|P\|_K = M_{P_\star}(1)$. Since f is transcendental, it follows that $\|\cdot\|_K$ is a norm on each vector space $\mathcal{P}_n$. As $\mathcal{P}_n$ are finite dimensional we have

$$E_n(f) = \sup\{\|P\|_{\Delta^2} : P \in \mathcal{P}_n, \|P\|_K \leq 1\} < +\infty,$$

for each $n \geq 0$. Note that $E_0(f) = 1$ and $E_n(f) \leq E_{n+1}(f)$.

Inequality (2) implies that the function

$$u_n(z) = \sup\{\log |P_\star(z)| : P \in \mathcal{P}_n, \|P\|_K \leq 1\}$$

is well defined. It is easy to see by a normal family argument that u_n is a non-negative continuous subharmonic function on $\mathbb{C}$ and $u_n = 0$ on Δ . We have $m_n(r) = \max\{u_n(z) : |z| \leq r\}$, hence $m_n(r)$ is a continuous increasing convex function of $\log r$.

We need the following simple lemma:

Lemma 2.1. *The following inequalities hold for every integer $m > 0$:*

$$\begin{aligned}\log(m+1) &\leq \sum_{j=1}^m \frac{1}{j} \leq \log m + 1, \\ m \log m - m + 1 &\leq \sum_{j=1}^m \log j \leq (m+1) \log m - m + 1, \\ \frac{m^2 \log m}{2} - \frac{m^2}{4} + \frac{1}{4} &\leq \sum_{j=1}^m j \log j \leq \frac{m^2 \log m}{2} - \frac{m^2}{4} + m \log m + \frac{1}{4}.\end{aligned}$$

Proof. The proof is elementary. For instance, the third inequality follows using $\int_{j-1}^j x \log x \, dx \leq j \log j \leq \int_j^{j+1} x \log x \, dx$. $\square$

3. PROOFS

We first prove Proposition 1.3, which was stated for arbitrary entire transcendental functions f . Recall the notations $N = (n^2 + 3n)/2$ and $P_\star(z) = P(z, f(z))$.

Proof of Proposition 1.3. Since $\dim \mathcal{P}_n = N + 1$, there exists $P \in \mathcal{P}_n$, $P \not\equiv 0$, such that the vanishing order of $P_\star$ at 0 is at least N . We let $g(z) = P_\star(z)/z^N$, so

$$M_{P_\star}(1) = M_g(1) \leq M_g(r) = M_{P_\star}(r)/r^N,$$

provided that $r \geq 1$. This and the definition of $m_n(r)$ clearly imply (4). Using (2) with $w = f(z)$ and $|z| \leq r$ we get

$$r^N \leq M_{P_\star}(r)/M_{P_\star}(1) \leq E_n(f) \exp[nL(r, M_f(r))],$$

so (5) follows.

In the case when f is of finite order of growth, we have $\log^+ M_f(r) \leq Cr^\rho$ for every $r \geq 1$. The conclusion follows by taking $r = n^{1/\rho}$ in (5) and by using the above estimate on $\log^+ M_f(r)$. $\square$

In the following proofs we have $f(z) = e^z$, so $P_\star(z) = P(z, e^z)$.

Proof of Theorem 1.1. The lower estimate of E_n follows from Proposition 1.3, since $f(z) = e^z$ has order of growth $\rho = 1$ and type 1 with respect to this order. Moreover, (3) follows from (2) and the upper bound of E_n . To prove the upper bound, we introduce the following notation. Let $d = \frac{d}{dz}$. For any polynomial $R(\lambda) = \sum_{j=0}^m c_j \lambda^j$ we denote by D_R the constant-coefficient differential operator

$$D_R = R(d) = \sum_{j=0}^m c_j \frac{d^j}{dz^j}.$$

Then for any integer $t \geq 0$ and any $\alpha \in \mathbb{C}$ we have

$$(6) \quad D_R[z^t e^{\alpha z}] \Big|_{z=0} = \sum_{j \geq t} c_j \frac{j!}{(j-t)!} \alpha^{j-t} = \frac{d^t R}{d\lambda^t} \Big|_{\lambda=\alpha} = R^{(t)}(\alpha).$$

Now fix $P \in \mathcal{P}_n$, $n \geq 1$, with $\|P\|_K \leq 1$. By Cauchy's estimates we have

$$(7) \quad |P_\star^{(l)}(0)| \leq l!, \quad \forall l \geq 0.$$

We write

$$(8) \quad P(z, w) = \sum_{k=0}^n P_{n-k}(z)w^k, \quad P_{n-k}(z) = \sum_{j=0}^{n-k} c_{kj}z^j.$$

In the sequel, we denote by C all absolute constants involved in our estimates. (They may change from one inequality to the next.)

Proposition 3.1. *There exists a constant C such that*

$$M_{P_{n-k}}(1) \leq \exp\left(\frac{n^2 \log n}{2} + Cn^2\right), \quad \forall k \in \{0, \dots, n\}.$$

Note that this proposition implies Theorem 1.1. Indeed,

$$\|P\|_{\Delta^2} \leq \sum_{k=0}^n M_{P_{n-k}}(1) \leq \exp\left(\frac{n^2 \log n}{2} + Cn^2\right),$$

so the same estimate holds for E_n since P is arbitrary with $\|P\|_K \leq 1$.

In order to prove Proposition 3.1, we fix $k \in \{0, \dots, n\}$. We will estimate the coefficients c_{kj} of P_{n-k} by using the differential operators given by the polynomials

$$(9) \quad \begin{aligned} R_k(\lambda) &= \prod_{l=0, l \neq k}^n (\lambda - l)^{n-l+1}, \\ R_{kj}(\lambda) &= R_k(\lambda)(\lambda - k)^j, \quad j = 0, \dots, n - k. \end{aligned}$$

Note that $\deg R_{k,n-k} = N$. By (6) we have

$$\begin{aligned} \alpha_{kj} &:= D_{R_{kj}} P_{\star}(z) \Big|_{z=0} = (D_{R_k}(d - k)^j) \left[\sum_{l=0}^{n-k} c_{kl} z^l e^{kz} \right] \Big|_{z=0} \\ &= D_{R_k} \left[\sum_{l=j}^{n-k} c_{kl} \frac{l!}{(l-j)!} z^{l-j} e^{kz} \right] \Big|_{z=0} \\ &= \sum_{l=j}^{n-k} c_{kl} \frac{l!}{(l-j)!} R_k^{(l-j)}(k). \end{aligned}$$

We write

$$(10) \quad c'_{kl} = l! c_{kl}, \quad l = 0, \dots, n - k,$$

$$(11) \quad r_{kt} = R_k^{(t)}(k)/t!, \quad t = 0, \dots, n - k.$$

Then c'_{kl} are the unique solution of the triangular system

$$\sum_{l=j}^{n-k} r_{k,l-j} c'_{kl} = \alpha_{kj}, \quad j = 0, \dots, n - k,$$

which yields

$$(12) \quad c'_{k,n-k-j} = \frac{\alpha_{k,n-k-j}}{r_{k0}} - \sum_{l=1}^j \frac{r_{kl}}{r_{k0}} c'_{k,n-k-j+l}, \quad j = 0, \dots, n - k.$$

In order to estimate the coefficients c'_{kj} , we obtain first bounds for α_{kj} , r_{k0} and r_{kl}/r_{k0} . This is done in a sequence of lemmas.

Lemma 3.2. *For all $k = 0, \dots, n$ and $j = 0, \dots, n - k$ we have*

$$|\alpha_{kj}| \leq e^{n^2 \log n + Cn \log n},$$

where $C > 0$ is an absolute constant.

Proof. We write $R_{kj}(\lambda) = \sum_{l=0}^{N_j} s_l \lambda^l$, $N_j = \deg R_{kj} = N - (n - k - j)$, and define

$$|R_{kj}|(\lambda) = \sum_{l=0}^{N_j} |s_l| \lambda^l = (\lambda + k)^j \prod_{l=0, l \neq k}^n (\lambda + l)^{n-l+1}.$$

Using Cauchy's estimates (7) and Stirling's formula $l! \leq e(l/e)^l \sqrt{l}$, for $l \geq 1$ (see [R]), we get

$$\begin{aligned} |\alpha_{kj}| &\leq \sum_{l=0}^{N_j} |s_l| l! \leq e \sum_{l=0}^{N_j} |s_l| (l/e)^l \sqrt{l} \\ &\leq e\sqrt{N} \sum_{l=0}^{N_j} |s_l| N^l = e\sqrt{N} |R_{kj}|(N) \\ &\leq e\sqrt{N} \prod_{l=0}^n (N + l)^{n-l+1} \leq e\sqrt{N} \exp \left((\log(N + n)) \sum_{l=1}^{n+1} l \right). \end{aligned}$$

Since $\log(N + n) \leq 2 \log n + 2/n$, this yields $|\alpha_{kj}| \leq \exp(n^2 \log n + Cn \log n)$. $\square$

Lemma 3.3. *If $0 \leq k \leq n$, then $|r_{k0}| \geq \exp[(n^2 \log n)/2 - Cn^2]$, where $C > 0$ is an absolute constant.*

Proof. We have by (9) and (11) that

$$|r_{k0}| = |R_k(k)| = \prod_{l=0, l \neq k}^n |k - l|^{n-l+1},$$

so after a direct calculation we get

$$\log |r_{k0}| = (n - k + 1) \left(\sum_{l=1}^k \log l + \sum_{l=1}^{n-k} \log l \right) + \sum_{l=1}^k l \log l - \sum_{l=1}^{n-k} l \log l.$$

Using Lemma 2.1, with the convention that $x \log x = 0$ for $x = 0$, it follows that for every $k = 0, \dots, n$ we have

$$\begin{aligned} \log |r_{k0}| &\geq (n - k + 1)[k \log k + (n - k) \log(n - k) - n] \\ &\quad + \frac{k^2 \log k}{2} - \frac{k^2}{4} - \frac{(n - k)^2 \log(n - k)}{2} \\ &\quad + \frac{(n - k)^2}{4} - (n - k) \log(n - k) \\ &\geq k \left(n - \frac{k}{2} \right) \log k + \frac{(n - k)^2}{2} \log(n - k) - Cn^2 \\ &= \frac{n^2 \log n}{2} + n^2 F \left(\frac{k}{n} \right) - Cn^2 \geq \frac{n^2 \log n}{2} - Cn^2, \end{aligned}$$

where $F(x) = x(1 - x/2) \log x + 0.5(1 - x)^2 \log(1 - x) > -C$ for all $x \in [0, 1]$ and some constant $C > 0$. $\square$

Lemma 3.4. *For any $k = 0, \dots, n$ and $j = 0, \dots, n - k$ we have*

$$|r_{kj}|/|r_{k0}| \leq e^2 n^2 (n+1)^j.$$

Proof. We fix $\epsilon \in (0, 1)$, to be chosen later in terms of n . Using the definition (11) of r_{kj} and applying Cauchy's estimates to R_k , we obtain that $|r_{kj}| \leq M/\epsilon^j$, where $M = \max\{|R_k(\lambda)| : |\lambda - k| = \epsilon\}$. If λ is on the circle $|\lambda - k| = \epsilon$, then

$$|R_k(\lambda)| \leq \prod_{l=0}^{k-1} (k + \epsilon - l)^{n-l+1} \prod_{l=k+1}^n (l - k + \epsilon)^{n-l+1} = |r_{k0}| F_k(\epsilon),$$

where

$$F_k(\epsilon) = \prod_{l=0}^{k-1} \left(1 + \frac{\epsilon}{k-l}\right)^{n-l+1} \prod_{l=k+1}^n \left(1 + \frac{\epsilon}{l-k}\right)^{n-l+1}.$$

Therefore $|r_{kj}|/|r_{k0}| \leq F_k(\epsilon)/\epsilon^j$. Using Lemma 2.1 we get

$$\begin{aligned} \log F_k(\epsilon) &\leq 2(n+1) \sum_{l=1}^n \log \left(1 + \frac{\epsilon}{l}\right) \\ &\leq 2\epsilon(n+1) \sum_{l=1}^n \frac{1}{l} \leq 2\epsilon(n+1)(\log n + 1). \end{aligned}$$

Choosing $\epsilon = 1/(n+1)$ we obtain $\log F_k(\epsilon) \leq 2(\log n + 1)$, hence the lemma follows. $\square$

We can now estimate the coefficients c'_{kj} by using the system of equations (12). Lemmas 3.2 and 3.3 imply that

$$(13) \quad \frac{|\alpha_{kj}|}{|r_{k0}|} \leq e^{t_n}, \quad t_n := \frac{n^2 \log n}{2} + Cn^2$$

holds for every $k = 0, \dots, n$ and $j = 0, \dots, n - k$, with an absolute constant $C > 0$. For fixed k , we prove by induction on $j = n - k, \dots, 0$ that

$$(14) \quad |c'_{kj}| \leq e^{t_n} [(n+1)(1 + e^2 n^2)]^{n-k-j}.$$

If $j = n - k$ this holds by (12) and (13). Assuming that the inequality is true for $l = n - k, \dots, n - k - j + 1$, we obtain by using Lemmas 3.2, 3.3 and 3.4 that

$$\begin{aligned} |c'_{k,n-k-j}| &\leq \frac{|\alpha_{k,n-k-j}|}{|r_{k0}|} + \sum_{l=1}^j \frac{|r_{kl}|}{|r_{k0}|} |c'_{k,n-k-j+l}| \\ &\leq e^{t_n} \left[1 + \sum_{l=1}^j e^2 n^2 (n+1)^l [(n+1)(1 + e^2 n^2)]^{j-l} \right] \\ &\leq e^{t_n} (n+1)^j \left[1 + e^2 n^2 \sum_{l=1}^j (1 + e^2 n^2)^{j-l} \right] \\ &= e^{t_n} (n+1)^j (1 + e^2 n^2)^j, \end{aligned}$$

so (14) is proved. Therefore, combining (10), (13) and (14), we get for every $j = 0, \dots, n - k$ that

$$|c_{kj}| \leq |c'_{kj}| \leq e^{t_n} (n+1)^n (1 + e^2 n^2)^n \leq \exp \left(\frac{n^2 \log n}{2} + Cn^2 \right),$$

with an absolute constant $C > 0$. Using this and (8) it follows that

$$M_{P_{n-k}}(1) \leq \sum_{j=0}^{n-k} |c_{kj}| \leq \exp\left(\frac{n^2 \log n}{2} + Cn^2\right).$$

The proof of Proposition 3.1, and hence the proof of Theorem 1.1, are complete. $\square$

Remark. Using the main steps and ideas from this proof, together with more careful estimates of the constants involved, we can prove for every $n \geq 1$ the inequalities

$$\exp\left(\frac{n^2 \log n}{2} - n^2\right) < E_n < \exp\left(\frac{n^2 \log n}{2} + n^2 + 63\right).$$

Proof of Theorem 1.2. Let x_n be defined as the unique solution of the equation $n(x_n - 1)e^{x_n} = \log E_n$, for $n \geq 1$. Using the polynomial $P(z, w) = w - 1 - z$ one checks that $E_1 > 1$. Therefore $x_n > 1$ and by Theorem 1.1 $\lim_{n \rightarrow \infty} x_n = +\infty$.

Let $\alpha_n = (\log E_n)/x_n$. We estimate first x_n and α_n . By Theorem 1.1

$$(x_n - 1)e^{x_n} = \frac{\log E_n}{n} = \frac{n \log n}{2} + O(n),$$

hence $x_n + \log(x_n - 1) = \log n - \log 2 + \log(\log n + O(1))$. This implies $\lim_{n \rightarrow \infty} x_n / \log n = 1$, so $\lim_{n \rightarrow \infty} 2\alpha_n/n^2 = 1$. Since

$$x_n = \log n - \log 2 + \log\left(\frac{\log n + O(1)}{x_n - 1}\right)$$

and since $x_n / \log n \rightarrow 1$ we get $x_n = \log n + O(1)$ and

$$(15) \quad \frac{2\alpha_n}{n^2} - 1 = \frac{2 \log E_n}{n^2 x_n} - 1 = \frac{\log n + O(1)}{\log n + O(1)} - 1 = \frac{O(1)}{\log n}.$$

We can now prove the estimate in the statement. Let $P \in \mathcal{P}_n$ with $\|P\|_K = 1$ and consider the subharmonic function $u(z) = \log^+ |P_\star(z)|$. Then $u \leq 0$ on Δ and $u(z) \leq \log E_n + n|z|$, for every $z \in \mathbb{C}$, by (2). Let $\tilde{u}(x) = \max\{u(z) : |z| = e^x\}$, where $x \geq 0$. Then $\tilde{u}$ is a convex increasing function, $\tilde{u}(0) = 0$ and

$$\tilde{u}(x) \leq \phi(x) = \log E_n + ne^x, \quad \forall x \geq 0.$$

Note that x_n verifies $x_n \phi'(x_n) = \phi(x_n)$, which means that the tangent line to the graph of ϕ at $(x_n, \phi(x_n))$ passes through the origin. Using the convexity of $\tilde{u}$ it follows that

$$\tilde{u}(x) \leq \begin{cases} \phi'(x_n)x, & \text{if } 0 \leq x \leq x_n, \\ \phi(x), & \text{if } x \geq x_n. \end{cases}$$

Since $x \leq e^x - 1$ and $ne^{x_n} = (\log E_n)/(x_n - 1)$ we obtain

$$\begin{aligned} \phi'(x_n)x &= (\log E_n + ne^{x_n}) \frac{x}{x_n} \\ &\leq \alpha_n x + \frac{ne^{x_n}}{x_n} (e^x - 1) = \alpha_n \left(x + \frac{e^x - 1}{x_n - 1}\right), \end{aligned}$$

for all $x \geq 0$. Moreover, it is easy to see that

$$\phi(x) = \log E_n + ne^x \leq \alpha_n x + \frac{ne^{x_n}}{x_n} (e^x - 1)$$

holds for $x \geq x_n$. We conclude that for all $x \geq 0$ we have

$$\tilde{u}(x) \leq \alpha_n \left(x + \frac{e^x - 1}{x_n - 1}\right) = \frac{n^2}{2} \left[x + \left(\frac{2\alpha_n}{n^2} - 1\right)x + \frac{2\alpha_n}{n^2} \frac{e^x - 1}{x_n - 1}\right].$$

Using (15) and the asymptotics of x_n we obtain

$$\tilde{u}(x) \leq \frac{n^2}{2} \left(x + \frac{Cx}{\log n} + C' \frac{e^x - 1}{x_n} \right) \leq \frac{n^2}{2} \left(x + C_2 \frac{e^x - 1}{1 + \log n} \right),$$

where $C_2 > 0$ is an absolute constant. If $x = \log |z|$, $|z| \geq 1$, this gives the desired inequality.

Combining the inequality we have just proved with (4), we get for all $n \geq 1$ and $r \geq 1$ that

$$\frac{\log r}{2} + \frac{3 \log r}{2n} \leq \frac{m_n(r)}{n^2} \leq \frac{\log r}{2} + \frac{C_2(r-1)}{2(1+\log n)}.$$

Therefore $\lim_{n \rightarrow \infty} m_n(r)/n^2 = \frac{1}{2} \log r$, locally uniformly for $r \geq 1$. □

REFERENCES

- [B] A. Baker, *Transcendental Number Theory*, Cambridge Univ. Press, 1975. MR **54**:10163
- [K] M. Klimek, *Pluripotential theory*, Clarendon Press, Oxford, 1991. MR **93h**:32021
- [R] W. Rudin, *Principles of Mathematical Analysis*, McGraw-Hill, Inc., 3rd ed., 1976. MR **52**:5893
- [T] R. Tijdeman, *On the number of zeros of general exponential polynomials*, Indag. Math., 37(1971), 1-7. MR **44**:4193

DEPARTMENT OF MATHEMATICS, SYRACUSE UNIVERSITY, SYRACUSE, NEW YORK 13244-1150
E-mail address: dcoman@syr.edu

DEPARTMENT OF MATHEMATICS, SYRACUSE UNIVERSITY, SYRACUSE, NEW YORK 13244-1150
E-mail address: eapolets@syr.edu