REGULARIZATION OF A_p WEIGHTS

RICHARD J. BAGBY AND BASEM MASAEDDEH

(Communicated by Andreas Seeger)

ABSTRACT. We show how to approximate a given weight function in the class A_p by weights that are bounded above by multiples of their infima in such a way that the A_p constant is not increased. As an application, we show that the precise range of p for which a given weight is in A_p cannot be determined by extrapolating the A_p constants.

1. INTRODUCTION

For $1 < p < \infty$, the class $A_p(\mathbf{R}^n)$ consists of those locally integrable weight functions w that are positive almost everywhere and satisfy an *a priori* bound of the form

$$(1.1) \quad \left(\int_Q w(x) \, dx \right)^{1/p} \left(\int_Q w(x)^{1-p'} \, dx \right)^{1/p'} \leq A |Q|$$

for all cubes Q in $\mathbf{R}^n$; here p' is the Hölder conjugate of p , A is a constant depending on w and $|Q|$ is the Lebesgue measure of Q . The smallest value of A for which (1.1) can hold is called $A_p(w)$, the A_p -constant for w . These classes were introduced by B. Muckenhoupt [5] in connection with weighted inequalities for the Hardy-Littlewood maximal function, and were soon seen to have a number of remarkable properties. Note that Hölder's inequality shows

$$|Q| = \int_Q w(x)^{1/p} w(x)^{-1/p'} \, dx \leq \left(\int_Q w(x) \, dx \right)^{1/p} \left(\int_Q w(x)^{1-p'} \, dx \right)^{1/p'};$$

for that reason a condition such as (1.1) is often referred to as a reverse Hölder inequality. A simple application of Hölder's inequality shows that $A_q(w) \leq A_p(w)$ whenever $p < q < \infty$, and several delicate arguments have been devised to show that, when $w \in A_p$, $A_q(w)$ must always be finite for all q in an open interval (p_0, ∞) that includes p . These arguments give upper bounds for p_0 and for $A_q(w)$ in terms of p, q and $A_p(w)$; see Muckenhoupt [5], Coifman and Fefferman [3], and Chung, Hunt, and Kurtz [2] as well as the related work by Gehring [4].

In [1], the authors showed how to derive Orlicz space bounds that substitute for $A_{p_0}(w)$ when p_0 is determined by such arguments, but left open the question of whether the best value of p_0 could be obtained by such methods. That is, must $A_{p_0}(w) = \infty$ for some weight having the given A_p -constant? Our results here do

Received by the editors September 12, 2001.

2000 *Mathematics Subject Classification*. Primary 42B25.

not answer that question, but do show that we can never determine $w \notin A_{p_0}$ by considering only values of $A_p(w)$ for finitely many $p > p_0$, each having $A_p(w) < \infty$.

Our investigations here develop a regularization principle that can be useful for establishing *a priori* bounds for classes of the form $\{w \in A_p : A_p(w) \leq M\}$. Given a locally integrable weight function w , we show how to construct a family of weights $\{w_t : t > 0\}$ such that each w_t and its reciprocal is uniformly bounded, with $A_p(w_t) \leq A_p(w)$ and $A_p(w) = \lim_{t \rightarrow 0+} A_p(w_t)$. By working with w_t in place of w , we can improve some arguments by guaranteeing that all the integrals involved are finite.

2. REGULARIZATION OF A_p WEIGHTS

Given a locally integrable function w on $\mathbf{R}^n$ that is positive almost everywhere, define

$$A_p(w) = \sup_Q \frac{1}{|Q|} \left(\int_Q w \right)^{1/p} \left(\int_Q w^{1-p'} \right)^{1/p'}$$

so that $w \in A_p$ if and only if $A_p(w) < \infty$. While $\int_Q w^{1-p'}$ need not be finite for all cubes, $\int_Q w$ is always positive so this supremum is well-defined, either as positive and real or as ∞ . For $t > 0$ we define

$$w_t(x) = \frac{1}{t + 1/(t + w(x))} = \frac{t + w(x)}{t^2 + tw(x) + 1};$$

the first form gives us $w_t(x) = 1/t$ when $w(x) = \infty$. Then $\lim_{t \rightarrow 0+} w_t(x) = w(x)$ pointwise, and $t/(t^2 + 1) \leq w_t(x) \leq 1/t$. Note that both w_t and $1/w_t$ are in Muckenhoupt's class A_1 , since that class consists of all weights for which the average over each cube is dominated by a fixed multiple of the essential infimum over the same cube. For $1 < p < \infty$ we easily calculate

$$A_p(w_t) \leq \left(\frac{1}{t} \right)^{1/p} \left[\left(\frac{t}{t^2 + 1} \right)^{-p'/p} \right]^{1/p'} = \left(\frac{t^2 + 1}{t^2} \right)^{1/p}$$

and find exactly the same bound for $A_p(1/w_t)$. Of course, these are only crude bounds; the theorem below shows that $A_p(w_t)$ mimics the behavior of $A_p(w)$ for small t .

Theorem 1. *For each a.e. positive locally integrable function w on $\mathbf{R}^n$ and for $1 < p < \infty$, $A_p(w_t) \leq A_p(w)$ on $(0, \infty)$, with $A_p(w) = \lim_{t \rightarrow 0+} A_p(w_t)$.*

Proof. For $s, t \geq 0$ call $u(x) = s + w(x)$ and $v(x) = tu(x) + 1$, and then define

$$F(s, t) = \left(\int_Q \frac{u}{v} \right)^{p'-1} \int_Q \left(\frac{u}{v} \right)^{1-p'}$$

with Q a fixed but arbitrary cube in $\mathbf{R}^n$. We begin by showing that the partial derivatives $\frac{\partial F}{\partial s}(s, 0)$ and $\frac{\partial F}{\partial t}(s, t)$ are never positive. The bound for $\frac{\partial F}{\partial s}$ was done earlier by the authors in [1]; we reprove it here with a slightly different argument.

Since

$$F(s, 0) = \left(\int_Q (s + w) \right)^{p'-1} \int_Q (s + w)^{1-p'},$$

for $s > 0$ we can differentiate with respect to s to find

$$\begin{aligned}\frac{\partial F}{\partial s}(s, 0) &= (p' - 1) \left(\int_Q (s + w) \right)^{p'-2} |Q| \int_Q (s + w)^{1-p'} \\ &\quad + (1 - p') \left(\int_Q (s + w) \right)^{p'-1} \int_Q (s + w)^{-p'} \\ &= (p' - 1) \left(\int_Q u \right)^{p'-2} \left[|Q| \int_Q u^{1-p'} - \left(\int_Q u \right) \int_Q u^{-p'} \right].\end{aligned}$$

We use a simple corollary of Hölder's inequality to show that this is never positive: for $u > 0$ and $0 \leq \theta \leq 1$ we have

$$\left(\int u^{a(1-\theta)+b\theta} \right) \left(\int u^{a\theta+b(1-\theta)} \right) \leq \left(\int u^a \right) \left(\int u^b \right).$$

Note $|Q| = \int_Q u^0$, and calling $\theta = 1/(p' + 1)$ lets us write

$$0 = 1 - \theta - p'\theta \quad \text{and} \quad 1 - p' = \theta - p'(1 - \theta).$$

Thus

$$|Q| \int_Q u^{1-p'} \leq \left(\int_Q u \right) \int_Q u^{-p'} < \infty,$$

proving that $\frac{\partial F}{\partial s}(s, 0) \leq 0$ for each $s > 0$.

Obviously $\lim_{s \rightarrow 0+} \int_Q (s + w) = \int_Q w$, and the monotone convergence theorem shows $\lim_{s \rightarrow 0+} \int_Q (s + w)^{1-p'} = \int_Q w^{1-p'}$ whether this last quantity is finite or infinite, so we have established that $F(s, 0) \leq F(0, 0)$.

Next we fix $s > 0$ and differentiate with respect to t . Since

$$\frac{\partial}{\partial t} \left(\frac{u}{v} \right) = \frac{\partial}{\partial t} \left(\frac{u}{tu + 1} \right) = -\frac{u^2}{v^2},$$

we find

$$\frac{\partial F}{\partial t}(s, t) = (p' - 1) \left(\int_Q \frac{u}{v} \right)^{p'-2} \left[\left(\int_Q \frac{u}{v} \right) \int_Q \left(\frac{u}{v} \right)^{2-p'} - \left(\int_Q \frac{u^2}{v^2} \right) \int_Q \left(\frac{u}{v} \right)^{1-p'} \right].$$

Again calling $\theta = 1/(p' + 1)$, we may write

$$1 = 2(1 - \theta) + \theta(1 - p') \quad \text{and} \quad 2 - p' = 2\theta + (1 - \theta)(1 - p')$$

and then show that $\frac{\partial F}{\partial t}(s, t) \leq 0$ as above; note all powers of u/v are locally integrable.

As for $F(s, t)$ itself, we have

$$\lim_{t \rightarrow 0+} \int_Q u/v = \int_Q u \quad \text{and} \quad \lim_{t \rightarrow 0+} \int_Q (u/v)^{1-p'} = \int_Q u^{1-p'}$$

by monotone convergence in the first case and by uniform convergence in the second, so that $\lim_{t \rightarrow 0+} F(s, t) = F(s, 0) \leq F(0, 0)$. Hence we have $F(s, t) \leq F(0, 0)$ for all $s, t > 0$. In particular, this holds true for $s = t$, and after taking p' roots we obtain

$$\left(\int_Q w_t \right)^{1/p} \left(\int_Q w_t^{1-p'} \right)^{1/p'} \leq \left(\int_Q w \right)^{1/p} \left(\int_Q w^{1-p'} \right)^{1/p'}$$

for all $t > 0$, all nonnegative locally integrable w , and all cubes Q in $\mathbf{R}^n$. Consequently, $A_p(w_t) \leq A_p(w)$ for each nonnegative locally integrable w on $\mathbf{R}^n$, and this is valid as long as $t > 0$ and $1 < p < \infty$.

We conclude the proof by appealing to Fatou's lemma. Since $\int_Q w$ and $\int_Q w^{1-p'}$ are always positive or infinite, for each cube Q we have

$$\begin{aligned} \frac{1}{|Q|} \left(\int_Q w \right)^{1/p} \left(\int_Q w^{1-p'} \right)^{1/p'} &\leq \frac{1}{|Q|} \left(\liminf_{t \rightarrow 0+} \int_Q w_t \right)^{1/p} \left(\liminf_{t \rightarrow 0+} \int_Q w_t^{1-p'} \right)^{1/p'} \\ &\leq \liminf_{t \rightarrow 0+} \frac{1}{|Q|} \left(\int_Q w_t \right)^{1/p} \left(\int_Q w_t^{1-p'} \right)^{1/p'} \\ &\leq \liminf_{t \rightarrow 0+} A_p(w_t) \leq \limsup_{t \rightarrow 0+} A_p(w_t) \leq A_p(w). \end{aligned}$$

Taking the supremum over all cubes completes the proof. $\square$

3. CONCLUDING REMARKS

Given any weight w that satisfies $A_{p_1}(w) > C_1$ and $A_{p_2}(w) \leq C_2$, for sufficiently small t the weight w_t will satisfy exactly the same inequalities, and w_t is in every A_p class. Consequently, no such system of inequalities can be used to show that the given weight w fails to belong to some weight class.

Our theorem fills a gap in some of the proofs that $\{p > 1 : w \in A_p\}$ is open. For example, the A_p condition and a Calderón-Zygmund covering argument can be used to show

$$\int_Q w^{1-q'} \leq C(p, q, A_p(w)) \int_Q w^{1-q'} + |Q|^{q'(1-p/q)} \left(\int_Q w^{1-p'} \right)^{(p/q)q'/p}$$

for $1 < q < p$, with $C(p, q, A_p(w)) < 1$ when q is close enough to p . Then subtraction gives a bound for $\int_Q w^{1-q'}$ that leads to a bound for $A_q(w)$ in terms of $A_p(w)$, but to justify the subtraction $\int_Q w^{1-q'}$ needs to be finite. Such arguments are always valid when w is replaced by w_t , and then the bound for $A_q(w)$ in terms of $A_p(w)$ can be obtained by letting $t \rightarrow 0$.

REFERENCES

- [1] R. J. Bagby and B. Masedeh, *Optimal Extrapolation of Weight Conditions*, Panamerican Mathematical Journal 7 (1997), Number 1, 55-63. MR **97k**:42038
- [2] H.-M. Chung, R. A. Hunt, and D. S. Kurtz, *The Hardy-Littlewood maximal function on $L(p, q)$ spaces with weights*, Indiana Univ. Math. J. 31 (1982), 109-120. MR **83b**:42021
- [3] R. R. Coifman and C. Fefferman, *Weighted norm inequalities for maximal functions and singular integrals*, Studia Math. 51 (1974), 241-250. MR **50**:10670
- [4] F. W. Gehring, *The L^p -integrability of the partial derivatives of a quasiconformal mapping*, Acta Math. 130 (1973), 265-277. MR **53**:5861
- [5] B. Muckenhoupt, *Weighted inequalities for the Hardy maximal function*, Trans. Amer. Math. Soc. 165 (1972), 207-226. MR **45**:2461

DEPARTMENT OF MATHEMATICAL SCIENCES, NEW MEXICO STATE UNIVERSITY, LAS CRUCES, NEW MEXICO 88003

E-mail address: `rbagby@nmsu.edu`

DEPARTMENT OF MATHEMATICS, MU'TAH UNIVERSITY, MU'TAH, JORDAN

E-mail address: `basmas_59@yahoo.com`