PROCEEDINGS OF THE AMERICAN MATHEMATICAL SOCIETY Volume 131, Number 6, Pages 1907–1916 S 0002-9939(03)06662-0 Article electronically published on January 2, 2003

A TALL SPACE WITH A SMALL BOTTOM

ISTVÁN JUHÁSZ, SAHARON SHELAH, LAJOS SOUKUP, AND ZOLTÁN SZENTMIKLÓSSY

(Communicated by Alan Dow)

ABSTRACT. We introduce a general method of constructing locally compact scattered spaces from certain families of sets and then, with the help of this method, we prove that if $\kappa^{<\kappa} = \kappa$, then there is such a space of height κ^+ with only κ many isolated points. This implies that there is a locally compact scattered space of height ω_2 with ω_1 isolated points in ZFC, solving an old problem of the first author.

1. Introduction

Let us start by recalling that a topological space X is called *scattered* if every non-empty subspace of X has an isolated point and that such a space has a natural decomposition into levels, the so-called Cantor-Bendixson levels. The α^{th} Cantor-Bendixson level of X will be denoted by $I_{\alpha}(X)$. We shall write $I_{<\lambda}(X) = \bigcup_{\alpha < \lambda} I_{\alpha}(X)$. The height of X, $\operatorname{ht}(X)$, is the least α with $I_{\alpha}(X) = \emptyset$. The sequence $\langle |I_{\alpha}(X)| : \alpha \in \operatorname{ht}(X) \rangle$ is said to be the *cardinal sequence* of X. The width of X, $\operatorname{wd}(X)$, is defined by $\operatorname{wd}(X) = \sup\{|I_{\alpha}(X)| : \alpha < \operatorname{ht}(X)\}$.

The cardinality of a T_3 , in particular of a locally compact, scattered (in short: LCS) space X is at most $2^{\lceil I(X) \rceil}$, hence clearly $\operatorname{ht}(X) < (2^{\lceil I(X) \rceil})^+$. Therefore under CH there is no LCS space of height ω_2 with only countably many isolated points. On the other hand, I. Juhász and W. Weiss, [2, theorem 4], proved in ZFC that for every $\alpha < \omega_2$ there is an LCS space X with $\operatorname{ht}(X) = \alpha$ and $\operatorname{wd}(X) = \omega$. The natural question of whether the existence of an LCS space of height ω_2 with countable width follows from $\neg CH$ was answered in the negative by W. Just, who proved, [3, theorem 2.13], that if one adds Cohen reals to a model of CH, then in the generic extension there are no LCS spaces of height ω_2 and $\operatorname{wd}(X) = \omega$. On the other hand, Baumgartner and Shelah proved in [1] that it is consistent (with $\neg CH$) that such an LCS space exists.

Received by the editors April 4, 2001 and, in revised form, December 4, 2001.

²⁰⁰⁰ Mathematics Subject Classification. Primary 54A25, 06E05, 54G12, 03E20.

Key words and phrases. Locally compact scattered space, superatomic Boolean algebra.

The first, third and fourth authors were supported by the Hungarian National Foundation for Scientific Research grant no. 25745.

The second author was supported by The Israel Science Foundation founded by the Israel Academy of Sciences and Humanities. Publication 714.

The third author was partially supported by Grant-in-Aid for JSPS Fellows No. 98259 of the Ministry of Education, Science, Sports and Culture, Japan.

The above-mentioned estimate $\operatorname{ht}(X) < (2^{|I(X)|})^+$ is sharp for LCS spaces with countably many isolated points: it is easy to construct an LCS space with countable "bottom" and of height α for each $\alpha < (2^{\omega})^+$ (see Theorem 2.20). Much less is known about LCS spaces with ω_1 isolated points, for example it is a long-standing open problem of whether there is, in ZFC, an LCS space of height ω_2 and width ω_1 . In fact, as was noticed by Juhász in the mid-eighties, even the much simpler question of whether there is a ZFC example of an LCS space of height ω_2 with only ω_1 isolated points turned out to be surprisingly difficult. On the other hand, Martínez in [5, theorem 1] proved that it is consistent that for each $\alpha < \omega_3$ there is an LCS space of height α and width ω_1 . As the main result of the present paper, we shall give an affirmative answer to the above question of Juhász: in section 2 we construct, in ZFC, an LCS space of height ω_2 with ω_1 isolated points. Since the space we construct in Corollary 2.21 has width ω_2 , the following question remains:

Problem 1. Is there an LCS space X of height ω_2 and width ω_1 in ZFC?

The methods used in the proof of Corollary 2.21 do not seem to suffice to get LCS spaces with ω_1 isolated points of arbitrary height $<\omega_3$. Thus we have the following problem:

Problem 2. Is there, in ZFC, an LCS space with ω_1 isolated points and of height α for each $\alpha < \omega_3$?

Although one of our main results, Theorem 2.19, can be applied to higher cardinals, it does not seem to suffice to get the analogous result, e.g. for ω_3 instead of ω_2 . If $2^{\omega} \leq \omega_2$ but $2^{\omega_1} > \omega_2$, then neither Theorem 2.20 nor Theorem 2.19 can be applied to get an LCS space of height ω_3 with only ω_2 many isolated points. Thus the following version of Juhász' problem remains open:

Problem 3. Is there, in ZFC, an LCS space X of height ω_3 having ω_2 isolated points?

Let us mention here that the problem of the existence of (λ^+, λ) -thin-tall spaces, i.e. LCS spaces of width λ and height λ^+ , is mentioned in [7, Problem 6.4, p. 53]. However, it is erroneously stated there that the existence of a (λ^+, λ) -thin-tall space follows from $\lambda^{<\lambda} = \lambda$ or from the existence of a λ^+ -tree.

The authors would like to express their thanks to the referee for pointing out several errors in a previous version of the paper.

2. Constructing a space of height ω_2 with ω_1 isolated points

Definition 2.1. Given a family of sets \mathcal{A} we define the topological space $X(\mathcal{A}) = \langle \mathcal{A}, \tau_{\mathcal{A}} \rangle$ as follows: $\tau_{\mathcal{A}}$ is the coarsest topology in which the sets $U_{\mathcal{A}}(A) = \mathcal{A} \cap \mathcal{P}(A)$ are clopen for each $A \in \mathcal{A}$. In other words, $\{U_{\mathcal{A}}(A), \mathcal{A} \setminus U_{\mathcal{A}}(A) : A \in \mathcal{A}\}$ is a subbase for $\tau_{\mathcal{A}}$.

We shall write U(A) instead of $U_{\mathcal{A}}(A)$ if \mathcal{A} is clear from the context.

Clearly $X(\mathcal{A})$ is a 0-dimensional T_2 -space. A family \mathcal{A} is called well-founded iff $\langle \mathcal{A}, \subset \rangle$ is well-founded. In this case we can define the rank-function $\mathrm{rk} : \mathcal{A} \longrightarrow On$ as usual:

$$\mathrm{rk}(A) = \sup \{ \mathrm{rk}(B) + 1 : B \in \mathcal{A} \ \land \ B \subsetneq A \},\$$

and write $R_{\alpha}(A) = \{A \in A : rk(A) = \alpha\}.$

The family \mathcal{A} is said to be \cap -closed iff $A \cap B \in \mathcal{A} \cup \{\emptyset\}$ whenever $A, B \in \mathcal{A}$.

It is easy to see that if \mathcal{A} is \cap -closed, then a neighbourhood base in $X(\mathcal{A})$ of $A \in \mathcal{A}$ is formed by the sets

$$W(A; B_1, \ldots, B_n) = U(A) \setminus \bigcup_{1}^{n} U(B_i),$$

where $n \in \omega$ and $B_i \subseteq A$ for i = 1, ..., n. (For n = 0 we have W(A) = U(A).)

The following simple result enables us to obtain LCS spaces from certain families of sets. Let us point out, however, that not every LCS space is obtainable in this manner, but we do not dwell upon this because we will not need it.

Lemma 2.2. Assume that A is both \cap -closed and well-founded. Then X(A) is an LCS space.

Proof. Given a non-empty subset \mathcal{Y} of \mathcal{A} let A be a \subset -minimal element of \mathcal{Y} . Then $U(A) \cap \mathcal{Y} = \{A\}$, i.e. A is isolated in \mathcal{Y} . Thus $X(\mathcal{A})$ is scattered.

Next we prove that every U(A) is compact by well-founded induction on $\langle \mathcal{A}, \subset \rangle$. Assume that U(B) is compact for each $B \subseteq A$. By Alexander's subbase lemma it is enough to prove that any cover of U(A) with subbase elements contains a finite subcover. So let $\mathcal{B}, \mathcal{C} \subset \mathcal{A}$ be such that

$$U(A) \subset \bigcup_{B \in \mathcal{B}} U(B) \cup \bigcup_{C \in \mathcal{C}} (A \setminus U(C)).$$

If $A \in U(B)$ for some $B \in \mathcal{B}$, then $A \subset B$ and so $U(A) \subset U(B)$.

Hence we can assume that $A \in \mathcal{A} \setminus \mathrm{U}(C)$, i.e. $A \not\subset C$ for some $C \in \mathcal{C}$. If $A \cap C = \emptyset$, then $\mathrm{U}(A) \setminus \{\emptyset\} \subset \mathcal{A} \setminus \mathrm{U}(C)$, and we are clearly done. So we can assume that $A \cap C \neq \emptyset$, and consequently $A \cap C \in \mathcal{A}$. Then $\mathrm{U}(A) \setminus (\mathcal{A} \setminus \mathrm{U}(C)) = \mathrm{U}(A) \cap \mathrm{U}(C) = \mathrm{U}(A \cap C)$. Since $A \cap C \neq A$ the set $\mathrm{U}(A \cap C)$ is compact by the induction hypothesis, hence $\mathrm{U}(A \cap C)$ is covered by a finite subfamily \mathcal{F} of $\{\mathrm{U}(B) : B \in \mathcal{B}\} \cup \{\mathcal{A} \setminus \mathrm{U}(D) : D \in \mathcal{C}\}$. Therefore $\mathcal{F} \cup \{\mathcal{A} \setminus \mathrm{U}(C)\}$ is a finite cover of $\mathrm{U}(A)$. Consequently $\mathrm{U}(A)$ is compact.

To simplify notation, if X(A) is scattered, then we write $I_{\alpha}(A) = I_{\alpha}(X(A))$. Clearly each minimal element of A is isolated in X(A); more generally we have $\alpha \leq \operatorname{rk}(A)$ if $A \in I_{\alpha}(A)$, as is shown by an easy induction on $\operatorname{rk}(A)$.

Example 2.3. Assume that $\langle T, \prec \rangle$ is a well-ordering, $\operatorname{tp} \langle T, \prec \rangle = \alpha$, and let \mathcal{A} be the family of all initial segments of $\langle T, \prec \rangle$, i.e. $\mathcal{A} = \{T\} \cup \{T_x : x \in T\}$, where $T_x = \{t \in T : t \prec x\}$. Then \mathcal{A} is well-founded, \cap -closed and it is easy to see that $X(\mathcal{A}) \cong \alpha + 1$, i.e. the space $X(\mathcal{A})$ is homeomorphic to the space of ordinals up to and including α .

Example 2.3 above shows that, in general, $R_{\alpha}(A)$ and $I_{\alpha}(A)$ may differ even for $\alpha = 0$. Indeed, if x is the successor of y in $\langle T, \prec \rangle$, then T_x is isolated in X(A) because $\{T_x\} = W(T_x; T_y) = U_A(T_x) \setminus U_A(T_y)$ is open, but $\mathrm{rk}(T_x) = \mathrm{tp}(T_x) > 0$. However, for a wide class of families, the two kinds of levels do agree. Let us call a well-founded family A rk-qood iff the following condition is satisfied:

$$\forall A \in \mathcal{A} \ \forall \alpha < \mathrm{rk}(A) \ |\{A' \in \mathcal{A} : A' \subset A \land \mathrm{rk}(A') = \alpha\}| \ge \omega.$$

Then we have the following result.

Lemma 2.4. If \mathcal{A} is well-founded, \cap -closed and $\operatorname{rk-good}$, then $I_{\alpha}(\mathcal{A}) = R_{\alpha}(\mathcal{A})$ for each α .

Proof. We prove this by induction on α . Assume that $I_{\xi}(\mathcal{A}) = R_{\xi}(\mathcal{A})$ for all $\xi < \alpha$. If $A \in R_{\alpha}(\mathcal{A})$, then $U(A) \setminus \{A\} \subset \bigcup_{\xi < \alpha} R_{\xi}(\mathcal{A}) = \bigcup_{\xi < \alpha} I_{\xi}(\mathcal{A})$ and so A is an isolated point of $\mathcal{A} \setminus \bigcup_{\xi < \alpha} I_{\xi}(\mathcal{A})$, i.e. $A \in I_{\alpha}(\mathcal{A})$. Thus we have $R_{\alpha}(\mathcal{A}) \subset I_{\alpha}(\mathcal{A})$.

Now assume that $A \in I_{\alpha}(A) \setminus R_{\alpha}(A)$. Then by our above remark $\alpha < \text{rk}(A)$, moreover there are $B_1, \ldots, B_n \in U_A(A) \setminus \{A\}$ such that

$$(\star) \qquad W(A; B_1, \dots, B_n) \setminus \{A\} \subset I_{<\alpha}(A) = R_{<}(\alpha)A.$$

Let $\eta = \max\{\alpha, \max_{i=1,\dots,n} \operatorname{rk} B_i\}$. Then $\eta < \operatorname{rk}(A)$, moreover we have $U_{\mathcal{A}}(A) \cap R_{\eta}(A) \subset \{B_1, \dots B_n\}$ by (\star) , contradicting $|U_{\mathcal{A}}(A) \cap R_{\eta}(A)| \geq \omega$. Note that this argument is valid for n = 0 as well. Indeed, in this case we have $U_{\mathcal{A}}(A) \setminus \{A\} \subset I_{<\alpha}(A)$, moreover $\eta = \alpha$. Thus we have concluded that $I_{\alpha}(A) = R_{\alpha}(A)$.

Example 2.5. For a fixed cardinal κ and any ordinal $\gamma < \kappa^+$ we define the family $\mathcal{E}_{\gamma} \subset \mathcal{P}(\kappa^{\gamma})$ as follows:

$$\mathcal{E}_{\gamma} = \Big\{ \big[\kappa^{1+\alpha} \cdot \xi, \kappa^{1+\alpha} \cdot (\xi+1) \big) : \ \alpha \le \gamma, \ \kappa^{1+\alpha} \cdot (\xi+1) \le \kappa^{1+\gamma} \Big\}.$$

Of course, throughout this definition, exponentiation means ordinal exponentiation.

 \mathcal{E}_{γ} is clearly well-founded, \cap -closed, moreover $\operatorname{rk}\left(\left[\kappa^{1+\alpha}\cdot\xi,\kappa^{1+\alpha}\cdot(\xi+1)\right)\right)=\alpha$, hence \mathcal{E}_{γ} is also rk-good. Consequently $X(\mathcal{E}_{\gamma})$ is an LCS space of height $\gamma+1$ in which the $\alpha^{\operatorname{th}}$ level is $\left\{\left[\kappa^{1+\alpha}\cdot\xi,\kappa^{1+\alpha}\cdot(\xi+1)\right):\kappa^{1+\alpha}\cdot(\xi+1)\leq\kappa^{1+\gamma}\right\}$, i.e. all levels except the top one are of size κ .

To get an LCS space of height κ^+ with "few" isolated points, our plan is to amalgamate the spaces $\{X(\mathcal{E}_\gamma): \gamma < \kappa^+\}$ into one LCS space X in such a way that $|\operatorname{I}_0(X)| \leq \kappa^{<\kappa}$. The following definition describes a situation in which such an amalgamation can be done.

Definition 2.6. A system of families $\{A_i : i \in I\}$ is called *coherent* iff $A \cap B \in A_i \cup \{\emptyset\}$ whenever $\{i, j\} \in [I]^2$, $A \in A_i$ and $B \in A_j$.

To simplify notation, we introduce the following convention. Whenever the system of families $\{A_i : i \in I\}$ is given, we will write $U_i(A)$ for $U_{A_i}(A)$, and τ_i for τ_{A_i} . If the family A is defined, then we will write U(A) for $U_A(A)$, and τ for τ_A .

Lemma 2.7. Assume that $\{A_i : i \in I\}$ is a coherent system of well-founded, \cap closed families and $A = \bigcup \{A_i : i \in I\}$. Then for each $i \in I$ and $A \in A_i$ we have $U(A) = U_i(A)$, A is also well-founded and \cap -closed, moreover $\tau_i \upharpoonright U(A) = \tau \upharpoonright U(A)$. Consequently each $X(A_i)$ is an open subspace of X(A) and thus $\{X(A_i) : i \in I\}$ forms an open cover of X(A).

Proof. Let $A \in \mathcal{A}_i$. Then it is clear from coherence that

$$U_i(A) \subseteq U(A) = \bigcup_{j \in I} \{B : B \in A_j \land B \subset A\} \subseteq U_i(A),$$

hence $U_i(A) = U(A)$.

Next let $B \in \mathcal{A}_j$. If $A \cap B = \emptyset$, then $U(A) \cap U(B) \subset \{\emptyset\}$. Now assume that $A \cap B \neq \emptyset$. Then, again by coherence, $A \cap B \in \mathcal{A}_i$ and we have

$$U(A) \cap U(B) = U_i(A) \cap U(B) = \{ C \in \mathcal{A}_i : C \subset A \land C \subset B \}$$
$$= \{ C \in \mathcal{A}_i : C \subset A \cap B \} = U_i(A \cap B).$$

In both cases $U(A) \cap U(B)$ is τ_i -open. Similarly we can see that $U(A) \setminus U(B) = U_i(A) \setminus U(B)$ is τ_i -open, hence the topologies $\tau_i \upharpoonright U(A)$ and $\tau \upharpoonright U(A)$ coincide.

To show that \mathcal{A} is well-founded, assume that $\{A_n : n \in \omega\} \subset \mathcal{A}$ and $A_0 \supseteq A_1 \supseteq \ldots$ If $A_0 \in \mathcal{A}_i$, then $\{A_n : n \in \omega\} \subset \mathcal{A}_i$ because $\mathrm{U}(A_0) = \mathrm{U}_i(A_0)$. Thus there is $n \in \omega$ with $A_m = A_n$ for each $m \ge n$ because \mathcal{A}_i is well-founded. Finally, that \mathcal{A} is \cap -closed is an easy consequence of coherence and the \cap -closedness of the families \mathcal{A}_i .

Given a system of families $\{A_i : i \in I\}$ we would like to construct a coherent system of families $\{\widehat{A}_i : i \in I\}$ such that A_i and \widehat{A}_i are isomorphic for all $i \in I$. A sufficient condition for when this can be done will be given in Lemma 2.9 below.

First, however, we need a definition. While reading it, one should remember that an ordinal is identified with the family of its proper initial segments.

Definition 2.8. Given a limit ordinal ρ and a family \mathcal{A} with $\rho \subset \mathcal{A} \subset \mathcal{P}(\rho)$, let us define the family $\widehat{\mathcal{A}}$ as follows. First consider the function $k_{\mathcal{A}}$ on ρ determined by the formula $k_{\mathcal{A}}(\eta) = U_{\mathcal{A}}(\eta + 1)$ for $\eta \in \rho$ and put

$$\widehat{\mathcal{A}} = \{ \mathbf{k}_{\mathcal{A}}^{"} A : A \in \mathcal{A} \}.$$

Since $\rho \subset \mathcal{A}$, for each $\eta \in \rho$ we clearly have $\bigcup U_{\mathcal{A}}(\eta) = \eta$ and so $k_{\mathcal{A}}(\eta) = U_{\mathcal{A}}(\eta+1) \neq U_{\mathcal{A}}(\xi+1) = k_{\mathcal{A}}(\xi)$ whenever $\{\eta,\xi\} \in \left[\rho\right]^2$. Consequently, $k_{\mathcal{A}}$ is a bijection that yields an isomorphism between \mathcal{A} and $\widehat{\mathcal{A}}$ (and so the spaces $X(\mathcal{A})$ and $X(\widehat{\mathcal{A}})$ are homeomorphic).

If the system of families $\{A_i : i \in I\}$ is given, then we write k_i for k_{A_i} for each $i \in I$.

If $\mathcal{A} \subset \mathcal{P}(\rho)$ and $\xi \leq \rho$, then we let

$$\mathcal{A} \upharpoonright \xi = \{ A \cap \xi : A \in \mathcal{A} \}.$$

For $A_0 \neq A_1 \subset \mathcal{P}(\rho)$ we let

$$\Delta(\mathcal{A}_0, \mathcal{A}_1) = \min\{\delta : \mathcal{A}_0 \upharpoonright \delta \neq \mathcal{A}_1 \upharpoonright \delta\}.$$

Clearly we always have $\Delta(\mathcal{A}_0, \mathcal{A}_1) \leq \rho$. If, in addition, $\rho + 1 \subset \mathcal{A}_0 \cap \mathcal{A}_1$, moreover both \mathcal{A}_0 and \mathcal{A}_1 are \cap -closed, then we also have

$$\Delta(\mathcal{A}_0, \mathcal{A}_1) = \min\{\delta : U_0(\delta) \neq U_1(\delta)\},\$$

because then $\mathcal{A}_i \upharpoonright \delta = U_i(\delta)$ whenever $i \in 2$ and $\delta \leq \rho$.

Lemma 2.9. Assume that κ is a cardinal, $\{A_i : i \in I\} \subset \mathcal{PP}(\kappa)$ are \cap -closed families, $\kappa \dotplus 1 \subset \mathcal{A}_i$ for each $i \in I$, and $\Delta(\mathcal{A}_i, \mathcal{A}_j)$ is a successor ordinal whenever $\{i, j\} \in [I]^2$. Then the system $\{\widehat{\mathcal{A}}_i : i \in I\}$ is coherent.

Proof. Let $A \in \mathcal{A}_i$ and $B \in \mathcal{A}_j$, where $\Delta(\mathcal{A}_i, \mathcal{A}_j) = \rho + 1$. Then $B \cap \rho \in U_j(\rho) = U_i(\rho)$ by the choice of ρ and so $A \cap B \cap \rho \in \mathcal{A}_i \cup \{\emptyset\}$ because \mathcal{A}_i is \cap -closed. Consequently we have

$$k_i'' A \cap k_j'' B = \left\{ U_i(\eta + 1) : \eta \in A \cap B \land U_i(\eta + 1) = U_j(\eta + 1) \right\}$$
$$= \left\{ U_i(\eta + 1) : \eta \in A \cap B \land \eta < \rho \right\} = k_i'' (A \cap B \cap \rho) \in \widehat{\mathcal{A}}_i \cup \{\emptyset\},$$

as required by the definition of coherence.

More is still needed if we want the "amalgamated" family to provide us a space with a small bottom, i.e. having not too many isolated points. This will be made clear by the following lemma.

Lemma 2.10. Let κ be a cardinal and $\{A_i : i \in I\} \subset \mathcal{PP}(\kappa)$ be a system of families such that

- (i) $\kappa + 1 \subset A_i$ and A_i is well-founded and \cap -closed for each $i \in I$,
- (ii) $\Delta(A_i, A_j)$ is a successor ordinal for each $\{i, j\} \in [I]^2$. Then
 - (a) the system $\{\widehat{A}_i : i \in I\}$ is coherent and thus $A = \bigcup \{\widehat{A}_i : i \in I\}$ is well-founded, \cap -closed and X(A) is covered by its open subspaces $\{X(\widehat{A}_i) : i \in I\}$.

If, in addition, we also have

- (iii) $I_0(\mathcal{A}_i) \subset [\kappa]^{<\kappa}$ for each $i \in I$ and
- (iv) $|U_i(\xi)| < \kappa$ for each $i \in I$ and $\xi \in \kappa$, then

(b)
$$I_0(\mathcal{A}) \subset \left[\left[[\kappa]^{<\kappa} \right]^{<\kappa} \right]^{<\kappa}$$
.

Proof of Lemma 2.10. The system $\{\widehat{A}_i : i \in I\}$ is coherent by Lemma 2.9, thus (a) holds by Lemma 2.7.

Consequently we have

$$I_0(\mathcal{A}) = \bigcup \{I_0(\widehat{\mathcal{A}}_i) : i \in I\}.$$

Now if $A \in I_0(A_i)$ and $\eta \in \kappa$, then $|A| < \kappa$ by (iii) and $U_i(\eta) \in \left[\left[\kappa\right]^{<\kappa}\right]^{<\kappa}$ by (iv), hence

$$\mathbf{k}_{i}^{"} A = \{\mathbf{U}_{i}(\eta + 1) : \eta \in A\} \in \left[\left[\left[\kappa\right]^{<\kappa}\right]^{<\kappa}\right]^{<\kappa}.$$

This, by $I_0(\widehat{\mathcal{A}}_i) = \{k_i'' A : A \in I_0(\mathcal{A}_i)\}, \text{ proves (b)}.$

Definition 2.11. If ρ is an ordinal and $\mathcal{A} \subset \mathcal{P}(\rho)$ let us put

$$\mathcal{A}^* = \{A \cap \xi : A \in \mathcal{A} \land \xi \leq \rho\} = \mathcal{A} \cup \{A \cap \xi : A \in \mathcal{A} \land \xi < \rho\}.$$

Definition 2.12. A family \mathcal{A} is called *chain-closed* if for each non-empty $\mathcal{B} \subset \mathcal{A}$, if \mathcal{B} is ordered by \subset (i.e. if \mathcal{B} is a chain), then $\bigcup \mathcal{B} \in \mathcal{A}$.

Lemma 2.13. If ρ is an ordinal and $\mathcal{A}_0, \mathcal{A}_1 \subset \mathcal{P}(\rho)$ are chain-closed, \cap -closed and well-founded families such that $\mathcal{A}_0^* \neq \mathcal{A}_1^*$, then $\Delta(\mathcal{A}_0^*, \mathcal{A}_1^*)$ is a successor ordinal.

Proof. Assume that δ is a limit ordinal and $\mathcal{A}_0^* \upharpoonright \gamma = \mathcal{A}_1^* \upharpoonright \gamma$ for all $\gamma < \delta$. We want to show that $\mathcal{A}_0^* \upharpoonright \delta = \mathcal{A}_1^* \upharpoonright \delta$. Since $\mathcal{A}_i^* \upharpoonright \delta = \bigcup_{\xi \leq \delta} \mathcal{A}_i \upharpoonright \xi$ and $\bigcup_{\xi < \delta} \mathcal{A}_i \upharpoonright \xi = \bigcup_{\xi < \delta} \mathcal{A}_i^* \upharpoonright \xi$, moreover $\emptyset \in \mathcal{A}_0^* \cap \mathcal{A}_1^*$, it is enough to show that $(\mathcal{A}_0 \upharpoonright \delta) \setminus \{\emptyset\} = (\mathcal{A}_1 \upharpoonright \delta) \setminus \{\emptyset\}$.

So assume that $A \in \mathcal{A}_0$ with $\sup(A \cap \delta) = \delta$ and verify that then $A \cap \delta \in \mathcal{A}_1 \upharpoonright \delta$. For each $\gamma \in A \cap \delta$ let $\mathcal{B}_{\gamma} = \{B \in \mathcal{A}_1 : B \cap (\gamma + 1) = A \cap (\gamma + 1)\}$. Since $A \cap (\gamma + 1) \in \mathcal{A}_0^* \upharpoonright (\gamma + 1) = \mathcal{A}_1^* \upharpoonright (\gamma + 1)$, there is $B \in \mathcal{A}_1$ and $\xi \leq \kappa$ such that $A \cap (\gamma + 1) = (B \cap \xi) \cap (\gamma + 1)$. Since $\gamma \in A \cap (\gamma + 1)$ it follows that $\xi > \gamma$ and so $B \cap (\gamma + 1) = (B \cap \xi) \cap (\gamma + 1)$. Thus $B \in \mathcal{B}_{\gamma}$, i.e. $\mathcal{B}_{\gamma} \neq \emptyset$. Let \mathcal{B}_{γ} be the \subset -minimal element of \mathcal{B}_{γ} , which exists because \mathcal{A}_1 is both \cap -closed and well-founded.

Then $\{B_{\gamma}: \gamma \in A \cap \delta\}$ is a chain because $B_{\gamma'} \in \mathcal{B}_{\gamma}$ for $\gamma < \gamma' < \delta, \gamma, \gamma' \in A$. Thus $B = \bigcup \{B_{\gamma}: \gamma \in A \cap \delta\} \in \mathcal{A}_1$ and clearly $A \cap \delta = B \cap \delta$ because $\sup(A \cap \delta) = \delta$. \square

The last result shows us that the operation * is useful because its application yields us families that satisfy condition (ii) of Lemma 2.10. On the other hand, the following result tells us that the LCS spaces associated with certain families modified by * do not differ significantly from the spaces given by the original families, moreover they also satisfy condition (iii) of Lemma 2.10.

Lemma 2.14. Let κ be a cardinal and $\mathcal{A} \subset [\kappa]^{\kappa}$ be well-founded and \cap -closed. Then so is \mathcal{A}^* , moreover

- (a) X(A) is a closed subspace of $X(A^*)$,
- (b) $I_0(\mathcal{A}) \subseteq I_{\kappa}(\mathcal{A}^*)$,
- (c) $ht(\mathcal{A}^*) \ge \kappa + ht(\mathcal{A}),$
- (d) $I_0(\mathcal{A}^*) \subset [\kappa]^{<\kappa}$.

Proof of Lemma 2.14. We shall write U(A) for $U_{\mathcal{A}}(A)$, and $U_*(A)$ for $U_{\mathcal{A}^*}(A)$. First observe that because

$$\mathbf{U}_*(A)\cap\mathcal{A}=\left\{\begin{array}{ll}\mathbf{U}(A) & \text{if } A\in\mathcal{A},\\ \emptyset & \text{if } A\in\mathcal{A}^*\setminus\mathcal{A}, \end{array}\right.$$

X(A) is a closed subspace of $X(A^*)$, hence (a) holds.

Now let $A \in I_0(A)$. Then there are $B_1, \ldots, B_n \in U(A) \setminus \{A\}$ such that

$${A} = W(A; B_1, \dots, B_n) = U(A) \setminus \bigcup_{i=1}^n U(B_i).$$

Since here $B_i \subsetneq A$ and $|A| = \kappa$, we can fix $\eta \in A$ such that $(A \cap \eta) \not\subset B_i$ for every $i = 1, \ldots, n$.

Now consider the basic neighbourhood

$$\mathcal{Z} = W_*(A; A \cap \eta, B_1, \dots, B_n) = U_*(A) \setminus U_*(A \cap \eta) \setminus \bigcup_{i=1}^n U_*(B_i)$$

of A in $X(A^*)$. We claim that $\mathcal{Z} = \{A \cap \xi : \eta < \xi \leq \kappa\}$. The inclusion \supset is clear from the choice of η . On the other hand, if $C \cap \xi \in \mathcal{Z}$ with $C \in \mathcal{A}$ and $\xi \leq \kappa$, then $C \cap \xi \subset A$ hence $C \cap \xi = A \cap C \cap \xi$, so as \mathcal{A} is \cap -closed we can assume that $C \subseteq A$. If we had $C \neq A$, then $\{A\} = W(A; B_1, \ldots, B_n)$ would imply $C \subset B_i$ for some i, hence $C \cap \xi \in U_*(B_i)$ and so $C \cap \xi \notin \mathcal{Z}$, a contradiction. Thus we must have C = A. Moreover, since $U_*(A \cap \eta) \supset \{A \cap \nu : \nu \leq \eta\}$, we must also have $\xi > \eta$.

By Example 2.3 we have $X(\mathcal{Z}) \cong \kappa + 1$. Moreover, the topologies $\tau_{\mathcal{Z}}$ and $\tau_{\mathcal{A}^*} \upharpoonright \mathcal{Z}$ coincide because the above argument also shows that for each $C \in \mathcal{A}$ and $\zeta \leq \kappa$ we have

$$\mathrm{U}_*(C\cap\zeta)\cap\mathcal{Z}=\mathrm{U}_*(A\cap C\cap\zeta)\cap\mathcal{Z}=\left\{\begin{array}{ll}\mathrm{U}_{\mathcal{Z}}(A\cap\zeta) & \text{if } A\subset C \text{ and } \zeta>\eta,\\ \emptyset & \text{otherwise.}\end{array}\right.$$

Hence $X(\mathcal{Z}) \cong \kappa + 1$ is a clopen subspace of $X(\mathcal{A}^*)$ and so $\{A\} = I_{\kappa}(\mathcal{Z}) = I_{\kappa}(\mathcal{A}^*) \cap \mathcal{Z}$, which proves (b).

(c) follows immediately from (a) and (b).

Finally, $I_0(\mathcal{A}^*) \subset I_{<\kappa}(\mathcal{A}^*) \subset (\mathcal{A}^* \setminus \mathcal{A}) \subset [\kappa]^{<\kappa}$, as follows immediately from (b), proving (d).

Before we can apply the amalgamation result to the families \mathcal{E}_{γ} , however, we need some further preparation that will be useful in ensuring the fulfillment of condition (iv) in Lemma 2.10.

Definition 2.15. A family \mathcal{A} is called *tree-like* iff $A \cap A' \neq \emptyset$ implies that $A \subset A'$ or $A' \subset A$, whenever $A, A' \in \mathcal{A}$.

It is easy to see that the families \mathcal{E}_{γ} given in Example 2.5 are both tree-like and chain-closed. Also, tree-like families are clearly \cap -closed.

Lemma 2.16. If δ is an ordinal and $A \subset \mathcal{P}(\delta)$ is tree-like, well-founded and chain-closed, then so is $A \upharpoonright \xi$ for each $\xi \leq \delta$.

Proof of Lemma 2.16. It is obvious that $\mathcal{A} \upharpoonright \xi$ is tree-like. To show that $\mathcal{A} \upharpoonright \xi$ is chain-closed, let $\emptyset \neq \mathcal{B} \subset \mathcal{A} \upharpoonright \xi$ be ordered by \subset . If $\mathcal{B} = \{\emptyset\}$, then $\bigcup \mathcal{B} = \emptyset \in \mathcal{A} \upharpoonright \xi$. If, however, $\mathcal{B} \neq \{\emptyset\}$, then put $\tilde{\mathcal{B}} = \{A \in \mathcal{A} : A \cap \xi \in \mathcal{B} \setminus \{\emptyset\}\}$. Since \mathcal{A} is tree-like, $\tilde{\mathcal{B}}$ is also ordered by \subset and clearly $\tilde{\mathcal{B}} \neq \emptyset$. So $\bigcup \tilde{\mathcal{B}} \in \mathcal{A}$ and $\bigcup \mathcal{B} = \bigcup \tilde{\mathcal{B}} \cap \xi \in \mathcal{A} \upharpoonright \xi$, which was to be shown.

To show that $\mathcal{A} \upharpoonright \xi$ is well-founded assume that $A_0 \cap \xi \supseteq A_1 \cap \xi \supseteq \ldots$, where each $A_n \in \mathcal{A}$. If $A_n \cap \xi = \emptyset$ for some n, then we are done. Otherwise for each $n \in \omega$ we have $A_n \cap \xi = (\bigcap_{m \le n} A_m) \cap \xi \neq \emptyset$, hence as \mathcal{A} is \cap -closed we can assume that $A_0 \supseteq A_1 \supseteq \ldots$. Since \mathcal{A} is well-founded, there is n such that $A_m = A_n$, and so $A_m \cap \xi = A_n \cap \xi$ as well, for each $m \ge n$.

Definition 2.17. Given a family $A \subset \mathcal{P}(\delta)$ and $\alpha, \beta \in \delta$ let us put

$$\mathfrak{S}^{\mathcal{A}}(\alpha,\beta) = \bigcup \{ A \in \mathcal{A} : \alpha \in A \text{ and } \beta \notin A \}.$$

Lemma 2.18. Assume that δ is an ordinal and $A \subset \mathcal{P}(\delta)$ is a tree-like, well-founded and chain-closed family with $\delta \in A$. Then

$$\mathcal{A}\setminus\{\emptyset\}=\{\delta\}\cup\{\mathfrak{S}^{\mathcal{A}}(\alpha,\beta):\alpha,\beta\in\delta\}\setminus\{\emptyset\}.$$

Consequently, $|\mathcal{A}| \leq |\delta|^2$.

Proof of Lemma 2.18. Given $\alpha, \beta \in \delta$, the family $\mathcal{S} = \{A \in \mathcal{A} : \alpha \in A, \beta \notin A\}$ is ordered by \subset because \mathcal{A} is tree-like. Thus either $\mathcal{S} = \emptyset$ and so $\mathfrak{S}^{\mathcal{A}}(\alpha, \beta) = \bigcup \mathcal{S} = \emptyset$, or if $\mathcal{S} \neq \emptyset$, then $\mathfrak{S}^{\mathcal{A}}(\alpha, \beta) = \bigcup \mathcal{S} \in \mathcal{A}$, for \mathcal{A} is chain-closed.

Assume now that $A \in \mathcal{A} \setminus \{\emptyset, \delta\}$ and let $\mathcal{D} = \{D \in \mathcal{A} : A \subsetneq D\}$. Clearly $\delta \in \mathcal{D}$. Since \mathcal{A} is tree-like, \mathcal{D} is ordered by \subset , so it has a \subset -least element, say D, because $\langle \mathcal{A}, \subset \rangle$ is also well-founded. Pick $\beta \in D \setminus A$ and let $\alpha \in A$. We claim that $A = \mathfrak{S}^{\mathcal{A}}(\alpha, \beta)$. Clearly $A \subset \mathfrak{S}^{\mathcal{A}}(\alpha, \beta)$ because $\alpha \in A$ and $\beta \notin A$. On the other hand, if $A' \in \mathcal{A}$, $\alpha \in A'$ and $\beta \notin A'$, then either $A' \subset A$ or $A \subset A'$ because \mathcal{A} is tree-like. But $\beta \notin A'$ implies that $A' \notin \mathcal{D}$, i.e. $A \subsetneq A'$ cannot hold. Thus $A' \subset A$ and so $\mathfrak{S}^{\mathcal{A}}(\alpha, \beta) = A$ is proved.

Now we are ready to collect the fruits of all the preparatory work.

Theorem 2.19. If $\kappa^{<\kappa} = \kappa$, then there is an LCS space X of height κ^+ with $|I_0(X)| = \kappa$.

Proof of Theorem 2.19. For each $\gamma < \kappa^+$ consider the well-founded, \cap -closed, rkgood family \mathcal{E}_{γ} constructed in Example 2.5:

$$\mathcal{E}_{\gamma} = \left\{ \left[\kappa^{1+\alpha} \cdot \xi, \kappa^{1+\alpha} \cdot (\xi+1) \right) : \ \alpha \le \gamma, \ \kappa^{1+\alpha} \cdot \xi < \kappa^{\gamma} \right\}.$$

Fix a bijection $f_{\gamma}: \kappa^{\gamma} \longrightarrow \kappa$, and let $\mathcal{F}_{\gamma} = \{f_{\gamma}''E : E \in \mathcal{E}_{\gamma}\}$, i.e. \mathcal{F}_{γ} is simply an isomorphic copy of \mathcal{E}_{γ} on the underlying set κ . As \mathcal{E}_{γ} is also chain-closed and tree-like, hence so is \mathcal{F}_{γ} .

We shall now show that the *-modified families $\{\mathcal{F}_{\gamma}^* : \gamma < \kappa^+\}$ satisfy conditions (i)-(iv) of Lemma 2.10. Since $\kappa \in \mathcal{F}_{\gamma}$ it follows that $\kappa \dotplus 1 \subset \mathcal{F}_{\gamma}^*$ and so (i) is true. For $\{\gamma, \delta\} \in [\kappa^+]^2$, the height of $X(\mathcal{E}_{\gamma})$ is $\gamma + 1$ and the height of $X(\mathcal{E}_{\delta})$ is $\delta + 1$, hence \mathcal{E}_{γ} and \mathcal{E}_{δ} are not isomorphic. Thus $\mathcal{F}_{\gamma} \neq \mathcal{F}_{\delta}$ and so $\mathcal{F}_{\gamma}^* \neq \mathcal{F}_{\delta}^*$ as well because $\mathcal{F}_{\gamma} = \mathcal{F}_{\gamma}^* \cap [\kappa]^{\kappa}$ and $\mathcal{F}_{\delta} = \mathcal{F}_{\delta}^* \cap [\kappa]^{\kappa}$. Hence $\Delta(\mathcal{F}_{\gamma}^*, \mathcal{F}_{\delta}^*)$ is a successor ordinal by Lemma 2.13, i.e. (ii) is satisfied.

(iii) holds by Lemma 2.14(d).

To show (iv), let us fix $\xi < \kappa$. Then $U_{\mathcal{F}_{\gamma}^*}(\xi) = \mathcal{F}_{\gamma}^* \upharpoonright \xi = \bigcup \{\mathcal{F}_{\gamma} \upharpoonright \zeta : \zeta \leq \xi\}$ where $|\mathcal{F}_{\gamma} \upharpoonright \zeta| \leq |\zeta|^2$ for all $\zeta \leq \xi$ by Lemmas 2.16 and 2.18, consequently $|\mathcal{F}_{\gamma}^* \upharpoonright \xi| \leq |\xi|^3 < \kappa$.

Thus we may apply Lemma 2.10 to the family $\mathcal{F} = \bigcup \{\widehat{\mathcal{F}_{\gamma}}^* : \gamma < \kappa^+\}$ and conclude that the space $X = X(\mathcal{F})$ is LCS, moreover $|I_0(X)| \leq \left((\kappa^{<\kappa})^{<\kappa}\right)^{<\kappa} = \kappa$. Since for every $\gamma \in \kappa^+$ the space $X(\mathcal{F}_{\gamma}^*)$ is an open subspace of X, we have $\operatorname{ht}(X) \geq \operatorname{ht}(X(\mathcal{F}_{\gamma}^*)) > \gamma$, consequently $\operatorname{ht}(X) \geq \kappa^+$.

In particular, if $2^{\omega} = \omega_1$, then the above result yields an LCS space X with $\operatorname{ht}(X) = \omega_2$ and $|\operatorname{I}_0(X)| = \omega_1$. That such a space also exists under $\neg \operatorname{CH}$, hence in ZFC, follows from the following result.

Theorem 2.20. For each $\alpha < (2^{\omega})^+$ there is a locally compact, scattered space X_{α} with $|X_{\alpha}| = |\alpha| + \omega$, $\operatorname{ht}(X_{\alpha}) = \alpha$ and $|I_0(X_{\alpha})| = \omega$.

Proof. We do induction on α . If $\alpha = \beta + 1$, then we let X_{α} be the 1-point compactification of the disjoint topological sum of countably many copies of X_{β} .

If α is limit, then we first fix an almost disjoint family $\{A_{\beta}: \beta < \alpha\} \subset [\omega]^{\omega}$, which is possible by $|\alpha| \leq 2^{\omega}$. Applying the inductive hypothesis, for each $\beta < \alpha$ we can also fix a locally compact scattered space X_{β} of cardinality $\omega + |\beta|$ and height β such that $I_0(X_{\beta}) = A_{\beta}$ and $X_{\beta} \cap X_{\gamma} = A_{\beta} \cap A_{\gamma}$ for $\{\beta, \gamma\} \in [\alpha]^2$. Now amalgamate the spaces X_{β} as follows: consider the topological space $X = \langle \bigcup_{\beta < \alpha} X_{\beta}, \tau \rangle$ where τ is the topology generated by $\bigcup_{\beta < \alpha} \tau_{X_{\beta}}$. Since $A_{\beta} \cap A_{\gamma}$ is a finite and open subspace of both X_{β} and X_{γ} it follows that each X_{β} is an open subspace of X. Consequently, X is LCS with countably many isolated points, and $\operatorname{ht}(X) = \sup_{\beta < \alpha} \operatorname{ht} X_{\beta} = \alpha$. \square

Corollary 2.21. There is, in ZFC, a locally compact scattered space of height ω_2 and having ω_1 isolated points.

Proof. If $2^{\omega} = \omega_1$, then Theorem 2.19 gives such a space.

If $2^{\omega} > \omega_1$, then $(2^{\omega})^+ \ge \omega_3$ and so according to Theorem 2.20 for each $\alpha < \omega_3$ there is locally compact, scattered space of height α and with countably many isolated points.

References

- J. E. Baumgartner and S. Shelah, Remarks on superatomic Boolean algebras, Ann. Pure Appl. Logic 33 (1987), no. 2, 109-129. MR 88d:03100
- I. Juhász and W. Weiss, On thin-tall scattered spaces, Colloquium Mathematicum, vol. XL (1978) 63–68. MR 82k:54005
- 3. W. Just, Two consistency results concerning thin-tall Boolean algebras, Algebra Universalis 20 (1985) no. 2, 135–142. MR 87c:03101
- 4. K. Kunen, Set Theory, North-Holland, New York, 1980. MR 82f:03001
- J. C. Martínez, A forcing construction of thin-tall Boolean algebras, Fundamenta Mathematicae 159 (1999), no 2, 99-113. MR 2000g:03116
- Judy Roitman, Height and width of superatomic Boolean algebras, Proc. Amer. Math. Soc. 94 (1985), no. 1, 9–14. MR 86f:06024
- S. Shelah, On what I do not understand (and have something to say) Part I, Fundamenta Mathematicae 166 (2000), no. 1-2, pp. 1-82. MR 2002a:03091

ALFRÉD RÉNYI INSTITUTE OF MATHEMATICS, P.O. BOX 127, 1364 BUDAPEST, HUNGARY *E-mail address*: juhasz@renyi.hu

Institute of Mathematics, The Hebrew University of Jerusalem, 91904 Jerusalem, Israel

E-mail address: shelah@math.huji.ac.il

ALFRÉD RÉNYI INSTITUTE OF MATHEMATICS, P.O. BOX 127, BUDAPEST, HUNGARY *E-mail address*: soukup@renyi.hu

EÖTVÖS UNIVERSITY OF BUDAPEST, 1117, BUDAPEST, PÁZMÁNY PÉTER SÉTÁNY 1/C, HUNGARY $E\text{-}mail\ address$: zoli@renyi.hu