ORTHOCOMPLETE EFFECT ALGEBRAS

GEJZA JENČA AND SYLVIA PULMANNOVÁ

(Communicated by Lance W. Small)

ABSTRACT. We prove that for every orthocomplete effect algebra E the center of E forms a complete Boolean algebra. As a consequence, every orthocomplete atomic effect algebra is a direct product of irreducible ones.

1. INTRODUCTION

Effect algebras were introduced by Foulis and Bennett in their paper [5] for the study of logical foundations of quantum mechanics. Independently, Chovanec and Kôpka introduced essentially equivalent structures called *D-posets* (see [14]). Another equivalent structure was introduced by Giuntini and Greuling in [6]. For more information about effect algebras see [4].

The class of effect algebras is a common generalization of several classes of well-established algebraic structures, in particular orthomodular lattices and MV-algebras.

In the present paper we prove that in an orthocomplete effect algebra E , the sums of all orthogonal families of central elements are central elements and that joins and meets of all families of central elements exist in E and that they are central. For finite families, these results were proved in [8]. For countable families, see [11]. As a consequence, an orthocomplete atomic effect algebra is a direct product of irreducible effect algebras. In addition, we prove that an effect algebra is κ -orthocomplete iff every chain of cardinality κ has a supremum.

2. EFFECT ALGEBRAS

An *effect algebra* is a partial algebra $(E; \oplus, 0, 1)$ with a binary partial operation $\oplus$ and two nullary operations $0, 1$ satisfying the following conditions:

- (E1) If $a \oplus b$ is defined, then $b \oplus a$ is defined and $a \oplus b = b \oplus a$.
- (E2) If $a \oplus b$ and $(a \oplus b) \oplus c$ are defined, then $b \oplus c$ and $a \oplus (b \oplus c)$ are defined and $(a \oplus b) \oplus c = a \oplus (b \oplus c)$.
- (E3) For every $a \in E$ there is a unique $a' \in E$ such that $a \oplus a' = 1$.
- (E4) If $a \oplus 1$ exists, then $a = 0$.

Received by the editors April 3, 2002.

2000 *Mathematics Subject Classification.* Primary 06F05; Secondary 03G25, 81P10.

Key words and phrases. Effect algebra, central element, orthocomplete.

This research was supported by grant G-1/7625/20 of MŠ SR, Slovakia and grant VEGA 2/7193/20.

In an effect algebra E , we write $a \leq b$ iff there is $c \in E$ such that $a \oplus c = b$. It is easy to check that $\leq$ is a partial order on E . In this partial order, 0 is the least and 1 is the greatest element of E . Moreover, it is possible to introduce a new partial operation $\ominus$; $b \ominus a$ is defined iff $a \leq b$ and then $a \oplus (b \ominus a) = b$. It can be proved that $a \oplus b$ is defined iff $a \leq b'$ iff $b \leq a'$. Therefore, it is usual to denote the domain of $\oplus$ by $\perp$. We say that elements a and b in an effect algebra E are *orthogonal* if $a \perp b$. In what follows, when we write $a \oplus b$ we mean that $a \oplus b$ is defined (i.e., $a \perp b$). Owing to associativity (E2), we may omit parentheses in $a_1 \oplus a_2 \oplus a_3$ and $a_1 \oplus a_2 \oplus \cdots \oplus a_n$, the latter term being defined by induction. We will say that the elements $a_1, \dots, a_n$ are **orthogonal** if the element $a_1 \oplus \cdots \oplus a_n$ exists in L . More generally, we say that $\{a_\alpha\}_\alpha$ is an *orthogonal family* if every finite subfamily is orthogonal.

An effect algebra need not be lattice-ordered. However, as proved in [19] and [8], the following relationship between $\wedge$, $\vee$ and $\oplus$ holds: if $a \vee b$ exists and $a \perp b$, then $a \wedge b$ exists and

$$(1) \quad a \oplus b = (a \wedge b) \oplus (a \vee b).$$

Moreover, it is easy to check (see [1]) that, for every subset B of an effect algebra such that $\vee B$ exists and for every $x \geq B$,

$$(2) \quad x \ominus (\vee B) = \wedge \{x \ominus b : b \in B\}.$$

Example 2.1. Let $(L; \wedge, \vee, ', 0, 1)$ be an orthomodular lattice. Write $a \oplus b = a \vee b$ iff $a \leq b'$, otherwise let $a \oplus b$ be undefined. Then $(L, \oplus, 0, 1)$ is an effect algebra. Effect algebras, which are associated with orthomodular lattices in this way, can be characterized as lattice-ordered effect algebras satisfying the implication

$$a \perp b \implies a \wedge b = 0.$$

Example 2.2. An *MV-algebra* (cf. [2], [15]) is a commutative semigroup $(M; \oplus, \neg, 0)$, satisfying identities $x \oplus 0 = x$, $\neg\neg x = x$, $x \oplus \neg 0 = \neg 0$ and

$$x \oplus \neg(x \oplus \neg y) = y \oplus \neg(y \oplus \neg x).$$

There is a natural partial order in an MV-algebra, given by $y \leq x$ iff $x = x \oplus \neg(x \oplus \neg y)$. Every MV-algebra $(M; \oplus, \neg, 0)$ can be considered as an effect algebra $(M; \oplus, 0, \neg 0)$ when we restrict the operation $\oplus$ to the domain $\perp = \{(x, y) : x \leq \neg y\}$. Effect algebras, which are associated with MV-algebras, can be characterized as lattice-ordered effect algebras satisfying the implication

$$a \wedge b = 0 \implies a \perp b.$$

(Cf. [17].)

Example 2.3. Let H be a Hilbert space, and let $S(H)$ denote the partially ordered group of all bounded self-adjoint linear operators on H . Put $E(H) = \{A \in S(H) : 0 \leq A \leq 1\}$; the elements of $E(H)$ are called *effects*. For $a, b \in E(H)$, write $a \oplus b = a + b$ iff $a + b \in E(H)$, otherwise let $a \oplus b$ be undefined. Then $(E(H); \oplus, 0, 1)$ is an effect algebra. We remark that for $\dim(H) \geq 2$, $E(H)$ is not lattice-ordered.

Let E_1, E_2 be effect algebras. A map $\phi : E_1 \mapsto E_2$ is called a *morphism* iff it satisfies the following condition:

$$(H1) \quad \phi(1) = 1 \text{ and if } a \perp b, \text{ then } \phi(a) \perp \phi(b) \text{ and } \phi(a \oplus b) = \phi(a) \oplus \phi(b).$$

A morphism $\phi : E_1 \mapsto E_2$ of effect algebras is called *full* iff the following condition is satisfied:

(H2) If $\phi(a) \perp \phi(b)$ and $\phi(a) \oplus \phi(b) \in \phi(E)$, then there exist $a_1, b_1 \in E_1$ such that $a_1 \perp b_1$, $\phi(a) = \phi(a_1)$ and $\phi(b) = \phi(b_1)$.

A bijective, full morphism is called an *isomorphism*. A morphism ϕ is an isomorphism if it is bijective and ϕ^{-1} is also a morphism.

Let E_1 be an effect algebra. A subset $E_2 \subseteq E_1$ is a *subeffect algebra* of E_1 iff $0, 1 \in E_2$, E_2 is closed under the $'$ operation, and $a, b \in E_2$ with $a \perp b \implies a \oplus b \in E_2$.

Another possibility to create a substructure of an effect algebra E is to restrict $\oplus$ to an interval

$$[0, a] = \{x \in E : 0 \leq x \leq a\}$$

as follows. For $x, y \in [0, a]$, $x \oplus y$ is defined iff $x \oplus y$ exists in E and $x \oplus y \in [0, a]$. We can then consider $[0, a]$ as an effect algebra, letting a act as the unit element. In what follows, we denote such effect algebras by $[0, a]_E$.

Let E be an effect algebra. A subset I of E is called an *ideal* of E iff the following condition is satisfied:

$$x, y \in I \text{ and } x \perp y \Leftrightarrow x \oplus y \in I.$$

3. ORTHOCOMPLETE EFFECT ALGEBRAS AND CENTRAL ELEMENTS

In this section, we will prove that the center of an orthocomplete effect algebra is a complete Boolean algebra. This is a generalization of [8] and [11].

Let E be an effect algebra. Suppose that there is an isomorphism $\phi : E \mapsto E_1 \times E_2$. For every such ϕ , the elements $\phi^{-1}(1, 0)$ and $\phi^{-1}(0, 1)$ are called *central elements* of E . We write $C(E)$ for the set of all central elements of an effect algebra E . We say that an effect algebra E is *irreducible* iff $C(E) = \{0, 1\}$.

Recall that an element $a \in E$ is *sharp* if $a \wedge a' = 0$, and a is *principal* if $b, c \leq a$, $b \perp c$ implies $b \oplus c \leq a$. It is easy to see that a principal element is sharp; the opposite implication need not be true, in general. Central elements can be intrinsically characterized by the following properties: (i) c and c' are principal and (ii) every element $x \in E$ admits a decomposition $x = x_1 \oplus x_2$ with $x_1 \leq c$, $x_2 \leq c'$. It can be proved that this decomposition of x is unique. In fact, $x_1 = x \wedge c$, $x_2 = x \wedge c'$. Moreover, for every central element a , the map $x \mapsto a \wedge x$ is a full morphism, which maps E onto $[0, a]_E$ (cf. [12]).

It was proved in [8] that the set of all central elements forms a sub-effect algebra of E , which is a Boolean algebra. Moreover, the joins and meets of elements of $C(E)$ exist in E and coincide with their joins and meets in $C(E)$. If $a, b \in C(E)$ are orthogonal, we have $a \vee b = a \oplus b$ and $a \wedge b = 0$.

Lemma 3.1. *If $x, y \in E$ and $a \in C(E)$, then*

$$(3) \quad a \wedge (x \oplus y) = a \wedge x \oplus a \wedge y.$$

Moreover, if $a, b \in C(E)$, and $x \in E$, then

$$(4) \quad x \wedge (a \oplus b) = x \wedge a \oplus x \wedge b.$$

Proof. Let $x \perp y$, $x, y \in E$, and $a \in C(E)$. Then $x \oplus y = x \wedge a \oplus x \wedge a' \oplus y \wedge a \oplus y \wedge a' = (x \wedge a \oplus y \wedge a) \oplus (x \wedge a' \oplus y \wedge a')$, where the first summand is under a , the second under a' . Uniqueness of the decomposition of $x \oplus y$ then yields $(x \oplus y) \wedge a = x \wedge a \oplus y \wedge a$.

If $a, b \in C(E)$, $a \perp b$ and $x \in E$, then $(a \oplus b) \wedge x = (a \oplus b) \wedge (x \wedge a \oplus x \wedge a') = (a \vee b) \wedge (x \wedge a) \oplus (a \vee b) \wedge (x \wedge a') = a \wedge x \oplus b \wedge x$. $\square$

For all $a \in C(E)$, the interval $[0, a]$ is an ideal of E . These ideals are called *central ideals*. By [3], a central ideal in an effect algebra E can be characterized as an ideal I satisfying the following conditions:

- $I = [0, a]$ for some $a \in E$.
- I is a *Riesz ideal*, i.e., if $i \in I$ and $i \leq a \oplus b$, then there exist $i_1, i_2 \in I$, such that $i_1 \leq a$, $i_2 \leq b$, $i \leq i_1 \oplus i_2$.

Let E be an effect algebra, and $\{a_\alpha\}_\alpha$ be an orthogonal family. We define $\bigoplus_\alpha a_\alpha := \bigvee_F \bigoplus (a_\alpha : \alpha \in F)$, where the supremum goes over all finite subfamilies F of α 's, if the supremum on the right-hand side exists.

We will say that an effect algebra E is *m-orthocomplete* for an infinite cardinal m if every orthogonal family of at most m elements has an $\bigoplus$ -sum in E . We say that an effect algebra E is *orthocomplete* if it is *m-orthocomplete* for every cardinal m .

The following theorem is a generalization of [11, Lemma 3.3]. For analogues in orthomodular lattices see [10], in orthoalgebras [9], [18].

Theorem 3.2. *Let E be an effect algebra and let m be a cardinal. The following are equivalent:*

- (1) *E is m -orthocomplete.*
- (2) *Every chain of at most m elements has a supremum.*

Proof. The implication that (1) implies (2) was proved in [13]. We have to prove that (2) implies (1). Assume that every chain of at most m elements in E has a supremum. Let X be an orthogonal subset of E and let $\text{card}(X) \leq m$. We may assume that X is infinite, and let γ be the first ordinal with $\text{card}(\gamma) = \text{card}(X)$. We will prove that the $\bigoplus X$ exists and it is equal to $\bigvee \Sigma$, where

$$\Sigma := (\bigoplus (x_\alpha : \alpha < \beta) : \beta < \gamma),$$

$(x_\alpha : \alpha < \gamma)$ being an indexing of X with γ . We proceed by induction by $\text{card}(X)$. If X is finite, there is nothing to prove. Let X be infinite, $\text{card}(X) \leq m$, and $\text{card}(X) = \text{card}(\gamma)$. The induction hypothesis is that for all orthogonal sets Y , $\text{card}(Y) = \beta < \gamma$, $\bigoplus Y$ exists, and

$$\bigoplus Y = \bigvee (\bigoplus (x_\sigma : \sigma < \nu) : \nu < \beta).$$

Let $X = (x_\alpha : \alpha < \gamma)$ be an indexing as desired. By induction hypothesis, the chain

$$\Sigma := (\bigoplus (x_\alpha : \alpha < \beta) : \beta < \gamma)$$

exists in E . Since $\text{card}(\Sigma) = \text{card}(X) \leq m$, the supremum $s := \bigvee \Sigma$ exists in E . Let $x_{\alpha_1}, \dots, x_{\alpha_n}$ be an arbitrary finite sequence with $\alpha_1, \dots, \alpha_n < \gamma$. Without loss of generality, we may assume that $\alpha_1 \leq \alpha_2 \leq \dots \leq \alpha_n$. Then $(x_{\alpha_1}, \dots, x_{\alpha_n}) \subseteq (x_\alpha : \alpha < \alpha_n + 1)$ and hence

$$s \geq \bigoplus (x_\alpha : \alpha < \alpha_n + 1) \geq \bigoplus (x_{\alpha_1}, \dots, x_{\alpha_n}).$$

This proves that s is an upper bound of all $\bigoplus (x_\alpha : \alpha \in F)$, F being a finite subset of the index set $(\alpha : \alpha < \gamma)$. To see that s is the desired supremum, let p be an upper bound of all $\bigoplus (x_\alpha : \alpha \in F)$, where F is a finite subsets of $(\alpha : \alpha < \gamma)$. Then for all $\beta < \gamma$, p is an upper bound of $\bigoplus (x_\alpha : \alpha < \beta)$. From this it follows that p is an upper bound of Σ , hence $p \geq s$. $\square$

Consequently, in every orthocomplete effect algebra, every chain has a supremum.

Lemma 3.3. *Let E be an orthocomplete effect algebra. Let $(a_\alpha : \alpha \in \Sigma) \subseteq E$ be an orthogonal family of central elements. Let $(x_\alpha : \alpha \in \Sigma)$ be a family of elements satisfying $x_\alpha \leq a_\alpha$, for all $\alpha \in \Sigma$. Then $\vee(x_\alpha : \alpha \in \Sigma)$ exists and equals $\oplus(x_\alpha : \alpha \in \Sigma)$.*

Proof. Obviously, $(x_\alpha : \alpha \in \Sigma)$ is an orthogonal family, so that $\oplus(x_\alpha : \alpha \in \Sigma)$ exists in E by orthocompleteness. Let M be a finite subset of Σ , and let y be any upper bound of $(x_\alpha : \alpha \in M)$. Then clearly $\forall \alpha \in M$, $x_\alpha \leq y \wedge a_\alpha$. Therefore

$$\begin{aligned} \oplus(x_\alpha : \alpha \in M) &\leq \oplus(y \wedge a_\alpha : \alpha \in M) \\ &= y \wedge (\oplus(a_\alpha : \alpha \in M)) \leq y. \end{aligned}$$

Thus, $\oplus(x_\alpha : \alpha \in M)$ is under every upper bound of $(x_\alpha : \alpha \in M)$, and we see that for every finite nonempty $M \subset \Sigma$, we have $\oplus(x_\alpha : \alpha \in M) = \vee(x_\alpha : \alpha \in M)$. This implies that

$$\begin{aligned} \oplus(x_\alpha : \alpha \in \Sigma) &= \vee_F \oplus(x_\alpha : \alpha \in F) \\ &= \vee_F \vee(x_\alpha : \alpha \in F) \\ &= \vee(x_\alpha : \alpha \in \Sigma), \end{aligned}$$

where $\vee_F$ runs over all finite subsets F of Σ . □

Theorem 3.4. *Let E be an orthocomplete effect algebra. Let $(a_\alpha : \alpha \in \Sigma)$ be an orthogonal family of central elements. Denote $a = \oplus(a_\alpha : \alpha \in \Sigma)$. Then a is central and*

$$[0, a]_E \simeq \prod_{\alpha \in \Sigma} [0, a_\alpha]_E.$$

Proof. Define a mapping $\phi : [0, a]_E \rightarrow \prod_{\alpha \in \Sigma} [0, a_\alpha]_E$ by $\phi(x) = (x \wedge a_\alpha)_{\alpha \in \Sigma}$. We shall prove that ϕ is an isomorphism.

To prove that ϕ is onto, let $(x_\alpha)_{\alpha \in \Sigma} \in \prod_{\alpha \in \Sigma} [0, a_\alpha]_E$. Observe that $(x_\alpha : \alpha \in \Sigma)$ is an orthogonal family and put $x = \oplus(x_\alpha : \alpha \in \Sigma)$. We will prove that $\phi(x) = (x_\alpha)_{\alpha \in \Sigma}$.

We have

$$\phi(x) = (x \wedge a_\alpha)_{\alpha \in \Sigma} = ((\oplus(x_\alpha : \alpha \in \Sigma)) \wedge a_\alpha)_{\alpha \in \Sigma}.$$

Fix $\beta \in \Sigma$. By associativity of $\oplus$,

$$\begin{aligned} x \wedge a_\beta &= (\oplus(x_\alpha : \alpha \in \Sigma)) \wedge a_\beta \\ &= (x_\beta \oplus (\oplus(x_\alpha : \alpha \in \Sigma \setminus \{\beta\}))) \wedge a_\beta. \end{aligned}$$

Since the family $(x_\alpha : \alpha \in \Sigma \setminus \{\beta\})$ satisfies conditions of Lemma 3.3, we have

$$\begin{aligned} (x_\beta \oplus (\oplus(x_\alpha : \alpha \in \Sigma \setminus \{\beta\}))) \wedge a_\beta \\ = (x_\beta \oplus (\vee(x_\alpha : \alpha \in \Sigma \setminus \{\beta\}))) \wedge a_\beta \end{aligned}$$

and since a_β is a central element,

$$\begin{aligned} (x_\beta \oplus (\vee(x_\alpha : \alpha \in \Sigma \setminus \{\beta\}))) \wedge a_\beta &= x_\beta \wedge a_\beta \oplus (\vee(x_\alpha : \alpha \in \Sigma \setminus \{\beta\}) \wedge a_\beta) \\ &= x_\beta \oplus (\vee(x_\alpha : \alpha \in \Sigma \setminus \{\beta\}) \wedge a_\beta). \end{aligned}$$

Since, for all $\alpha \in \Sigma \setminus \{\beta\}$, $x_\alpha \wedge a_\beta = 0$, we have

$$(\vee(x_\alpha : \alpha \in \Sigma \setminus \{\beta\}) \wedge a_\beta) = 0.$$

Thus, for all $\alpha \in \Sigma$, $x \wedge a_\alpha = x_\alpha$.

To prove that ϕ is one-to-one, it suffices to prove that, for all $x \in [0, a]$,

$$x = \oplus(x \wedge a_\alpha : \alpha \in \Sigma).$$

As

$$\begin{aligned} \oplus (x \wedge a_\alpha : \alpha \in \Sigma) &= \vee_F(\oplus(x \wedge a_\alpha : \alpha \in F)) \\ &= \vee_F(x \wedge (\oplus a_\alpha : \alpha \in F)), \end{aligned}$$

where we used (4) in the last equality, we see that $\oplus(x \wedge a_\alpha : \alpha \in \Sigma) \leq x$. Moreover, using (2),

$$\begin{aligned} x \ominus (\oplus(x \wedge a_\alpha : \alpha \in \Sigma)) &= x \ominus (\vee_F(\oplus(x \wedge a_\alpha : \alpha \in F))) \\ &= x \ominus (\vee_F(x \wedge (\oplus(a_\alpha : \alpha \in F)))) \\ &= \wedge_F(x \ominus (x \wedge (\oplus(a_\alpha : \alpha \in F)))). \end{aligned}$$

Since $\forall F$, $\oplus(a_\alpha : \alpha \in F)$ is central, we have

$$\begin{aligned} (x \ominus (x \wedge (\oplus(a_\alpha : \alpha \in F)))) &= x \wedge (\oplus(a_\alpha : \alpha \in F))' \\ &= x \wedge (\vee(a_\alpha : \alpha \in F))' \\ &= x \wedge (\wedge(a'_\alpha : \alpha \in F)). \end{aligned}$$

Therefore

$$\begin{aligned} \wedge_F(x \ominus (x \wedge (\oplus(a_\alpha : \alpha \in F)))) &= \wedge_F x \wedge (\wedge(a'_\alpha : \alpha \in F)) \\ &\leq \wedge_F a \wedge (\wedge(a'_\alpha : \alpha \in F)) \\ &= \wedge_F(a \ominus (\vee(a_\alpha : \alpha \in F)))' \\ &= a \ominus (\vee_F \vee(a_\alpha : \alpha \in F)) \\ &= 0. \end{aligned}$$

Hence, for all $x \in [0, a]_E$, $x = \oplus(x \wedge a_\alpha : \alpha \in \Sigma)$, and this implies that ϕ is one-to-one.

Let us prove that $[0, a]$ is an ideal. Obviously, $x \oplus y \in [0, a]$ implies $x \perp y$ and $x, y \in [0, a]$. To prove the opposite implication, assume that $x, y \in [0, a]$ and $x \perp y$. By the preceding paragraph, $x = \oplus(x \wedge a_\alpha : \alpha \in \Sigma)$, $y = \oplus(y \wedge a_\alpha : \alpha \in \Sigma)$. Then

$$\begin{aligned} x \oplus y &= (\oplus(x \wedge a_\alpha : \alpha \in \Sigma)) \oplus (\oplus(y \wedge a_\alpha : \alpha \in \Sigma)) \\ &= (\oplus(x \wedge a_\alpha \oplus y \wedge a_\alpha : \alpha \in \Sigma)). \end{aligned}$$

Since $\forall \alpha$, a_α is central,

$$x \wedge a_\alpha \oplus y \wedge a_\alpha = (x \oplus y) \wedge a_\alpha \leq a_\alpha.$$

Using Lemma 3.3,

$$\begin{aligned} x \oplus y &= \vee(x \wedge a_\alpha \oplus y \wedge a_\alpha : \alpha \in \Sigma) \\ &= \vee((x \oplus y) \wedge a_\alpha : \alpha \in \Sigma) \leq a. \end{aligned}$$

Therefore, $[0, a]$ is an ideal. To prove that a is central, we need to prove that $[0, a]$ is a central ideal, i.e. that $[0, a]$ is a Riesz ideal.

Assume that $z \leq x \oplus y$, where $z \in [0, a]$, $x, y \in E$. Then $z = \oplus(z \wedge a_\alpha : \alpha \in \Sigma)$. For $\forall \alpha \in \Sigma$, $z \wedge a_\alpha \leq x \wedge a_\alpha \oplus y \wedge a_\alpha$. Put $z_1 = \oplus(x \wedge a_\alpha : \alpha \in \Sigma)$, $z_2 = \oplus(y \wedge a_\alpha : \alpha \in \Sigma)$. Obviously, $z \leq z_1 \oplus z_2$, $z_1, z_2 \in [0, a]$ and $z_1 \leq x$, $z_2 \leq y$. This proves that $[0, a]$ is a central ideal, i.e. a is central. $\square$

Theorem 3.5. *Let E be an orthocomplete effect algebra. Then the centre $C(E)$ of E is a complete Boolean algebra. Moreover, all suprema and infima in $C(E)$ coincide with those in E .*

Proof. Let $(a_\alpha : \alpha \in \Sigma)$ be a family of elements in $C(E)$ indexed by a set Σ . We will prove that $\vee(a_\alpha : \alpha \in \Sigma)$ exists in E and belongs to $C(E)$. The latter statement is true for any finite set, so we may assume that Σ is infinite. Let σ be the least ordinal corresponding to $\text{card}(\Sigma)$. We may assume that σ is a limit ordinal, and replace the set Σ by the set $(\alpha : \alpha < \sigma)$, so that we are dealing with an ordinal-indexed family. Further we proceed by a transfinite induction. Assume that $y_\alpha = \vee(x_\rho : \rho < \alpha)$ exists and belongs to $C(E)$ for every $\alpha < \sigma$. This family $(y_\alpha : \alpha < \sigma)$ is nondecreasing, and $(y_{\alpha+1} \ominus y_\alpha : \alpha + 1 < \sigma)$ is an orthogonal family. Indeed, choose a finite subset $\alpha_1 < \alpha_2 < \dots < \alpha_n$ with $\alpha_n + 1 < \sigma$. We then have $y_{\alpha_1} \leq y_{\alpha_1+1} \leq y_{\alpha_2} \leq y_{\alpha_2+1} \leq \dots \leq y_{\alpha_n} \leq y_{\alpha_n+1}$. Then

$$\begin{aligned} & (y_{\alpha_1+1} \ominus y_{\alpha_1}) \oplus (y_{\alpha_2} \ominus y_{\alpha_1+1}) \oplus \dots \oplus (y_{\alpha_n+1} \ominus y_{\alpha_n}) \\ &= y_{\alpha_n+1} \ominus y_{\alpha_1} \geq (y_{\alpha+1} \ominus y_{\alpha_1}) \oplus \dots \oplus (y_{\alpha_n+1} \ominus y_{\alpha_n}). \end{aligned}$$

Hence

$$z = \oplus(y_{\alpha+1} \ominus y_\alpha : \alpha + 1 < \sigma)$$

exists and belongs to $C(E)$ by Theorem 3.4. We will prove that z is the desired join $\vee(x_\rho : \rho < \sigma)$.

First, we note that if z is an upper bound of the set $(x_\rho : \rho < \sigma)$, then it is the least one. For if $w \geq x_\rho$ for all $\rho < \sigma$, then for all $\alpha + 1 < \sigma$,

$$w \geq \vee(x_\rho : \rho < \alpha + 1) = y_{\alpha+1} \geq y_{\alpha+1} \ominus y_\alpha.$$

By Lemma 3.3, $z = \vee(y_{\alpha+1} \ominus y_\alpha : \alpha + 1 < \sigma)$, which yields $w \geq z$. Hence it is enough to show that $z \geq x_\beta$ for every $\beta < \sigma$.

If $\beta < \sigma$, σ being a limit ordinal, we have $\beta + 2 < \sigma$, whence

$$\begin{aligned} x_\beta &\leq \vee(x_\rho : \rho \leq \beta + 1) = y_{\beta+1} \\ &= \vee(y_\alpha : \alpha < \beta + 2) = \oplus(y_{\rho+1} \ominus y_\rho : \rho + 1 < \beta + 2) \leq z. \end{aligned}$$

This proves the theorem. $\square$

4. ORTHOCOMPLETE AND ATOMIC EFFECT ALGEBRAS

Recall that an element $a \neq 0$ in an effect algebra E is called an *atom* if $x \leq a$ implies $x = a$ or $x = 0$. An effect algebra E is *atomic* if every element in E majorizes an atom.

Theorem 4.1. *Let E be an orthocomplete effect algebra and let $x \in E$ be an atom. Then $\wedge(z \in C(E) : z \geq x)$ is an atom in $C(E)$.*

Proof. Put $c(x) = \wedge(z \in C(E) : z \geq x)$. By Theorem 3.5, the element $c(x)$ exists and belongs to $C(E)$. To prove that $c(x)$ is an atom of $C(E)$, assume that $d \in C(E)$, $d \leq c(x)$. Then $c(x) = d \oplus (d' \wedge c(x))$, and

$$x = x \wedge c(x) = x \wedge d \oplus x \wedge (d' \wedge c(x)).$$

If $x \neq d$, then $x \wedge d = 0$, because x is an atom, and hence $x \leq d' \wedge c(x)$. Since $d' \wedge c(x)$ belongs to $C(E)$, we obtain, by the definition of $c(x)$, that $c(x) \leq d' \wedge c(x) \leq d'$. Then we obtain $d \leq c(x)$, $d \leq c(x)'$ hence $d = 0$. This concludes the proof. $\square$

Lemma 4.2. *Let c be a central element of an effect algebra E . The centre of the effect algebra $[0, c]_E$ consists of elements $z \wedge c$, $z \in C(E)$.*

Proof. Since c is central in E , we may write $E \simeq [0, c]_E \times [0, c']_E$. If d is central in $[0, c]_E$, then $E \simeq [0, d]_E \times [0, d' \wedge c]_E \times [0, c']_E$, so that $d \in C(E)$. If $z \in C(E)$, then $z \wedge c, z' \wedge c$ are orthogonal elements of $C(E)$ with $z \wedge c \oplus z' \wedge c = c$, and therefore $[0, c]_E = [0, c \wedge z]_E \times [0, c \wedge z']_E$, hence $z \wedge c$ is central in $[0, c]_E$. $\square$

Theorem 4.3. *Every orthocomplete atomic effect algebra is a direct product of irreducible effect algebras.*

Proof. Since E is atomic, under every element c in the centre $C(E)$ of E there is an atom x of E . Theorem 4.1 implies that the element $c(x) = \bigwedge \{z \in C : x \leq z\}$ is an atom of $C(E)$. Clearly, $c(x) \leq c$. It follows that $C(E)$ is an atomic Boolean algebra, and by Theorem 3.5, $C(E)$ is complete. Let $(c_\alpha : \alpha \in \Sigma)$ denote the set of all atoms of $C(E)$. Then $(c_\alpha : \alpha \in \Sigma)$ is an orthogonal set, and by Theorem 3.4, we have

$$E \simeq \prod_{\alpha \in \Sigma} [0, c_\alpha]_E.$$

By Lemma 4.2, the centre of $[0, c_\alpha]_E$ consists of $\{0, c_\alpha\}$, hence $[0, c_\alpha]_E$ is irreducible. $\square$

REFERENCES

1. Bennett, M.K., Foulis, D.J., *Phi-symmetric effect algebras*, Found. Phys. 25 (1995), 1699-1722. MR **97a**:81015
2. Chang, C.C., *Algebraic analysis of many-valued logic*, Trans. Amer. Math. Soc. 88 (1958), 467-490. MR **20**:821
3. Chevalier, G., Pulmannová, S., *Some ideal lattices in partial abelian monoids and effect algebras*, Order 17 (2000), 75-92. MR **2001g**:06004
4. Dvurečenskij, A., Pulmannová, S., *New Trends in Quantum Structures*, Kluwer Academic Publ., Dordrecht and Ister Science Press, Bratislava, 2000. MR **2002h**:81021
5. Foulis, D., Bennett, M.K., *Effect algebras and unsharp quantum logics*, Found. Phys. 24 (1994), 1331-1352. MR **95k**:06020
6. Giuntini, R., Greuling, H., *Toward a formal language for unsharp properties*, Found. Phys. 19 (1989), 931-945. MR **90j**:81017
7. Grätzer, G., *Universal algebra*, Springer-Verlag, second edition (1979), 80-81. MR **80g**:08001
8. Greechie, R., Foulis, D., Pulmannová, S., *The center of an effect algebra*, Order 12 (1995), 91-106. MR **96c**:81026
9. Hail, E., *Orthosummable orthoalgebras*, Inter. J. Theor. Phys. 33 (1994), 1957-1984. MR **95k**:81009
10. Holland, S. Jr., *An m -orthocomplete orthomodular lattice is m -complete*, Proc. Amer. Math. Soc. 24 (1970), 716-718. MR **41**:1604
11. Jenča, G., *A Cantor-Bernstein type theorem for effect algebras*, Algebra Universalis (to appear).
12. Jenča, G., *Subcentral ideals in generalized effect algebras*, Inter. J. Theor. Phys. 39 (2000), 745-755. MR **2001j**:81018
13. Jenča, G., Pulmannová, S., *Quotients of partial abelian monoids and the Riesz decomposition property*, Algebra Universalis 47 (2002), 443-477.
14. Köpka, F., Chovanec, F., *D-posets*, Math. Slovaca 44 (1994), 21-34. MR **95i**:03134
15. Mundici, D., *Interpretations of AF C^* -algebras in Lukasiewicz sentential calculus*, J. Funct. Anal. 65 (1986), 15-63. MR **87k**:46146

16. Pulmannová, S., *Congruences in partial abelian semigroups*, Algebra Universalis 37 (1997), 119-140. MR **97i**:06001
17. Pulmannová, S., *On connections among some orthomodular structures*, Demonstratio Math. **30** (1997), 313-328. MR **98m**:06004
18. Schröder, B., *On three notions of orthosummability in orthoalgebras*, Inter. J. Theor. Phys. 38 (1999), 3305-3313. MR **2001g**:81015
19. Riečanová, Z., *On proper orthoalgebras, difference posets and Abelian inverse semigroups*, Tatra Mt. Math. Publ. 10 (1997), 119-128. MR **98d**:06003

DEPARTMENT OF MATHEMATICS, FACULTY OF ELECTRICAL ENGINEERING AND INFORMATION TECHNOLOGY, ILKOVIČOVA 3, 812 19 BRATISLAVA, SLOVAKIA

E-mail address: `jenca@kmat.elf.stuba.sk`

MATHEMATICAL INSTITUTE, SLOVAK ACADEMY OF SCIENCES, ŠTEFÁNIKOVA 49, 814 73 BRATISLAVA, SLOVAKIA

E-mail address: `pulmann@mat.savba.sk`