A TAUBERIAN THEOREM FOR VILENKIN SERIES

W. R. WADE

(Communicated by Andreas Seeger)

ABSTRACT. There are a number of papers in the literature which contain Cesàro analogues of results already known for martingale sums of Vilenkin-Fourier series. We show that for Vilenkin systems of bounded type, these are not merely analogues but actually generalizations. Indeed, we prove that convergence of the Cesàro means of a Vilenkin series S implies convergence of martingale partial sums of S itself.

§1. INTRODUCTION

Let $\mathbf{N} := \{0, 1, 2, \dots\}$, and $\mathcal{P} := \{p_0, p_1, \dots\}$ be any sequence of integers which satisfies $p_n \geq 2$. For each $n \in \mathbf{N}$ set $P_n := p_0 p_1 \cdots p_{n-1}$, where the empty product is by definition 1. The *multiplicative Vilenkin group* associated with $\mathcal{P}$ is the set $G := \{(x_0, x_1, \dots) : x_k \in \mathbf{N} \text{ and } 0 \leq x_k < p_k\}$ together with the operation

$$x \dot{+} y := (x_0 \oplus y_0, x_1 \oplus y_1, \dots),$$

where $x = (x_0, x_1, \dots)$, $y = (y_0, y_1, \dots)$ and, for each k , $x_k \oplus y_k$ represents the sum of x_k and y_k modulo p_k . The dual group of G is the system $(w_n, n \in \mathbf{N})$ defined for $x = (x_0, x_1, \dots)$ by

$$w_n(x) := \prod_{k=0}^{\infty} \exp\left(\frac{2\pi i n_k x_k}{p_k}\right),$$

where the coefficients n_k are integers which satisfy $0 \leq n_k < p_k$ and $n = \sum_{k=0}^{\infty} n_k P_k$ (see Vilenkin [4] for details). When $p_k := 2$ for all k , the group G is called the *dyadic group* and the characters w_n are called the *Walsh system*. When $p_k = O(1)$, the system $\{w_n\}$ is called a (multiplicative) Vilenkin system of *bounded type*.

It is well known that G is a compact group for each collection of radices $\mathcal{P}$, and that the corresponding *Vilenkin system* $\{w_n\}$ is a complete orthonormal system on G . Moreover, the group G can be identified with the interval $[0, 1)$ by taking an $x = (x_0, x_1, \dots) \in G$ to the number

$$\bar{x} := \sum_{k=0}^{\infty} x_k P_{k+1}^{-1}.$$

Under this identification, Haar measure on G is taken to Lebesgue measure on $[0, 1)$.

Received by the editors April 10, 2002.

2000 *Mathematics Subject Classification*. Primary 42C10, 43A75.

A *Vilenkin series* is a series of the form $S := \sum_{k=0}^{\infty} a_k w_k$, where a_k is some sequence of complex numbers. For each $x \in G$ and $n \in \mathbf{N}$, the *partial sums* of a Vilenkin series S are defined by

$$S_n(x) := \sum_{k=0}^{n-1} a_k w_k(x).$$

The partial sums S_{P_n} form a martingale in $L^2(G)$ which allows one to use martingale convergence theorems on Vilenkin series.

§2. PRELIMINARIES

For each nonnegative integer n , define intervals on G by $I_0(0) := G$, and

$$(1) \quad I_n(j) := \left\{ x = (x_0, x_1, \dots) \in G : \sum_{k=0}^{n-1} x_k P_{k+1}^{-1} = \frac{j}{P_n} \right\}$$

for $j = 0, 1, \dots, P_n - 1$, $n = 1, 2, \dots$. Recall that $\{I_n(0)\}_{n=0}^{\infty}$ is a nested sequence of subgroups of G which forms a neighborhood base at the origin, and for each n , $\{I_n(j)\}_{j=0}^{P_n-1}$ is a collection of pairwise disjoint compact sets in G whose union is G . In particular, given $x \in G$ and $n \in \mathbf{N}$, there is a unique $0 \leq j < P_n$ such that $x \in I_n(j)$. We shall denote this interval by $I_n(x)$.

Denote the Haar measure of a subset E of G by $m(E)$ and the Lebesgue measure of a subset E of $[0, 1)$ by $|E|$. Notice that under the identification of G with $[0, 1)$, the interval $I_n(j)$ corresponds to the interval $[jP_n^{-1}, (j+1)P_n^{-1})$. In particular,

$$(2) \quad m(I_n(j)) = P_n^{-1}$$

for $0 \leq j < P_n$ and $n \in \mathbf{N}$.

Let $x \in G$. A sequence of measurable sets E_j in G is said to *shrink nicely* to x if there exist integers $r_j \rightarrow 0$, as $j \rightarrow \infty$, and an absolute constant $\alpha > 0$ such that $E_j \subset I_{r_j}(x)$ and

$$m(E_j) \geq \alpha \cdot m(I_{r_j}(x)),$$

for $j = 1, 2, \dots$. By using the identification of G with the unit interval, and of Haar measure with Lebesgue measure, it is easy to check that if E is a measurable subset of G and h represents the characteristic function of E , i.e., $h(x) = 1$ for $x \in E$ and $h(x) = 0$ for $x \notin E$, then

$$\frac{1}{m(E_j)} \int_{E_j} h \, dm \rightarrow 1$$

almost everywhere $[m]$ on E for any sequence of sets E_j which shrinks nicely to x (see Rudin [2], p. 140). In particular, if E_j shrinks nicely to x , then

$$(3) \quad \lim_{j \rightarrow \infty} \frac{m(E \cap E_j)}{m(E_j)} = 1$$

for almost every $x \in E$.

§3. THE MAIN RESULT

For each $x \in G$ and each integer $n > 0$, the Cesàro means of S are defined by

$$\sigma_n(x) := \frac{1}{n} \sum_{k=1}^n S_k(x).$$

It is well known (and easy to see) that if S_k converges to some limit f , as $k \rightarrow \infty$, then $\sigma_n \rightarrow f$, as $n \rightarrow \infty$. We will obtain the following partial converse to this result. (The Walsh version of this result was obtained by Shaginyan [3] in 1974.)

Theorem 1. *Suppose E is a measurable subset of G and S is a Vilenkin series whose Cesàro means satisfy*

$$(4) \quad \lim_{n \rightarrow \infty} \sigma_{P_n}(x) = f(x), \quad x \in E,$$

for some finite-valued, measurable function f . If the Vilenkin system is of bounded type, then

$$\lim_{n \rightarrow \infty} S_{P_n}(x) = f(x)$$

for almost every $x \in E$.

Proof. We may suppose that E is of positive Haar measure, i.e., $m(E) > 0$. Fix $0 < \epsilon < m(E)/2$. By hypothesis, $f_n(x) := \sigma_{P_{n+1}}(x)$ converges to $f(x)$ for $x \in E$, as $n \rightarrow \infty$. Thus by Egoroff's Theorem, we can choose a measurable subset E_0 of E such that $f_n \rightarrow f$ uniformly on E_0 and the measure of E_0 satisfies $m(E_0) > m(E) - \epsilon$.

Let $x \in E_0$ be a point which satisfies (3) for every sequence E_j which shrinks nicely to x . For each fixed positive integer j , let $\mathcal{W}_j$ represent the collection of indices ℓ which satisfy $I_{j+1}(\ell) \subset I_j(x)$. If $\ell \in \mathcal{W}_j$ and

$$B_j := I_{j+1}(\ell) \neq A_j := I_{j+1}(x),$$

let E_j represent the union of A_j and B_j . By construction, $E_j \subset I_j(x)$ and by (2),

$$m(E_j) = 2m(I_{j+1}(x)) = \frac{2}{p_j P_j} \geq \frac{2}{M P_j} = \frac{2}{M} \cdot m(I_j(x)),$$

where $M := \sup\{p_i : i \in \mathbb{N}\}$. Thus A_j and E_j both shrink nicely to x , as $j \rightarrow \infty$. It follows from (3) that $m(E_0 \cap A_j)/m(A_j) \rightarrow 1$ and $m(E_0 \cap E_j)/m(E_j) \rightarrow 1$ as $j \rightarrow \infty$.

Let $n \in \mathbb{N}$ be so large that $m(E_0 \cap A_n)/m(A_n) < 5/4$ and $m(E_0 \cap E_n)/m(E_n) > 3/4$. Since $B_n = E_n \setminus A_n$ and $m(E_n) = 2m(A_n) = 2m(B_n)$, such a choice for n implies

$$\begin{aligned} \frac{m(E_0 \cap B_n)}{m(B_n)} &= \frac{m(E_0 \cap E_n) - m(E_0 \cap A_n)}{m(B_n)} \\ &= \frac{2m(E_0 \cap E_n)}{m(E_n)} - \frac{m(E_0 \cap A_n)}{m(A_n)} > \frac{6}{4} - \frac{5}{4} > 0. \end{aligned}$$

In particular, $E_0 \cap B_n$ is uncountable.

We have proved that if n is sufficiently large, for each $\ell \in \mathcal{W}_n$ which satisfies $I_{n+1}(\ell) \neq I_{n+1}(x)$, there exists an $x_n(\ell) \in E_0 \cap I_{n+1}(\ell)$. The restriction $I_{n+1}(\ell) \neq I_{n+1}(x)$ is not needed. Indeed, if ℓ satisfies $I_{n+1}(\ell) = I_{n+1}(x)$, then $x_n(\ell) := x$ also belongs to $E_0 \cap I_{n+1}(\ell)$. Thus, for n sufficiently large, it is possible to choose points $x_n(\ell) \in E_0 \cap I_{n+1}(\ell)$ for all $\ell \in \mathcal{W}_n$.

We claim that for these points,

$$(5) \quad \sum_{\ell \in \mathcal{W}_n} \sigma_{P_{n+1}}(x_n(\ell)) = \sigma_{P_n}(x) + (p_n - 1)S_{P_n}(x).$$

To verify (5), let $y \in G$ and notice by definition that

$$\begin{aligned} \sigma_{P_{n+1}}(y) &= \frac{1}{P_{n+1}} \sum_{k=1}^{P_{n+1}} S_k(y) \\ (6) \quad &= \frac{1}{p_n P_n} \sum_{k=1}^{P_n} S_k(y) + \frac{1}{p_n P_n} \sum_{k=P_n+1}^{P_{n+1}} S_k(y) \\ &= \frac{1}{p_n} \sigma_{P_n}(y) + \frac{1}{p_n P_n} \sum_{k=P_n+1}^{P_{n+1}} S_k(y) =: I_1(y) + I_2(y). \end{aligned}$$

Look at a typical term $T(y) := a_k w_k(y)$ in one of these sums where $y = x_n(\ell)$. If T comes from I_1 , then k is less than P_n . Since the Vilenkin function w_k is constant on $I_n(x)$ and $x_n(\ell) \in I_n(x)$, it is clear that $T(x_n(\ell)) = T(x)$ for all $\ell \in \mathcal{W}_n$. Since the number of intervals of the form $I_{n+1}(\ell)$ which are subsets of $I_n(x)$ is p_n , it follows that

$$\sum_{\ell \in \mathcal{W}_n} I_1(x_n(\ell)) = \sum_{\ell \in \mathcal{W}_n} \frac{1}{p_n} \sigma_{P_n}(x) = \sigma_{P_n}(x).$$

On the other hand, suppose that T comes from I_2 and $P_n < k \leq P_{n+1}$. Notice that $\{I_{n+1}(\ell) : \ell \in \mathcal{W}_n\}$ contains every subinterval of $I_n(x)$, hence $\{w_k(x_n(\ell)), \ell \in \mathcal{W}_n\}$ ranges over every p_n th root of unity. Since the sum of p_n th roots of unity is zero, we have

$$\sum_{\ell \in \mathcal{W}_n} w_k(x_n(\ell)) = 0.$$

Consequently, adding up the values of I_2 at $y = x_n(\ell)$, as ℓ runs over $\mathcal{W}_n$, cancels higher order terms and multiplies lower order terms. Indeed, since the number of intervals of the form $I_{n+1}(\ell)$ which are subsets of $I_n(x)$ is p_n and $P_{n+1} - P_n = (p_n - 1)P_n$, we have

$$\sum_{\ell \in \mathcal{W}_n} \sum_{k=P_n+1}^{P_{n+1}} S_k(x_n(\ell)) = \sum_{\ell \in \mathcal{W}_n} \sum_{k=P_n+1}^{P_{n+1}} S_{P_n}(x) = p_n(p_n - 1)P_n S_{P_n}(x).$$

It follows from (6) that

$$\sum_{\ell \in \mathcal{W}_n} \sigma_{P_{n+1}}(x_n(\ell)) = \sigma_{P_n}(x) + (p_n - 1)S_{P_n}(x),$$

i.e., (5) holds.

Solve (5) for S_{P_n} to write

$$(7) \quad S_{P_n}(x) = \frac{1}{p_n - 1} \left(\sum_{\ell \in \mathcal{W}_n} \sigma_{P_{n+1}}(x_n(\ell)) - \sigma_{P_n}(x) \right).$$

By construction, $x_n(\ell)$ belongs to $I_n(x)$, hence $x_n(\ell) \rightarrow x$, as $n \rightarrow \infty$, for each $\ell \in \mathcal{W}_n$. Since the characters w_k are continuous on the group, $w_k(x_n(\ell)) \rightarrow w_k(x)$, as $n \rightarrow \infty$, for each $k \in \mathbf{N}$. But f is the uniform limit of f_n on E_0 and each f_n is a Vilenkin polynomial. Since $x_n(\ell) \in E_0$, it follows that $f(x_n(\ell)) \rightarrow f(x)$ as $n \rightarrow \infty$.

Let g (respectively, g_n) represent the real part of f (respectively, of f_n). By construction and the observation just made, we can choose n so large that

$$(9) \quad \operatorname{Re} [\sigma_{P_{n+1}}(x_n(\ell))] = g_n(x_n(\ell)) \geq g(x_n(\ell)) - \frac{\epsilon}{2} \geq g(x) - \epsilon.$$

Substituting this estimate back into (7), we obtain

$$(8) \quad \begin{aligned} \operatorname{Re} (S_{P_n}(x)) &\geq \frac{1}{p_n - 1} (p_n g(x) - p_n \epsilon - \operatorname{Re} [\sigma_{P_n}(x)]) \\ &\geq \frac{1}{p_n - 1} (p_n g(x) - \operatorname{Re} [\sigma_{P_n}(x)]) - M\epsilon. \end{aligned}$$

Take the limit infimum of this inequality as $n \rightarrow \infty$, apply hypothesis (4), and then let $\epsilon \rightarrow 0$. It follows that

$$\liminf_{n \rightarrow \infty} \operatorname{Re} [S_{P_n}(x)] \geq g(x) = \operatorname{Re} [f(x)]$$

for almost every $x \in E_0$. Since $|E_0| > |E| - \epsilon$, we can replace E_0 by E . Hence, $\liminf_{n \rightarrow \infty} \operatorname{Re} [S_{P_n}] \geq \operatorname{Re} [f]$ almost everywhere on E . By repeating the entire argument with the inequalities in (9) reversed, we see that $\limsup_{n \rightarrow \infty} \operatorname{Re} [S_{P_n}] \leq \operatorname{Re} [f]$ almost everywhere on E . A similar argument works for the imaginary parts. We conclude that $S_{P_n} \rightarrow f$ almost everywhere on E as $n \rightarrow \infty$. $\square$

REFERENCES

1. J.E. Daly and K.L. Phillips, *A note on H^1 multipliers for locally compact Vilenkin groups*, Canad. Math. Bull. **41** (1998), 392–397. MR **99j**:43005
2. W. Rudin, *Real and Complex Analysis, 3rd edition*, McGraw–Hill, Inc., 1987. MR **88k**:00002
3. L.A. Shaginyan, *On definite limits and sets of limit functions of Walsh series*, Mat. Sbornik **95** (1974), 263–271.
4. N. Ya. Vilenkin, *On a class of complete orthonormal systems*, Izv. Akad. Nauk. SSSR, Ser. Mat. **11** (1947), 363–400. MR **9**:224h

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF TENNESSEE, KNOXVILLE, TENNESSEE 37996