p -HYPONORMALITY IS NOT TRANSLATION-INVARIANT

MUNEO CHŌ AND JUN IK LEE

(Communicated by Joseph A. Ball)

ABSTRACT. In this note we provide an example of a semi-hyponormal Hilbert space operator T for which $T - \lambda$ is not p -hyponormal for some $\lambda \in \mathbb{C}$ and all $0 < p \leq \frac{1}{2}$.

Let $\mathcal{H}$ be a complex Hilbert space and $B(\mathcal{H})$ be the algebra of all bounded linear operators on $\mathcal{H}$. An operator $T \in B(\mathcal{H})$ is said to be p -hyponormal if

$$(T^*T)^p - (TT^*)^p \geq 0.$$

If $p = 1$, T is hyponormal and if $p = \frac{1}{2}$, T is called semi-hyponormal. It is well known that q -hyponormal operators are p -hyponormal for $p \leq q$, by Löwner's Theorem. Throughout this note we assume $0 < p < 1$. The notion of semi-hyponormality was introduced by D. Xia [Xi] and the notion of p -hyponormal operators was introduced by A. Aluthge [Al]. The p -hyponormal operators have been studied by many authors (cf. [Al], [AlW], [Ch], [ChH], [ChI], [ChJ]). An operator $T \in B(\mathcal{H})$ is said to be *paranormal* if $\|T^2x\| \geq \|Tx\|^2$ for every unit vector $x \in \mathcal{H}$. It is well known (cf. [An], [ChJ], [FHM]) that

$$(0.1) \quad p\text{-hyponormal} \implies \text{paranormal}.$$

The p -hyponormal operators share many properties with hyponormal operators: for example, if T is p -hyponormal, then (cf. [Al], [Ch], [ChH])

- (i) T is normaloid (i.e., norm equals spectral radius);
- (ii) T is reduced by its eigenspaces;
- (iii) if T is invertible, then T^{-1} is also p -hyponormal.

However, it has remained open whether p -hyponormality is translation-invariant. For a long time researchers have guessed that this is not true. But no concrete example has been found. The purpose of this note is to provide an example which shows that p -hyponormality is *not* translation-invariant.

Concrete examples of non-hyponormal p -hyponormal are scarce in the literature before [AlW]. But the following result in [AlW] helps provide an abundance of examples.

Theorem 1 ([AlW, Corollary 1]). *If $T \in B(\mathcal{H})$ is p -hyponormal and n is a positive integer, then T^n is $\frac{p}{n}$ -hyponormal for every $0 < p \leq 1$.*

Received by the editors February 14, 2000 and, in revised form, April 26, 2002.

2000 *Mathematics Subject Classification*. Primary 47B20, 47B35.

Key words and phrases. Semi-hyponormal, p -hyponormal, paranormal, translation-invariant.

To see how Theorem 1 gives rise to many examples of operators which are semi-hyponormal but not hyponormal, recall the definition of Toeplitz operators on the Hardy space $H^2(\mathbb{T})$ of the unit circle $\mathbb{T}$. Recall that the Hilbert space $L^2(\mathbb{T})$ has a canonical orthonormal basis given by the trigonometric functions $e_n(z) = z^n$, for all $n \in \mathbb{Z}$, and that the Hardy space $H^2(\mathbb{T})$ is the closed linear span of $\{e_n : n = 0, 1, \dots\}$. An element $f \in L^2(\mathbb{T})$ is said to be analytic if $f \in H^2(\mathbb{T})$, and co-analytic if $f \in L^2(\mathbb{T}) \ominus H^2(\mathbb{T})$. If P denotes the orthogonal projection from $L^2(\mathbb{T})$ to $H^2(\mathbb{T})$, then for every $\varphi \in L^\infty(\mathbb{T})$ the operator T_φ defined by

$$T_\varphi g := P(\varphi g) \quad (g \in H^2(\mathbb{T}))$$

is called the *Toeplitz operator* with symbol φ . If φ is a trigonometric polynomial of the form $\varphi(z) = \sum_{n=-m}^N a_n z^n$, where a_{-m} and a_N are nonzero, then T_φ is called a *trigonometric Toeplitz operator*.

The following theorem gives a useful necessary condition for hyponormality and normality of trigonometric Toeplitz operators.

Theorem 2 ([FaL]). *Suppose that φ is a trigonometric polynomial of the form $\varphi(z) = \sum_{n=-m}^N a_n z^n$, where a_{-m} and a_N are nonzero.*

- (i) *If T_φ is hyponormal, then $m \leq N$ and $|a_{-m}| \leq |a_N|$.*
- (ii) *If T_φ is normal, then $m = N$ and $|a_{-m}| = |a_N|$.*

The following theorem together with Theorem 1 gives abundant examples which are semi-hyponormal operators but not hyponormal.

Theorem 3 ([CuL, Theorem 3.2]). *Every trigonometric Toeplitz operator whose square is hyponormal must be normal or analytic.*

For example, if φ is a trigonometric polynomial of the form

$$\varphi(z) \equiv \sum_{n=-m}^N a_n z^n \quad (a_{-m} \neq 0 \text{ and } a_N \neq 0)$$

and satisfies

- (i) $m < N$,
- (ii) $|a_{-m}| = |a_N|$,
- (iii) $\overline{a_N} \begin{pmatrix} a_{-1} \\ a_{-2} \\ \vdots \\ a_{-m} \end{pmatrix} = a_{-m} \begin{pmatrix} \overline{a_N - m + 1} \\ \overline{a_N - m + 2} \\ \vdots \\ \overline{a_N} \end{pmatrix}$,

then T_φ is hyponormal by an argument of [FaL, Theorem 1.4], but by Theorem 3, T_φ^2 is not hyponormal since T_φ is neither analytic nor normal. However by Theorem 1, T_φ^2 is semi-hyponormal.

We now have

Theorem 4. *p -hyponormality is not translation-invariant; more precisely, there exists an operator T satisfying that T is semi-hyponormal, but $T - \lambda$ is not p -hyponormal for some $\lambda \in \mathbb{C}$ and any $p > 0$.*

Proof. If U is the unilateral shift on ℓ_2 , define

$$S \equiv 4U^2 + U^{*2} + 2UU^* + 2.$$

We now claim that (i) S is semi-hyponormal, and (ii) $S - 4$ is not p -hyponormal for any $p > 0$.

(i) Define $\varphi(z) = 2z + z^{-1}$. Then it is easy to see that T_φ is hyponormal, but T_φ^2 is not hyponormal by Theorem 2. A straightforward calculation shows that $T_\varphi^2 = S$. Thus, by Theorem 1, S is semi-hyponormal.

(ii) To show that $S - 4$ is not p -hyponormal for any $p > 0$, in view of (0.1) it suffices to show that $S - 4$ is not paranormal. Indeed,

$$\begin{aligned} \|(S - 4)e_0\|^2 &= \|(4U^2 + U^{*2} + 2UU^* - 2)e_0\|^2 \\ &= \|4e_2 - 2e_0\|^2 = 20 \end{aligned}$$

and

$$\begin{aligned} \|(S - 4)^2 e_0\| &= \|(4U^2 + U^{*2} + 2UU^* - 2)(4e_2 - 2e_0)\| \\ &= \|8e_0 - 8e_2 + 16e_4\| = \sqrt{384}, \end{aligned}$$

which shows that $\|(S - 4)e_0\|^2 > \|(S - 4)^2 e_0\|$. This implies that $S - 4$ is not paranormal. $\square$ $\square$

REFERENCES

- [Al] A. Aluthge, *On p -hyponormal operators for $0 < p < 1$* , Int. Eq. Op. Th. **13** (1990), 307–315. MR **91a**:47025
- [AlW] A. Aluthge and D. Wang, *Powers of p -hyponormal operators*, J. Inequal. Appl. **3** (1999), 279–284. MR **2000m**:47028
- [An] T. Ando, *Operators with a norm condition*, Acta Sci. Math. (Szeged) **33** (1972), 169–178. MR **47**:9334
- [Ch] M. Chō, *Spectral properties of p -hyponormal operators*, Glasgow Math. J. **36** (1994), 117–122. MR **94m**:47043
- [ChH] M. Chō and T. Huruya, *p -hyponormal operators ($0 < p < \frac{1}{2}$)*, Commentationes Math. **33** (1993), 23–29. MR **95b**:47021
- [ChI] M. Chō and M. Itoh, *Putnam's inequality for p -hyponormal operators*, Proc. Amer. Math. Soc. **123** (1995), 2435–2440. MR **95j**:47027
- [ChJ] M. Chō and H. Jin, *On p -hyponormal operators*, Nihonkai Math. J. **6** (1995), 201–206. MR **96j**:47019
- [CuL] R.E. Curto and W.Y. Lee, *Joint hyponormality of Toeplitz pairs*, Memoirs Amer. Math. Soc. **150** (2001). MR **2002c**:47042
- [FaL] D.R. Farenick and W.Y. Lee, *Hyponormality and spectra of Toeplitz operators*, Trans. Amer. Math. Soc. **348** (1996), 4153–4174. MR **97k**:47027
- [FHM] M. Fujii, R. Nakamoto and H. Watanabe, *The Heinz–Kato–Furuta inequality and hyponormal operators*, Math. Japonica **39** (1994), 469–472. MR **95k**:47033
- [Xi] D. Xia, *Spectral Theory of Hyponormal Operators*, Birkhäuser Verlag, Boston, 1983. MR **87j**:47036

DEPARTMENT OF MATHEMATICS, KANAGAWA UNIVERSITY, YOKOHAMA 221-8686, JAPAN
E-mail address: chiyom01@kanagawa-u.ac.jp

DEPARTMENT OF MATHEMATICS, SUNGKYUNKWAN UNIVERSITY, SUWON 440-746, KOREA
E-mail address: jilee@math.skku.ac.kr