THE MINIMUM NUMBER OF ACUTE DIHEDRAL ANGLES OF A SIMPLEX

GANGSONG LENG

(Communicated by Wolfgang Ziller)

ABSTRACT. For any n -dimensional simplex $\Omega \subset R^n$, we confirm a conjecture of Klamkin and Pook (1988) that there are always at least n acute dihedral angles in Ω .

Let Ω be an n -simplex in R^n with vertices $A_1, A_2, \dots, A_{n+1}$ (i.e., $\Omega = \langle A_1, A_2, \dots, A_{n+1} \rangle$), let $\Omega_i = \langle A_1, \dots, A_{i-1}, A_{i+1}, \dots, A_{n+1} \rangle$ denote its facet which lies in a hyperplane π_i , and let e_i be the unit outer normal vector of π_i ($i = 1, 2, \dots, n+1$). Denote by θ_{ij} the supplement of the angle between e_i and e_j ; we call θ_{ij} the dihedral angle between Ω_i and Ω_j .

Obviously, for any n -simplex, there exist $\binom{n+1}{2}$ dihedral angles. In [1], Klamkin shows that there are always at least three acute dihedral angles in any tetrahedron. Further, he posed the question: can one generalize this result to an n -simplex? Pook [1] conjectures that n is the required minimum number of acute dihedral angles in any n -simplex.

In this paper, we confirm this supposition.

Theorem 1. *There exist at least n acute dihedral angles in any n -simplex, and there exists an n -simplex which has only n acute dihedral angles.*

It is clear that Theorem 1 can be replaced equivalently by the following statement.

Theorem 1'. *There exist at most $\frac{1}{2}n(n-1)$ obtuse dihedral angles in any n -simplex, and there exists an n -simplex which has only $\frac{1}{2}n(n-1)$ obtuse dihedral angles.*

To prove Theorem 1', we need the following three lemmas.

Lemma 1. *Let $\{x_1, x_2, \dots, x_n\}$ be a given linearly independent set of vectors from R^n . Then there exists an n -simplex Ω which takes $\frac{x_1}{\|x_1\|}, \frac{x_2}{\|x_2\|}, \dots, \frac{x_n}{\|x_n\|}$ as n unit outer normal vectors of facets.*

Proof. Since $x_1, x_2, \dots, x_n$ are linearly independent, there exist n linearly independent vectors $v_1, v_2, \dots, v_n$ such that

$$\langle x_i, v_j \rangle = \delta_{ij} \|x_i\|^2 \quad (i, j = 1, 2, \dots, n),$$

Received by the editors May 31, 2000 and, in revised form, August 8, 2001.

2000 *Mathematics Subject Classification.* Primary 52A20.

Key words and phrases. Simplex, dihedral angle, dual basis.

This work was supported by the National Natural Sciences Foundation of China (10271071).

where δ_{ij} is the Kronecker delta symbol and $\langle, \rangle$ denotes the ordinary inner product of R^n .

Without loss of generality, we may assume that $v_i = \overrightarrow{A_{n+1}A_i}$ ($i = 1, 2, \dots, n$). We now consider the n -simplex $\Omega = \langle A_1, A_2, \dots, A_{n+1} \rangle$. Since $x_i \perp v_j$ ($i \neq j$), it follows that

$$x_i \perp \pi_i \quad (i = 1, 2, \dots, n),$$

where π_i is the hyperplane spanned by the facet Ω_i . This completes the proof of Lemma 1. $\square$

Lemma 2 (Yang and Zhang [3]). *Let Ω and Ω' be two n -simplices with dihedral angles θ_{ij} and θ'_{ij} ($1 \leq i < j \leq n+1$), respectively. If*

$$\theta_{ij} \leq \theta'_{ij} \quad (1 \leq i < j \leq n+1),$$

then Ω and Ω' are similar, namely,

$$\theta_{ij} = \theta'_{ij} \quad (1 \leq i < j \leq n+1). \quad \square$$

Let $\{u_1, \dots, u_n\}$ and $\{s_1, \dots, s_n\}$ be two bases of R^n . They are called dual if

$$\langle u_i, s_j \rangle = \delta_{ij} \quad (1 \leq i \leq j \leq n).$$

It is well known that there exists a unique dual basis for a given basis of R^n .

Lemma 3. *Let $U = \{u_1, \dots, u_n\}$ be a basis of R^n and $\{s_1, \dots, s_n\}$ be the dual basis of U . If $\langle u_i, u_j \rangle < 0$ ($i \neq j$), then $\langle s_i, s_j \rangle > 0$ for all i, j .*

Proof. We first consider $n = 2$. Suppose that $\langle u_1, u_2 \rangle < 0$ and $\{s_1, s_2\}$ is the dual basis. Then $s_1 = \langle s_1, s_1 \rangle u_1 + \langle s_1, s_2 \rangle u_2$. Hence

$$0 = \langle s_1, u_2 \rangle = \langle s_1, s_1 \rangle \langle u_1, u_2 \rangle + \langle s_1, s_2 \rangle \|u_2\|^2.$$

Thus $\langle s_1, s_2 \rangle > 0$.

Turning to the case of general n , let

$$H_1 = \{x \in R^n \mid \langle u_1, x \rangle = 0\}.$$

Then the vectors $u_2, \dots, u_n$ are on the opposite side of H_1 from u_1 . For $j \geq 2$, let $u_j = y_j + z_j$, $y_j \perp H_1$, $z_j \in H_1$; then

$$y_j = \frac{\langle u_1, u_j \rangle}{\|u_1\|^2} u_1, \quad z_j = u_j - \frac{\langle u_1, u_j \rangle}{\|u_1\|^2} u_1.$$

If $i \neq j$ and $i, j \geq 2$, then

$$\begin{aligned} \langle z_i, z_j \rangle &= \langle u_i - \frac{\langle u_1, u_i \rangle}{\|u_1\|^2} u_1, u_j - \frac{\langle u_1, u_j \rangle}{\|u_1\|^2} u_1 \rangle \\ &= \langle u_i, u_j \rangle - 2 \frac{\langle u_1, u_i \rangle \langle u_1, u_j \rangle}{\|u_1\|^2} + \frac{\langle u_1, u_i \rangle \langle u_1, u_j \rangle}{\|u_1\|^2} \\ &= \langle u_i, u_j \rangle - \frac{\langle u_1, u_i \rangle \langle u_1, u_j \rangle}{\|u_1\|^2} < 0. \end{aligned}$$

Note that $s_k \in H_1$ for $k \geq 2$, and

$$\langle z_j, s_k \rangle = \langle u_j - \frac{\langle u_1, u_j \rangle}{\|u_1\|^2} u_1, s_k \rangle = \langle u_j, s_k \rangle = \delta_{jk}$$

for $k, j \geq 2$.

Since $\{z_2, \dots, z_n\}$ and $\{s_2, \dots, s_n\}$ are biorthogonal sets of $n-1$ vectors in the $(n-1)$ -dimensional subspace H_1 , they must be dual bases in H_1 , and $\langle z_i, z_j \rangle < 0$

when $i \neq j$. If $\dim H_1 > 2$, then we can continue projecting to lower-dimensional subspaces until $\{u_1, \dots, u_n\}$ is mapped to a basis $G = \langle g_{n-1}, g_n \rangle$ in a two-dimensional subspace with $\langle g_{n-1}, g_n \rangle < 0$ and with $\{s_{n-1}, s_n\}$ the dual basis of G . But then in dimension 2, $\langle s_{n-1}, s_n \rangle > 0$, completing the proof. $\square$

Proof of Theorem 1'. We will first construct an n -simplex that has exactly $\frac{1}{2}n(n-1)$ obtuse dihedral angles. Denote by $\bar{\Omega}$ an n -simplex all of whose dihedral angles are acute (obviously, such an $\bar{\Omega}$ exists, e.g., regular simplex).

Let $u_1, u_2, \dots, u_n$ be unit outer normal vectors of n facets of $\bar{\Omega}$. Then

$$\langle u_i, u_j \rangle < 0, \quad i \neq j,$$

and $U = \{u_1, \dots, u_n\}$ is a basis of R^n . Further let $\{s_1, \dots, s_n\}$ be the dual basis of U . By Lemma 1, there exists an n -simplex Ω which takes $e_1, \dots, e_n$ as unit outer normal vectors of n facets, where $e_i = \frac{s_i}{\|s_i\|}$. By Lemma 3, we have

$$(1.1) \quad \langle e_i, e_j \rangle = \frac{1}{\|s_i\| \|s_j\|} \langle s_i, s_j \rangle > 0, \quad 1 \leq i, j \leq n.$$

Let e_{n+1} be the unit outer normal vector of the $(n+1)$ -th facet of Ω , and f_i the $(n-1)$ -dimensional volume of the facet Ω_i of Ω . By Minowski's realization theorem (see [2, p. 390]), we have

$$\sum_{i=1}^{n+1} f_i e_i = 0.$$

Therefore, for $j \in \{1, 2, \dots, n\}$,

$$(1.2) \quad \begin{aligned} \langle e_{n+1}, e_j \rangle &= \left\langle -\sum_{i=1}^n e_i \frac{f_i}{f_{n+1}}, e_j \right\rangle \\ &= -\sum_{i=1}^n \langle e_i, e_j \rangle \frac{f_i}{f_{n+1}} < 0. \end{aligned}$$

By (1.1) and (1.2), we know that Ω has $\frac{1}{2}n(n-1)$ obtuse dihedral angles and n acute dihedral angles.

Assume that there exists an n -simplex whose obtuse dihedral angles are more than $\frac{1}{2}n(n-1)$. Without loss of generality, we may assume that

$$\alpha_1, \dots, \alpha_{\frac{1}{2}n(n-1)}, \alpha_{\frac{1}{2}n(n-1)+1}, \dots, \alpha_{\frac{1}{2}n(n-1)+j}$$

are its obtuse dihedral angles, while $\alpha_{\frac{1}{2}n(n-1)+j+1}, \dots, \alpha_{\frac{1}{2}n(n+1)}$ are acute. Thus, we can construct a rectangular n -simplex with acute dihedral angles $\alpha_{\frac{1}{2}n(n-1)+j+1}, \dots, \alpha_{\frac{1}{2}n(n+1)}$. In fact, let $\{\bar{e}_1, \dots, \bar{e}_n\}$ be a standard base of R^n , and let

$$\bar{e}_{n+1} = -\sqrt{\frac{1-\lambda}{j}}\bar{e}_1 - \dots - \sqrt{\frac{1-\lambda}{j}}\bar{e}_j - \cos \alpha_{\frac{1}{2}n(n-1)+j+1}\bar{e}_{j+1} - \dots - \cos \alpha_{\frac{1}{2}n(n+1)}\bar{e}_n,$$

where

$$\lambda = \cos^2 \alpha_{\frac{1}{2}n(n-1)+j+1} + \dots + \cos^2 \alpha_{\frac{1}{2}n(n+1)}.$$

Then the rectangular simplex that takes $\bar{e}_1, \dots, \bar{e}_n, \bar{e}_{n+1}$ as $n+1$ unit normal vectors of facets has n acute dihedral angles

$$\arccos \sqrt{\frac{1-\lambda}{j}}, \dots, \arccos \sqrt{\frac{1-\lambda}{j}}, \alpha_{\frac{1}{2}n(n-1)+j+1}, \dots, \alpha_{\frac{1}{2}n(n+1)}.$$

By Lemma 2, this is contradictory. This completes the proof. $\square$

ACKNOWLEDGEMENTS

The author would like to thank the referees for many valuable suggestions and comments.

REFERENCES

- [1] M. S. Klamkin, L. P. Pook, Acute dihedral angles, Problems 1281, Math. Mag. 61, 5 (1988), 320.
- [2] R. Schneider, Convex Bodies: the Brunn–Minkowski Theory, Cambridge University Press, Cambridge, 1993. MR **94d**:52007
- [3] L. Yang and J. Z. Zhang, A necessary and sufficient condition for embedding a simplex with prescribed dihedral angles in E^n , Acta Math. Sinica 26 (1983), 250–256. MR **84g**:51023

DEPARTMENT OF MATHEMATICS, SHANGHAI UNIVERSITY, SHANGHAI, 200436, PEOPLE'S REPUBLIC OF CHINA

E-mail address: gleng@mail.shu.edu.cn