ON THE INVARIANCE OF CLASSES $\Phi BV, \Lambda BV$ UNDER COMPOSITION

PAMELA B. PIERCE AND DANIEL WATERMAN

(Communicated by Andreas Seeger)

ABSTRACT. The necessary and sufficient condition for $g \circ f$ to be in the class $\Phi BV, \Lambda BV$ for every f of that class whose range is in the domain of g is that g be in $\text{Lip } 1$.

Several different classes of functions arise naturally in the study of the convergence of Fourier series [W1]–[W4], [S], [Y]. We concern ourselves here with two such classes: ΛBV and ΦBV . Waterman [W1] defined the class ΛBV as follows: Suppose $\Lambda = \{\lambda_n\}$ is an increasing sequence such that $\sum_{n=1}^{\infty} 1/\lambda_n = \infty$. We say that $f \in \Lambda BV$ on $[a, b]$ if $\sum_{n=1}^{\infty} |f(I_n)|/\lambda_n < \infty$ for every set $\{I_n\}$ of nonoverlapping intervals in $[a, b]$. If $\{\lambda_n\} = \{n\}$, the resulting class is called HBV , the functions of *harmonic bounded variation*.

To define the class ΦBV , we let φ be a convex function with domain $[0, \infty)$ having the following three properties:

1. $\varphi(0) = 0$, $\varphi(x) > 0$ for $x > 0$;
2. $\frac{\varphi(x)}{x} \rightarrow 0$ as $x \rightarrow 0$;
3. $\frac{\varphi(x)}{x} \rightarrow \infty$ as $x \rightarrow \infty$.

Another property which is sometimes assumed is:

4. there exists $a > 0$ and $\delta > 0$ such that

$$\frac{\varphi(2x)}{\varphi(x)} \leq \delta \quad \text{for } x \in (0, a].$$

The last condition above is usually called the condition Δ_2 (often called Δ_2 for small values).

Let $P = \{a = x_0, x_1, \dots, x_m = b\}$ be a partition of $[a, b]$. We use the notation $I_n = [x_{n-1}, x_n]$, and we write $f(I_n) = f(x_n) - f(x_{n-1})$. Since φ may be represented as an integral of a nondecreasing function, clearly φ itself is a strictly increasing continuous function on $[0, \infty)$.

We say that $f \in \Phi BV$ [MO] on an interval $[a, b]$ if there exists a constant M such that whenever $P = \{I_n\}$ is an arbitrary partition of $[a, b]$, we have

$$\sum_P \phi(|f(I_n)|) < M.$$

Received by the editors October 21, 2002.

2000 *Mathematics Subject Classification*. Primary 26A45, 42A16.

Key words and phrases. Bounded variation, Fourier series.

Alternatively we may require the infinite sum $\sum_{n=1}^{\infty} \phi(|f(I_n)|)$ to be finite whenever $\{I_n\}_{n=1}^{\infty}$ is a collection of *nonoverlapping* intervals in $[a, b]$. These two definitions have been shown to be equivalent. We will henceforth omit all reference to the interval $[a, b]$ and simply write ΦBV to denote this class. We see the importance of the condition Δ_2 in the following theorem.

Theorem (Musielak and Orlicz). *The class ΦBV is linear if and only if Δ_2 is satisfied.*

In what follows we shall assume that φ satisfies Δ_2 .

GW represents the class of functions (necessarily having only simple discontinuities) that have a convergent Fourier series for every change of variable, and the class UGW is defined analogously with respect to uniform convergence. Chaika and Waterman [CW] proved the following.

Theorem (Chaika and Waterman). *$g \circ f$ is in one of the classes GW, UGW or HBV for each f of that class whose range is in the domain of g if and only if $g \in \text{Lip } 1$.*

Josephy [J] had proved an analogous theorem for the class BV .

We are interested now in necessary and sufficient conditions for a function f to be preserved as a member of the classes $\Lambda BV, \Phi BV$ when it is composed with a function g on the left. We prove here the following extension of the theorem of Chaika and Waterman:

Theorem. *$g \circ f$ is in the class ΛBV or ΦBV for each f of that class whose range is in the domain of g if and only if $g \in \text{Lip } 1$.*

Proof. We observe that the continuity of g is necessary. Suppose g is discontinuous. We may suppose $g(0) = 0, g(t_n) \geq 1$, where $t_n \searrow 0$ and $\sum_{n=1}^{\infty} t_n < 1$. We will define a function $f \in BV$ such that $g \circ f \notin \Lambda BV \cup \Phi BV$. Let $\{a_n\}$ be a sequence in $(0, 1)$ converging downward to 0. Let f be 0 at 0 and on $[a_1, 1]$. On each interval $[a_{n+1}, a_n]$ we define f to be a tent function such that $f(a_{n+1}) = f(a_n) = 0$ and $f((a_{n+1} + a_n)/2) = t_n$. Note that $f \in BV$ and therefore $f \in \Lambda BV$ and $f \in \Phi BV$. Let $I_n = [(a_n + a_{n+1})/2, a_n]$. Then $(g \circ f)(I_n) = g(t_n) \geq 1$. Hence

$$\sum_{n=1}^{\infty} \frac{|(g \circ f)(I_n)|}{\lambda_n} \geq \sum_{n=1}^{\infty} \frac{1}{\lambda_n} = \infty$$

and

$$\sum_{n=1}^{\infty} \varphi(|g \circ f(I_n)|) \geq \sum_{n=1}^{\infty} \varphi(1) = \infty.$$

Let us now suppose $g \in \text{Lip } 1$ and I_n is a collection of nonoverlapping intervals in the domain of f . Then there is a $c > 0$ such that for any interval I_n , $|g \circ f(I_n)| < c|f(I_n)|$. Thus if f is in ΛBV or ΦBV , then so is $g \circ f$.

Now we assume that g is continuous but $g \notin \text{Lip } 1$, and the range of f is in the domain of g . Without loss of generality, we may suppose that the domain of g contains $[0, a]$ for some positive a and that there is a sequence of disjoint intervals $J_n = [p_n, q_n]$ in the domain of g , with $p_n, q_n \searrow 0$, and a sequence of positive numbers $c_n \rightarrow \infty, c_1 > 1$, such that

$$\sum_{n=1}^{\infty} \frac{1}{c_n} < \infty, \quad |J_n| c_n \leq g(J_n),$$

and

$$\sum_{n=1}^{\infty} \varphi(q_n) < \infty.$$

Since g is continuous and $|J_n| \rightarrow 0$, we see that $|g(J_n)| \rightarrow 0$ implying $c_n|J_n| \rightarrow 0$. We may choose $\{J_n\}$ so that $\{c_n|J_n|\}$ is monotone.

We now construct a function $f \in \Lambda BV$ such that $g \circ f \notin \Lambda BV$. To do this, we define a function $L : [0, \infty) \rightarrow [0, \infty)$ by setting

$$L(q) = \sum_{k=1}^q \frac{1}{\lambda_k} \quad \text{for } q \in Z^+, \quad L(0) = 0,$$

and extending L to be linear on each interval $[q-1, q], q \in Z^+$.

We observe that L is strictly increasing on $[0, \infty)$, $L(x) \rightarrow \infty$ as $x \rightarrow \infty$, and hence L^{-1} exists. There is no loss of generality if we assume that $\lambda_1 \geq 1$. Let $k_n = \lceil L^{-1}(1/(c_n|J_n|) + 1) \rceil$, where $\lceil t \rceil$ denotes the greatest integer less than or equal to t . Then

$$k_n \rightarrow \infty, \quad L(k_n) > \frac{1}{c_n|J_n|} \quad \text{and} \quad g(J_n)L(k_n) > \frac{g(J_n)}{c_n|J_n|} \geq 1.$$

There is a $C > 1$ such that

$$\frac{1}{c_n} \leq |J_n|L(k_n) < C \frac{1}{c_n}.$$

For this C we then have

$$(1) \quad \sum_{n=1}^{\infty} |J_n|L(k_n) < C \sum_{n=1}^{\infty} \frac{1}{c_n} < \infty.$$

For each $n = 1, 2, \dots$, let $I_{n,1}, I_{n,2}, \dots, I_{n,k_n}$ be a collection of disjoint closed intervals in $(2^{-n}, 2^{-(n-1)})$ with $I_{n,m}$ to the left of $I_{n,m+1}$ for each m .

Let f be defined on each $I_{n,m}$ ($m = 1, 2, \dots, k_n; n = 1, 2, \dots$) to be the increasing linear map of $I_{n,m}$ onto J_n . Set $f(0) = f(1) = 0$. Define f to be linear on each of the component intervals of the remainder of $[0, 1]$ and continuous on $[0, 1]$. We now claim that $f \in \Lambda BV$ while $g \circ f \notin \Lambda BV$.

To see that $g \circ f \notin \Lambda BV$ we observe that

$$\begin{aligned} V_{\Lambda} \left(g \circ f, \left[\frac{\pi}{2^n}, \frac{\pi}{2^{n-1}} \right] \right) &\geq \sum_{m=1}^{k_n} \frac{(g \circ f)(I_{n,m})}{\lambda_m} \\ &= g(J_n)L(k_n) \geq 1. \end{aligned}$$

If $g \circ f$ were in ΛBV , then the right continuity of $g \circ f$ at 0 would imply $V_{\Lambda}(g \circ f, [0, \epsilon]) \rightarrow 0$ as $\epsilon \rightarrow 0$ (see [W2]). Hence $g \circ f \notin \Lambda BV$. $\square$

To see that $f \in \Lambda BV$ we first recall a definition of Banach:

Definition. If f is a continuous function, then $\mathcal{N}(y) = \mathcal{N}(f; y) = \text{card}\{x | f(x) = y\}$. (This function $\mathcal{N}$ has been called the Banach indicatrix of f .)

This notion is easily extended to regulated functions, i.e., those functions with only simple discontinuities, by adjoining a vertical line connecting $f(x+)$ and $f(x-)$ at each point x of discontinuity (see [W1]). $\mathcal{N}(y)$ is then defined to be the number of intersection points of this “extended” graph of f and the horizontal line of height

y . The following result of Waterman [W2], which generalizes a result of Goffman [G], will prove useful:

Theorem. *If $\inf f = A$ and $\sup f = B$, f has only simple discontinuities, and $L(x)$ is an increasing function such that*

$$L(n) \sim \sum_1^n \frac{1}{\lambda_k} \quad \text{as } n \rightarrow \infty,$$

then $\int_A^B L(\mathcal{N}(y)) dy < \infty$ implies that f is in the class ΛBV .

The importance of the class of functions satisfying $\int_A^B L(\mathcal{N}(y)) dy < \infty$ was first observed by Garsia and Sawyer [GS] for continuous functions and $L(x) = \log(x)$.

Now we need only observe

$$\begin{aligned} \int L(\mathcal{N}(f; y)) dy &\leq q_1 L(2) + \sum_{n=1}^{\infty} |J_n| L(2k_n) \\ &< q_1 L(2) + 2 \sum_{n=1}^{\infty} |J_n| L(k_n) \\ &< \infty \end{aligned}$$

by (1). Hence $f \in \Lambda BV$, and we have the portion of the desired result which is concerned with ΛBV .

To complete the proof for ΦBV , we now define a function P by setting

$$P(n) = \frac{1}{\varphi(c_n |J_n|)} \quad \text{for } n \in Z^+, \quad P(0) = 0,$$

and extending P to be linear on each interval $[k-1, k], k \in Z^+$. Then P is a one-to-one increasing mapping of $[0, \infty)$ onto $[0, \infty)$. Let $k_n = [P(n) + 1]$. Thus $k_n > P(n) = 1/(\varphi(c_n |J_n|))$.

Let f be defined as in the construction for ΛBV but with this definition of k_n . We now claim that $f \in \Phi BV$ while $g \circ f \notin \Phi BV$.

To see that $g \circ f \notin \Phi BV$ we observe that

$$\begin{aligned} \sum_{m=1}^{k_n} \varphi(|(g \circ f)(I_{n,m})|) &= \sum_{m=1}^{k_n} \varphi(|g(J_n)|) \\ &= k_n \varphi(|g(J_n)|) \\ &\geq k_n \varphi(c_n |J_n|) \\ &> \frac{\varphi(c_n |J_n|)}{\varphi(c_n |J_n|)} \\ &= 1. \end{aligned}$$

So for this collection $\{I_{n,m}\}$ of intervals, we have

$$\begin{aligned} \sum_{n=1}^{\infty} \sum_{m=1}^{k_n} \varphi(|(g \circ f)(I_{n,m})|) &\geq \sum_{n=1}^{\infty} 1 \\ &= \infty, \end{aligned}$$

implying $g \circ f \notin \Phi BV$.

We now show that $f \in \Phi BV$. Let $\{0 = x_0, x_1, \dots, x_m = 1\}$ be an arbitrary partition of $[0, 1]$. We will find an upper bound for the sum

$$\sum_{i=1}^m \varphi(|f(x_i) - f(x_{i-1})|).$$

Consider the contribution of an arbitrary term, $\varphi(|f(x_i) - f(x_{i-1})|)$. For $i = 1$ we have

$$\varphi(|f(x_1) - f(0)|) = \varphi(|f(x_1)|) \leq \varphi(q_1)$$

and similarly for $i = m$ we get $\varphi(|0 - f(x_{m-1})|) \leq \varphi(q_1)$. For $i \neq 1, m$ we have the following cases:

Case 1. There exists n such that $f(x_i), f(x_{i-1}) \in J_n$. If $I_{n,m} = [a_{n,m}, b_{n,m}]$, then $x_i, x_{i-1} \in [a_{n,1}, b_{n,k_n}]$. Clearly,

$$\varphi(|f(x_i) - f(x_{i-1})|) < \varphi(|J_n|).$$

We observe that increasing the number of partition points along intervals of monotonicity does not increase the sum, and hence the contribution of all such terms is bounded by $2k_n\varphi(|J_n|)$.

Case 2. There does not exist an n such that $f(x_i), f(x_{i-1}) \in J_n$. Then there is a smallest integer j satisfying $f(x_i) \leq q_j$.

Suppose there are r such intervals having this same smallest integer j satisfying the above. If these intervals are

$$[x_{i_0}, x_{i_0+1}], [x_{i_0+1}, x_{i_0+2}], \dots, [x_{i_0+r-1}, x_{i_0+r}],$$

we then have, since φ is convex,

$$\begin{aligned} \sum_{l=1}^r \varphi(|f(x_{i_0+l}) - f(x_{i_0+l-1})|) &\leq \varphi(|f(x_{i_0+r}) - f(x_{i_0})|) \\ &\leq \varphi(q_j). \end{aligned}$$

We observe that the above situation can arise at most once for each index j , $j = 1, \dots, m$.

By combining all of the above information we show that an upper bound for $\sum_1^m \varphi(|f(x_i) - f(x_{i-1})|)$ is

$$\begin{aligned} M &= 2\varphi(q_1) + \sum_{n=1}^{\infty} 2k_n\varphi(|J_n|) + \sum_{n=1}^{\infty} \varphi(q_n) \\ &= \text{I} + \text{II} + \text{III}. \end{aligned}$$

Clearly I, II, and III are independent of the choice of the partition. Obviously I is finite and q_n was chosen so that III is finite. Since $c_n \geq 1$, it follows that

$\varphi(c_n|J_n|) > c_n\varphi(|J_n|)$. Thus

$$\begin{aligned} \sum_{n=1}^{\infty} k_n \varphi(|J_n|) &= \sum_{n=1}^{\infty} [P(n) + 1] \varphi(|J_n|) \\ &\leq \sum_{n=1}^{\infty} \left(\frac{1}{\varphi(c_n|J_n|)} + 1 \right) \varphi(|J_n|) \\ &\leq \sum_{n=1}^{\infty} \left(\frac{\varphi(|J_n|)}{c_n \varphi(|J_n|)} + \varphi(|J_n|) \right) \\ &= \sum_{n=1}^{\infty} \frac{1}{c_n} + \sum_{n=1}^{\infty} \varphi(|J_n|) \\ &\leq \sum_{n=1}^{\infty} \frac{1}{c_n} + \sum_{n=1}^{\infty} \varphi(q_n) < \infty. \end{aligned}$$

Thus $f \in \Phi BV$, and the proof is complete.

REFERENCES

- [CW] M. Chaika and D. Waterman, *On the invariance of certain classes of functions under composition*, Proc. Amer. Math. Soc. **43** (1974), 345–348. MR **48**:8704
- [G] C. Goffman, *Everywhere convergence of Fourier series*, Indiana Univ. Math. J. **20** (1970), 107–112. MR **42**:4941
- [GW] C. Goffman and D. Waterman, *Functions whose Fourier series converge for every change of variable*, Proc. Amer. Math. Soc. **19** (1968), 80–86. MR **36**:4245
- [GS] A. Garsia and S. Sawyer, *On some classes of continuous functions with convergent Fourier series*, J. of Math. and Mech. **13** (1964), 586–601. MR **33**:7777
- [J] M. Josephy, *Composing functions of bounded variation*, Proc. Amer. Math. Soc. **83** (1981), 354–356. MR **83c**:26009
- [MO] J. Musielak and W. Orlicz, *On generalized variations I*, Studia Math. **18** (1959), 11–41. MR **21**:3524
- [S] R. Salem, *Sur un test général pour la convergence uniforme des séries de Fourier*, Comptes Rend. Acad. Sci. Paris v. **207** (1938), 662–664.
- [W1] D. Waterman, *On convergence of Fourier series of functions of generalized bounded variation*, Studia Math. **44** (1972), 107–117. MR **46**:9623
- [W2] D. Waterman, *On Λ -bounded variation*, Studia Math. **57** (1976), 33–45. MR **54**:5408
- [W3] D. Waterman, *On the summability of Fourier series of functions of Λ -bounded variation*, Studia Math. **55** (1976), 87–95. MR **53**:6212
- [W4] D. Waterman, *Fourier series of functions of Λ -bounded variation*, Proc. Amer. Math. Soc. **74** (1979), 119–123. MR **80j**:42010
- [Y] L. C. Young, *Sur une généralisation de la notion de variation de puissance p -ième bornée au sens de M. Wiener, et sur la convergence des séries de Fourier*, Comptes Rend. Acad. Sci. Paris **204** (1937), 470–472.

DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCE, THE COLLEGE OF WOOSTER,
WOOSTER, OHIO 44691

E-mail address: ppierce@acs.wooster.edu

DEPARTMENT OF MATHEMATICS, FLORIDA ATLANTIC UNIVERSITY, BOCA RATON, FLORIDA
33431-0991

E-mail address: fourier@earthlink.net