ZEROS OF FUNCTIONS WITH FINITE DIRICHLET INTEGRAL

STEFAN RICHTER, WILLIAM T. ROSS, AND CARL SUNDBERG

(Communicated by Juha M. Heinonen)

ABSTRACT. In this paper, we refine a result of Nagel, Rudin, and Shapiro (1982) concerning the zeros of holomorphic functions on the unit disk with finite Dirichlet integral.

This is a remark about the zeros of functions $f = \sum_{n \geq 0} a_n z^n$ holomorphic on the unit disk $U = \{z \in \mathbb{C} : |z| < 1\}$ that have finite *Dirichlet integral*

$$D(f) := \frac{1}{\pi} \int_U |f'|^2 dA = \sum_{n=0}^{\infty} n |a_n|^2,$$

where dA is Lebesgue measure in the plane. Clearly such functions belong to the classical Hardy space H^2 , and so the zeros $(z_n)_{n \geq 1} \subset U$ of f (repeated according to multiplicity) satisfy the Blaschke condition $\sum_{n \geq 0} (1 - |z_n|^2) < \infty$ [4, p. 18]. However, not every Blaschke sequence are the zeros of a holomorphic f with $D(f) < \infty$ [2].

In 1962, Shapiro and Shields [6] improved a result of Carleson [3] and showed that if

$$(1) \quad \sum_{n=1}^{\infty} \frac{1}{-\log(1 - |z_n|)} < \infty,$$

then there is a nontrivial holomorphic f on U with $D(f) < \infty$ such that $f(z_n) = 0$ for all n .

This condition does not completely characterize the zero sets of analytic functions with finite Dirichlet integral. For example, if $(z_n)_{n \geq 0} \subset (0, 1)$ is a Blaschke sequence for which (1) fails, then $f = (1 - z)^2 B$ has finite Dirichlet integral, where B is the Blaschke product with zeros $(z_n)_{n \geq 0}$. Nevertheless, in the converse direction, Nagel, Rudin, and Shapiro [5] proved that if $(r_n)_{n \geq 0} \subset (0, 1)$ is such that

$$\sum_{n=0}^{\infty} \frac{1}{-\log(1 - r_n)} = \infty,$$

then there is a sequence of angles $(\theta_n)_{n \geq 0}$ such that the sequence $(r_n e^{i\theta_n})_{n \geq 0}$ is not the zeros of any nontrivial holomorphic function f on U with $D(f) < \infty$. They do this by first noting that when $D(f) < \infty$, the limit

$$\lim_{z \rightarrow e^{i\theta}, z \in \Omega_{e^{i\theta}}} f(z)$$

Received by the editors October 22, 2002 and, in revised form, May 6, 2003.
 2000 *Mathematics Subject Classification*. Primary 30C15; Secondary 30C85.

exists for almost every $e^{i\theta}$, where $\Omega_{e^{i\theta}}$ is the *exponential* contact region

$$\Omega_{e^{i\theta}} := \{re^{i\phi} : 1 - r^2 > e^{-\frac{1}{|\theta-\phi|}}\}.$$

Beginning at $z = 1$, lay down arcs $I_n \subset \partial U$ of length

$$\frac{1}{-\log(1 - r_n)}$$

end-to-end (repeatedly traversing the unit circle). Since $\sum_{n \geq 1} |I_n| = \infty$, by hypothesis, each $e^{i\theta} \in \partial U$ will be contained in infinitely many of the intervals $(I_n)_{n \geq 0}$. Let $e^{i\theta_n}$ be the center of the interval I_n , and note that simple geometry shows that for every $e^{i\theta}$, the exponential contact region $\Omega_{e^{i\theta}}$ contains infinitely many of the points $r_n e^{i\theta_n}$. Thus if f has finite Dirichlet integral and $f(r_n e^{i\theta_n}) = 0$ for all n , the above limit result says that the boundary function for f will vanish almost everywhere on ∂U , forcing f to be identically zero. This argument actually shows that the sequence $(r_n e^{i\theta_n})_{n \geq 0}$ cannot be the zeros of a nontrivial *harmonic* function f on U with finite Dirichlet integral (where $|f'|^2$ is replaced by $|\nabla f|^2$ in the definition of the Dirichlet integral).

In this note, we refine this result and show that for analytic functions the angles θ_n can be chosen so that the zeros $(r_n e^{i\theta_n})_{n \geq 0}$ need not accumulate at every point of the circle, but instead accumulate at a *single* point.

Theorem 2. *Suppose $(r_n)_{n \geq 0} \subset (0, 1)$ with $r_n \rightarrow 1$ and*

$$\sum_{n=0}^{\infty} \frac{1}{-\log(1 - r_n)} = \infty.$$

Then there are angles $(\theta_n)_{n \geq 0}$ such that $\text{clos}(r_n e^{i\theta_n})_{n \geq 0} \cap \partial U = \{1\}$ and such that if f is holomorphic on U with $D(f) < \infty$ and $f(r_n e^{i\theta_n}) = 0$ for all n , then f is identically the zero function.

Our proof is based on the following lemma. In order to make our construction easier, we will work in the upper half plane.

Lemma 3. *Let $J \subset \mathbb{R}$ be a finite open interval with center x_0 and $0 < y_0 < |J|$. Set*

$$S := \{x + iy : x \in J, 0 < y < |J|\}$$

and $\lambda_0 := x_0 + iy_0$. Suppose f is holomorphic on S with

$$\int_S |f'|^2 dA < \infty$$

and $f(\lambda_0) = 0$. If

$$E = \{x \in J : |f(x)| \geq 1\},$$

and $|E| \geq \frac{1}{2}|J|$, then

$$\int_S |f'|^2 dA \geq \frac{c}{\log(|J|/y_0)},$$

where c is a universal constant.

Proof. Elementary considerations show that $\omega_{\lambda_0}^S(E)$, the harmonic measure of E with respect to S at λ_0 , is bounded below by a universal constant times $y_0/|J|$. Indeed,

$$\omega_{\lambda_0}^S(E) \geq \omega_{\lambda_0}^S(F),$$

where F is the union of two intervals in the real line of length $\frac{1}{4}|J|$ located at the lower corners of S . Let $\psi : S \rightarrow U$ be the conformal map that takes the centroid of S to the origin and the line segment containing λ_0 and x_0 to the positive real axis. Thus $\psi(x_0) = 1$ and $\psi(\lambda_0) = r$ with $1 - r \asymp y_0/|J|$. Then

$$\omega_{\lambda_0}^S(F) = |\psi(F)| \asymp \omega_r^U(\psi(F)) = \int_{\psi(F)} \frac{1-r^2}{|e^{it}-r|^2} dt.$$

But since $\psi(F)$ is a fixed distance from the point $z = 1$, the denominator in the above integral does not matter. Thus, since the measure of $\psi(F)$ is fixed, the above integral is comparable to $1 - r \asymp y_0/|J|$.

Let $\varphi : U \rightarrow S$ be the conformal map with $\varphi(0) = \lambda_0$, and let $g := f \circ \varphi$. Then $g(0) = 0$, $|g| \geq 1$ on $\varphi^{-1}(E)$, and $|\varphi^{-1}(E)| = \omega_{\lambda_0}^S(E) \geq cy_0/|J|$. This means that g is a “test function” for the logarithmic capacity of $\varphi^{-1}(E)$ [1, Theorem 2], and so

$$\int_U |g'|^2 dA \geq c \operatorname{cap}(\varphi^{-1}(E)) \geq \frac{c}{\log(|J|/y_0)}.$$

Here we are using the well-known fact that if $W \subset \partial U$ with $a = |W|$, then

$$\operatorname{cap}(W) \geq \frac{c}{\log(1/a)}.$$

Finally, note that

$$\int_U |g'|^2 dA = \int_S |f'|^2 dA.$$

□

We are now ready to prove our main theorem. To make the construction easier, we work in the upper half plane and replace the sequence $(r_n)_{n \geq 0}$ with a sequence $(y_n)_{n \geq 0} \subset (0, 1)$ with $y_n \rightarrow 0$ and such that

$$\sum_{n=0}^{\infty} \frac{1}{\log(1/y_n)} = \infty.$$

We will construct a sequence $(x_n + iy_n)_{n \geq 0}$ in the upper half plane whose closure intersects the real axis only at $x = 0$ and such that the only holomorphic function f in the upper half plane with finite Dirichlet integral for which $f(x_n + iy_n) = 0$ for all n is the zero function.

Assuming that $y_n \searrow 0$, we can find

$$1 \leq n_1 < m_1 < n_2 < m_2 < \cdots$$

such that, whenever $n \geq n_k$,

$$y_n \log \frac{1}{y_n} < \frac{1}{k^2} e^{-2k^2}$$

and

$$ke^{2k^2} < \sum_{n=n_k}^{m_k} \frac{1}{\log(1/y_n)} < ke^{2k^2} + 1.$$

For each k , lay out intervals

$$J_{n_k}, J_{n_k+1}, \dots, J_{m_k}$$

on the real axis end-to-end starting at $x = 0$ and such that

$$|J_n| = \frac{1}{k^2 e^{2k^2} \log(1/y_n)}, \quad n_k \leq n \leq m_k.$$

Then

$$\begin{aligned} \log \frac{|J_n|}{y_n} &= \log \frac{1}{k^2 e^{2k^2} y_n \log(1/y_n)} \\ &= \log(1/y_n) - \log k^2 - 2k^2 - \log \log(1/y_n) \\ &< \log(1/y_n). \end{aligned}$$

Let x_n be the center of J_n , and set $\lambda_n := x_n + iy_n$ and

$$S_n := \{x + iy : x \in J_n, 0 < y < |J_n|\}.$$

Suppose that f is holomorphic on the upper half plane with finite Dirichlet integral and such that $f(\lambda_n) = 0$ for all $n_k \leq n \leq m_k$. Set

$$\begin{aligned} A_k &:= \{n : n_k \leq n \leq m_k \text{ and } |f| \geq e^{-k^2} \text{ on a set } E_n \subset J_n \text{ with } |E_n| \geq \tfrac{1}{2}|J_n|\}, \\ B_k &:= \{n : n_k \leq n \leq m_k, n \notin A_k\}. \end{aligned}$$

Apply Lemma 3 to see that if $n \in A_k$, then

$$\int_{S_n} |f'|^2 dA \geq ce^{-2k^2} \frac{1}{\log(|J_n|/y_n)} \geq c \frac{e^{-2k^2}}{\log(1/y_n)},$$

and if $n \in B_k$, then

$$\int_{J_n} \log \frac{1}{|f|} dx \geq \frac{1}{2} |J_n| k^2 = \frac{1}{2} \frac{e^{-2k^2}}{\log(1/y_n)}.$$

We conclude that

$$\sum_{n \in A_k} \int_{S_n} |f'|^2 dA + \sum_{n \in B_k} \int_{J_n} \log \frac{1}{|f|} dx \geq ce^{-2k^2} \sum_{n=n_k}^{m_k} \frac{1}{\log(1/y_n)} \geq ck.$$

Thus by the log-integrability of f on the boundary [4, p. 17], f must be the zero function.

It follows that the set

$$\bigcup_{k=1}^{\infty} (\lambda_n)_{n_k \leq n \leq m_k}$$

cannot be the zeros of a holomorphic function with finite Dirichlet integral. Choose the remaining points (from the unused y_n 's) on the imaginary axis to obtain a sequence $(\lambda_n)_{n \geq 0}$ that is not the zero set of a function with finite Dirichlet integral. Finally, since

$$\sum_{n=n_k}^{m_k} |J_n| = \frac{1}{k^2 e^{2k^2}} \sum_{n=n_k}^{m_k} \frac{1}{\log(1/y_n)} < \frac{ke^{2k^2} + 1}{k^2 e^{2k^2}} \rightarrow 0, \quad k \rightarrow \infty,$$

it follows that the closure of the sequence $(\lambda_n)_{n \geq 0}$ intersects the real axis only at $x = 0$.

REFERENCES

- [1] T. Bagby, *Quasi topologies and rational approximation*, J. Functional Analysis **10** (1972), 259–268. MR **50**:7535
- [2] K. Bogdan, *On the zeros of functions with finite Dirichlet integral*, Kodai Math. J. **19** (1996), no. 1, 7–16. MR **96k**:30005
- [3] L. Carleson, *On the zeros of functions with bounded Dirichlet integrals*, Math. Z. **56** (1952), 289–295. MR **14**:458e
- [4] P. L. Duren, *Theory of H^p spaces*, Academic Press, New York, 1970. MR **42**:3552
- [5] A. Nagel, W. Rudin, and J. Shapiro, *Tangential boundary behavior of functions in Dirichlet-type spaces*, Ann. of Math. (2) **116** (1982), no. 2, 331–360. MR **84a**:31002
- [6] H. S. Shapiro and A. L. Shields, *On the zeros of functions with finite Dirichlet integral and some related function spaces*, Math. Z. **80** (1962), 217–229. MR **26**:2617

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF TENNESSEE, KNOXVILLE, TENNESSEE 37996
E-mail address: richter@math.utk.edu

DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCE, UNIVERSITY OF RICHMOND, RICHMOND, VIRGINIA 23173
E-mail address: wross@richmond.edu

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF TENNESSEE, KNOXVILLE, TENNESSEE 37996
E-mail address: sundberg@math.utk.edu