ERRATUM TO “SUPERCYCLIC AND CHAOTIC TRANSLATION SEMIGROUPS”

MAI MATSUI, MINO YAMADA, AND FUKIKO TAKEO

(Communicated by Joseph A. Ball)

In [1], the proof of Theorem 3 in Section 4 is incorrect, as pointed out to us by Alfredo Peris. In proving that “(2) implies (1)”, v_p has not been confirmed to belong to the space $C_{0,\rho}(I)$. As for the characterizations of chaoticity of the semigroup in the space $C_{0,\rho}(I)$, we give the following Theorem 3' instead of Theorems 3 and 4 in [1]. As the result, Example 2 in [1] does not give a chaotic translation semigroup.

Theorem 3'. *Let I be the interval $(-\infty, \infty)$ (resp. $I = [0, \infty)$), and let X be $C_{0,\rho}(I)$. For the translation semigroup $\{T(t)\}$ on X , the following assertions are equivalent:*

- (1) $\lim_{\tau \rightarrow \pm\infty} \rho(\tau) = 0$ (resp. $\tau \rightarrow \infty$);
- (2) $\{T(t)\}$ is chaotic.

Proof. (1) implies (2): It is clear that $\{T(t)\}$ is hypercyclic by Theorem A in [1]. So we only have to show that the set X_{per} of periodic points is dense in X . Since the set $X_{0,0}$ of all the functions with compact support is dense in X , we shall show that for any $z \in X_{0,0}$ with $\text{supp}(z) \subset [-l, l]$ and $\varepsilon > 0$, there exists $v \in X_{per}$ such that $\|z - v_p\| < \varepsilon$.

From Lemma B in [1], there exists $M_{2l} \geq 1$ satisfying

$$\frac{1}{M_{2l}} \rho(\sigma - l) \leq \rho(\tau) \leq M_{2l} \rho(\sigma + l)$$

for any $\sigma \in I$ and any $\tau \in [\sigma - l, \sigma + l]$.

Take ε' such as $0 < \varepsilon' < \frac{\rho(-l)}{M_{2l}\|z\|} \varepsilon$. By (1), there exists τ_0 such that $\rho(\tau) < \varepsilon'$ for any $|\tau| > \tau_0$. Let P satisfy $P > \max\{\tau_0 + l, 2l\}$. We shall put $v_p(\tau) = \sum_{n \in \mathbf{Z}} z(\tau - nP)$. Then clearly $T(P)v_p = v_p$ and $\rho(l + nP) < \varepsilon'$ for $n \in \mathbf{Z} \setminus \{0\}$. We calculate $\|z - v_p\|$:

$$\begin{aligned} \|z - v_p\| &= \left\| \sum_{n \in \mathbf{Z} \setminus \{0\}} z(\tau - nP) \right\| \leq \sup_{n \in \mathbf{Z} \setminus \{0\}} \sup_{\tau \in [-l+nP, l+nP]} \rho(\tau) \cdot |z(\tau - nP)| \\ &\leq \sup_{n \in \mathbf{Z} \setminus \{0\}} \frac{M_{2l}^2 \rho(l + nP)}{\rho(-l)} \cdot \|z\| < \varepsilon. \end{aligned}$$

Received by the editors December 3, 2003.

2000 *Mathematics Subject Classification.* Primary 47D60; Secondary 47D06.

Key words and phrases. Admissible weight function, supercyclic semigroups, chaotic semigroups.

(In the case when $I = [0, \infty)$, replace $-l$, $2l$, $\mathbf{Z}$ and $\mathbf{Z} \setminus \{0\}$ with 0 , l , $\mathbf{Z}_+$ and $\mathbf{N}$.) Since $v_p \in C_{0,\rho}(I)$ follows from (1) and v_p belongs to X_{per} , the set X_{per} is dense in X . Therefore $\{T(t)\}$ is chaotic.

(2) implies (1): We shall consider the case $I = [0, \infty)$. Let $z \in X$ satisfy $z(0) = 1$ and $z(\tau) = 0$ for $\tau \geq 1$.

Since $\{T(t)\}$ is chaotic, there exists $v \in X_{per}$ such that $\|z - v\| < \rho(0)/2$ and $P > 0$ such that $v(\tau) = v(\tau + nP)$ for any $n \in \mathbf{N}$. Then

$$v(0)\rho(0) \geq z(0)\rho(0) - \rho(0)/2 \geq \rho(0) - \rho(0)/2 = \rho(0)/2.$$

Therefore $v(0) \geq \frac{1}{2}$.

By Lemma B in [1], there exists M_P satisfying

$$(*) \quad \frac{1}{M_P} \rho(nP) \leq \rho(\tau) \leq M_P \rho((n+1)P)$$

for any $n \in \mathbf{N}$ and any $\tau \in [nP, (n+1)P]$.

Since v belongs to X , $\lim_{\tau \rightarrow \infty} v(\tau)\rho(\tau) = 0$ holds. So for any $\varepsilon > 0$, putting $\varepsilon_1 = \frac{\varepsilon}{2M_P}$, there exists τ_0 such that $|v(\tau)\rho(\tau)| < \varepsilon_1$ for $\tau > \tau_0$. If $|nP| > \tau_0$, then $|v(nP)\rho(nP)| < \varepsilon_1$ holds. By the relation $v(0) = v(nP)$, we have

$$\rho(nP) < \frac{\varepsilon_1}{|v(0)|} < 2\varepsilon_1.$$

For $\tau > \tau_0$, there exists $n \in \mathbf{Z}$ such that $nP \leq \tau < (n+1)P$. By (*), we have

$$\rho(\tau) < M_P \rho((n+1)P) < M_P \cdot 2\varepsilon_1 = \varepsilon,$$

which implies that $\lim_{\tau \rightarrow \infty} \rho(\tau) = 0$. In the case when $I = (-\infty, \infty)$, we can prove the theorem in a similar way. $\square$

ACKNOWLEDGEMENT

The authors would like to thank Alfredo Peris for pointing out the mistake in [1].

REFERENCES

1. Mai Matsui, Mino Yamada and Fukiko Takeo, *Supercyclic and chaotic translation semigroups*, Proc. Amer. Math. Soc. **131** (2003), 3535–3546. MR 2004c:47018

DOCTORAL RESEARCH COURSE IN HUMAN CULTURE, OCHANOMIZU UNIVERSITY, 2-1-1 OTSUKA, BUNKYO-KU, TOKYO 112-8610, JAPAN

E-mail address: mai@tke.att.ne.jp

DEPARTMENT OF INFORMATION SCIENCES, OCHANOMIZU UNIVERSITY, 2-1-1 OTSUKA, BUNKYO-KU, TOKYO 112, JAPAN

DEPARTMENT OF INFORMATION SCIENCES, OCHANOMIZU UNIVERSITY, 2-1-1 OTSUKA, BUNKYO-KU, TOKYO 112, JAPAN

E-mail address: takeo@is.ocha.ac.jp