THE VANISHING OF $\mathrm{Tor}_1^R(R^+, k)$ IMPLIES THAT R IS REGULAR

IAN M. ABERBACH

(Communicated by Bernd Ulrich)

ABSTRACT. Let (R, m, k) be an excellent local ring of positive prime characteristic. We show that if $\mathrm{Tor}_1^R(R^+, k) = 0$, then R is regular. This improves a result of Schoutens, in which the additional hypothesis that R was an isolated singularity was required for the proof.

Let R be an integral domain. Then we denote by R^+ the integral closure of R in an algebraic closure of the fraction field of R . Under the assumption that R is a local excellent domain with positive prime characteristic p , the ring R^+ is a balanced big Cohen-Macaulay algebra [3]. We assume for the rest of this paper that R is a commutative ring with positive prime characteristic p . Let $F : R \rightarrow R$ be the Frobenius endomorphism given by $r \mapsto r^p$. It is a theorem of Kunz [6] that R is regular if and only if F is a flat map. From this theorem it is not difficult to show that R is regular if and only if R^+ is flat over R . The more general question of whether $\mathrm{Tor}_1^R(R^+, k) = 0$ implies that R is regular for a local ring $(R, \mathfrak{m}, k)$ of positive characteristic is posed in the exercises in section 8 of [5]. (When $\mathrm{Tor}_1^R(S, k) = 0$ for a module-finite extension, then Nakayama's lemma shows that S is flat over R ; however, R^+ is far from finitely generated over R .) Schoutens has shown that for an excellent local ring the condition $\mathrm{Tor}_1^R(R^+, k) = 0$ implies that R is weakly F -regular, and if R has an isolated singularity, then R is regular ([8], Theorems 1.3 and 1.1). We show here that, in fact, the vanishing of $\mathrm{Tor}_1^R(R^+, k)$ suffices to imply regularity for excellent rings of positive prime characteristic.

Assume that $(R, \mathfrak{m}, k)$ is a reduced excellent local ring. R is then approximately Gorenstein, and so there is a sequence of irreducible $\mathfrak{m}$ -primary ideals $\{I_t\}$ cofinal with the powers of $\mathfrak{m}$ (see [2]). By taking a subsequence we may assume that the sequence is non-increasing. Let u_t be an element of R representing the socle modulo I_t . Then the injective hull of the residue field is $E = E_R(k) = \varinjlim_t R/I_t$, and the image of u_t in E is the socle element u of E for all t . Moreover, because the sequence is non-increasing, we may assume that for all t there is an injection $R/I_t \hookrightarrow R/I_{t+1}$ sending $u_t + I_t \mapsto u_{t+1} + I_{t+1}$.

Recall that a ring R of positive prime characteristic is called *F-finite* if the Frobenius endomorphism is module-finite. Such rings are excellent [7]; thus if, in addition, R is reduced, then it is approximately Gorenstein. We will use q to denote

Received by the editors July 15, 2003 and, in revised form, September 18, 2003.

2000 *Mathematics Subject Classification.* Primary 13A35; Secondary 13H05.

Key words and phrases. Regular rings, characteristic p , absolute integral closure.

The author was partially supported by the National Security Agency. He also wishes to thank the referee for a careful reading of this paper and several corrections.

a power of p . Whenever R is reduced, there is a well-defined ring of q th roots of R , denoted $R^{1/q}$, that is a finitely generated R -module for some (equivalently, all) q precisely when R is F -finite. In this case we will write $R^{1/q} \cong R^{a_q} \oplus_R M_q$, where M_q is a module with no free R summands.

The characterization of the injective hull given above is very helpful in proving the next lemma, which shows how to compute the values of a_q in a special case. By $I^{[q]}$ we mean the ideal $(i^q : i \in I)$.

Lemma 1. *Let $(R, \mathfrak{m}, k)$ be a reduced, F -finite ring with perfect residue field k . Then $a_q = \lambda_R(R/(I_t^{[q]} : u_t^q))$ for all $t \gg 0$.*

Proof. This result is a special case of Corollary 2.8 of [1]. However, we give a proof here for the benefit of the reader. We will use the fact that over an approximately Gorenstein ring, a homomorphism $f : R \rightarrow M$, where M is finitely generated, has a splitting over R if and only if for all t , $f(u_t) \notin I_t M$ (see [2]).

Fix q , and write $R^{1/q} \cong R^{a_q} \oplus_R M_q$ as above. We first claim that for $t \gg 0$, $u_t M_q \subseteq I_t M_q$, since for any minimal generator of M_q , the map $R x \rightarrow M_q$ does not split, and hence, $x u_t \in I_t M_q$. The claim follows since M_q is a finitely generated R -module. We will also use the fact that if I is an $\mathfrak{m}$ -primary ideal, then $\lambda_R(R/I^{[q]}) = \lambda_R(R^{1/q}/IR^{1/q})$, since k is perfect.

Thus, for any $t \gg 0$, we have

$$\begin{aligned} \lambda(R/(I_t^{[q]} : u_t^q)) &= \lambda(R/I_t^{[q]}) - \lambda(R/(I_t, u_t)^{[q]}) \\ &= \lambda(R^{1/q}/I_t R^{1/q}) - \lambda(R^{1/q}/(I_t, u_t) R^{1/q}) \\ &= \lambda(R^{a_q}/I_t R^{a_q}) + \lambda(M_q/I_t M_q) \\ &\quad - (\lambda(R^{a_q}/(I_t, u_t) R^{a_q}) + \lambda(M_q/(I_t, u_t) M_q)) \\ &= a_q \cdot 1 + \lambda(M_q/I_t M_q) - \lambda(M_q/(I_t, u_t) M_q) = a_q, \end{aligned}$$

since $(I_t, u_t) M_q = I_t M_q$ (for $t \gg 0$). $\square$

We will need to pass to a Γ construction as described in [4], Section 6. We refer the reader to [4] for details. What we need to know is as follows. Let $(R, \mathfrak{m}, k)$ be a complete ring of characteristic p . Then $R \rightarrow R^\Gamma$ is a faithfully flat, purely inseparable extension, the maximal ideal of R^Γ is $\mathfrak{m} R^\Gamma$, and R^Γ is F -finite. Note that if $I \subseteq R$ is an irreducible $\mathfrak{m}$ -primary ideal of R , then IR^Γ is also an irreducible $\mathfrak{m} R^\Gamma$ -primary ideal of R^Γ . Moreover, if $E_R(R/\mathfrak{m}) = \varinjlim_t R/I_t$, then $E_{R^\Gamma}(R^\Gamma/\mathfrak{m} R^\Gamma) = E_R(R/\mathfrak{m}) \otimes_R R^\Gamma = \varinjlim_t R^\Gamma/I_t R^\Gamma$.

Our main theorem is

Theorem 2. *Let $(R, \mathfrak{m}, k)$ be an excellent local domain of positive prime characteristic. Suppose that $\text{Tor}_1(R^+, k) = 0$. Then R is regular.*

Proof. By [8], Theorem 1.2, the ring R is weakly F -regular, therefore a Cohen-Macaulay, normal domain. In particular, R is approximately Gorenstein. Also, $R \rightarrow R^+$ is cyclically pure. The assumption that $\text{Tor}_1(R^+, k) = 0$ and an induction on length shows that for any $\mathfrak{m}$ -primary ideal $I \subseteq R$ and element x we have $IR^+ :_{R^+} x = (I :_R x) R^+$.

We first claim that for all q and all t , $I_t^{[q]} :_R u_t^q \subseteq \mathfrak{m}^{[q]}$. To see this suppose that $v u_t^q \in I_t^{[q]}$. Taking q th roots shows that $v^{1/q} \in I_t R^+ :_{R^+} u_t = \mathfrak{m} R^+$, and hence

that $v \in (\mathfrak{m}^{[q]})^+ = \mathfrak{m}^{[q]}$ (by cyclic purity of R in R^+). This shows that for all q and for all t , $\lambda(R/(I_t^{[q]} : u_t^q)) \geq \lambda(R/\mathfrak{m}^{[q]})$, which is greater than or equal to q^d ([6]).

We consider $R \rightarrow \hat{R} \rightarrow (\hat{R})^\Gamma = S$ for any Gamma extension of $\hat{R}$. In particular, we may take Γ to be the empty set, in which case the residue field of S is perfect. Then by faithful flatness and the fact that the maximal ideal of S is $\mathfrak{m}S$, $\lambda_R(R/(I_t^{[q]} :_R u_t^q)) = \lambda_S(S/(I_t S^{[q]} :_S u_t^q))$. Since $u_t^q \notin I_t S^{[q]}$ for all t , the ring S is F -pure, and hence reduced. Thus by Lemma 1, for large enough t (depending on q), $\lambda_R(R/(I_t^{[q]} :_R u_t^q)) = a_q(S)$ is the number of S -free summands in $S^{1/q}$. Since S has perfect residue field, the rank of $S^{1/q}$ as an S -module is precisely q^d ; hence $a_q(S) \leq q^d$. We have now shown that $q^d \geq \lambda(R/(I_t^{[q]} : u_t^q)) \geq \lambda(R/\mathfrak{m}^{[q]}) \geq q^d$. Thus $\lambda(R/\mathfrak{m}^{[q]}) = q^d$ and R is regular [6]. $\square$

REFERENCES

1. I. M. Aberbach and F. Enescu, *The structure of F -pure rings*, preprint, 2003.
2. M. Hochster, *Cyclic purity versus purity in excellent Noetherian rings*, Trans. Amer. Math. Soc. **231** (1977), no. 2, 463–488. MR57:3111
3. M. Hochster and C. Huneke, *Infinite integral extensions and big Cohen-Macaulay algebras*, Ann. of Math. (2) **135** (1992), no. 1, 53–89. MR92m:13023
4. M. Hochster and C. Huneke, *F -regularity, test elements, and smooth base change*, Trans. Amer. Math. Soc. **346** (1994), no. 1, 1–62. MR95d:13007
5. Craig Huneke, *Tight closure and its applications*, With an appendix by Melvin Hochster. CBMS Regional Conference Series in Mathematics, **88**. Published for the Conference Board of the Mathematical Sciences, Washington, DC; by the American Mathematical Society, Providence, RI, 1996. x+137 pp. ISBN: 0-8218-0412-X. MR96m:13001
6. E. Kunz, *Characterizations of regular local rings for characteristic p* , Amer. J. Math. **91** (1969), 772–784. MR40:5609
7. E. Kunz, *On Noetherian rings of characteristic p* , Amer. J. Math. **98** (1976), no. 4, 999–1013. MR55:5612
8. H. Schoutens, *On the vanishing of Tor for the absolute integral closure*, J. Algebra **275** (2004), 567–574.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MISSOURI, COLUMBIA, MISSOURI 65211

E-mail address: aberbach@math.missouri.edu

URL: <http://www.math.missouri.edu/people/iaberbach.html>