THE OSC DOES NOT IMPLY THE SOSC FOR INFINITE ITERATED FUNCTION SYSTEMS

TOMASZ SZAREK AND STANISŁAW WĘDRYCHOWICZ

(Communicated by Michael Handel)

ABSTRACT. It is shown that every class of contracting similitudes $\{f_1, \dots, f_N\}$ on R^s satisfying the OSC and such that $\dim_H \mathcal{K}_0 < s$, where $\mathcal{K}_0$ denotes the corresponding fractal, can be extended to an infinite family of contracting similitudes which still satisfies the OSC but the SOSC does not hold.

1. NOTATION AND DEFINITIONS

We consider R^s ($s \geq 1$) with the Euclidean norm $|\cdot|$. By $\text{dist}(A, B)$ we denote the Hausdorff distance of two compact subsets A, B of R^s . Recall that a map $f : R^s \rightarrow R^s$ is called a *similitude* with Lipschitz constant $c \in (0, 1)$ if $|f(x) - f(y)| = c|x - y|$ for $x, y \in R^s$. Let I be a countable index set with at least two elements, and let $\{f_i : R^s \rightarrow R^s : i \in I\}$ be a collection of contracting similitudes – a so-called *ifs*. The class of maps $\{f_i : i \in I\}$ is said to satisfy the *Open Set Condition (OSC)* if there is a nonempty and bounded open set $U \subset R^s$ such that $f_i(U) \subset U$ for all i , and $f_i(U) \cap f_j(U) = \emptyset$ for $i \neq j$.

Put $I^* = \bigcup_{n \geq 1} I^n$ and for $\mathbf{i} \in I^n$, $\mathbf{i} = (i_1, \dots, i_n)$, $n \geq 1$, set

$$f_{\mathbf{i}} = f_{i_1} \circ \dots \circ f_{i_n}.$$

If $\mathbf{i} \in I^* \cup I^\infty$ and $n \geq 1$ does not exceed the length of $\mathbf{i}$, we denote by $\mathbf{i}_{|n}$ the word $(i_1, \dots, i_n)$. Let $\{f_i : i \in I\}$ satisfy the OSC with a set U . Set $X = \text{cl } U$. Observe now that given $\mathbf{i} \in I^\infty$, the compact sets $f_{\mathbf{i}_{|n}}(X)$, $n \geq 1$, are decreasing and their diameters converge to zero. This implies that the set

$$\pi(\mathbf{i}) = \bigcap_{n=0}^{\infty} f_{\mathbf{i}_{|n}}(X)$$

is a singleton and therefore this formula defines a map $\pi : I^\infty \rightarrow X$. The main object of our interest will be the set

$$\mathcal{K} = \pi(I^\infty) = \bigcup_{\mathbf{i} \in I^\infty} \bigcap_{n=0}^{\infty} f_{\mathbf{i}_{|n}}(X),$$

Received by the editors July 2, 2003.

2000 *Mathematics Subject Classification.* Primary 28A80; Secondary 28A78.

Key words and phrases. Fractal, open set condition, strong open set condition.

The first author's research was supported by the State Committee for Scientific Research Grant No. Z PO3A 031 25.

called the *attractor* or *fractal* for $\{f_i : i \in I\}$. We easily check that

$$\mathcal{K} = \bigcup_{i \in I} f_i(\mathcal{K}).$$

The *Strong Open Set Condition (SOSC)* holds if the set U in the definition of the OSC can be chosen in such a way that $\mathcal{K} \cap U \neq \emptyset$.

For every finite word $\mathbf{i} = (i_1, \dots, i_n)$ in the alphabet I we will use the following abbreviation:

$$c_{\mathbf{i}} = c_{i_1} \cdots c_{i_n}.$$

2. THE THEOREM

Let $U \subset R^s$ be an open set, and let $f_1, \dots, f_N$ be given. Assume that $\{f_1, \dots, f_N\}$ satisfies the OSC with the set U . A. Schief proved that if f_i , $i \in \{1, \dots, N\}$, are contracting similitudes, then the OSC implies the SOSC (see [6]). Y. Peres *et al.* showed that the above implication is valid for conformal contractions (see [5]). Recently, infinite families of contracting maps have been extensively studied (see for example [3]). However, the question (see Problem 7.5 in [3]) of whether the above implication holds in this case seems to be still open. The main aim of our note is to show that for infinite iterated function systems the OSC does not imply the SOSC.

Let $\mathcal{K}_0$ denote the attractor for $\{f_1, \dots, f_N\}$. Assume that $\dim_H \mathcal{K}_0 < s$. Hence $\sum_{i=1}^N c_i^s < 1$, where c_i denotes the Lipschitz constant of f_i , $i = 1, \dots, N$ (see [2], [4]). Without loss of generality we may assume that $c_i \leq c_1$ for all $i \in \{1, \dots, N\}$.

Let $(a_n)_{n \geq 0}$ be a strictly decreasing sequence of reals such that $1 < a_n \leq 2$ for all $n \in \mathbb{N}$.

Since $\mathcal{L}^s(\mathcal{K}_0) = 0$, where $\mathcal{L}^s$ is the Lebesgue measure in R^s , we have $U \setminus \mathcal{K}_0 \neq \emptyset$. Choose $x \in U \setminus \mathcal{K}_0$. Since $x \notin \mathcal{K}_0$, there exists $r > 0$ such that $f_{j_1} \circ \dots \circ f_{j_m}(B(x, 2r)) \cap B(x, 2r) = \emptyset$ for $j_1, \dots, j_m \in \{1, \dots, N\}$, $m \in \mathbb{N}$, and $B(x, 2r) \subset U$. We may also assume that $\text{dist}(B(x, 2r), \mathcal{K}_0) > r$.

The following lemma is crucial for our considerations.

Lemma. *Let $((x_n, r_n))_{n \geq 1}$ be a dense sequence in $\text{cl}U \times (0, r/2]$, and let $B_n = B(x_n, r_n)$ for $n \in \mathbb{N}$. Then there are contracting similitudes $f_{N+1}, f_{N+2}, \dots$ such that there exist sequences $(i_n)_{n \geq 0}$ of integers, $(U_n)_{n \geq 0}$ of open subsets of U and $(F_n)_{n \geq 0}$ of compact subsets of R^s satisfying:*

- (1) $(i_n)_{n \geq 0}$ is nondecreasing, $i_n \rightarrow \infty$ as $n \rightarrow \infty$;
- (2) $\{f_1, \dots, f_{i_n}\}$ satisfies the OSC with the set U_n , $\text{dist}(B(x, 2r), \mathcal{K}_n) > r$ and $\dim_H \mathcal{K}_n < s$, where $\mathcal{K}_n$ denotes the attractor for $\{f_1, \dots, f_{i_n}\}$, $n \in \mathbb{N} \cup \{0\}$;
- (3) $f_{\mathbf{j}}(B(x, 2r)) \cap B(x, 2r) = \emptyset$, where $\mathbf{j}$ is a finite word in the alphabet $\{1, \dots, i_n\}$, $n \in \mathbb{N} \cup \{0\}$;
- (4) $B(x, a_n r) \subset U_n$ for $n \in \mathbb{N} \cup \{0\}$;
- (5) $U_n = U_{n-1} \setminus F_n$, $f_{\mathbf{j}}^{-1}(F_n) \cap U_{n-1} \subset F_n$, where $\mathbf{j}$ is a finite word in the alphabet $\{1, \dots, i_n\}$, $n \in \mathbb{N}$;
- (6) $f_{i_n}(\mathcal{K}_{n-1}) \subset B_n$ if $B_n \cap \mathcal{K}_{n-1} \neq \emptyset$ for $n \in \mathbb{N}$.

Proof. We set $i_0 = N$, $U_0 = U$ and $F_0 = \emptyset$. Now assume that we are given $i_p, f_1, \dots, f_{i_p}$, F_p and U_p for some $p \in \mathbb{N} \cup \{0\}$. We will define i_{p+1} , $f_{i_{p+1}}$, F_{p+1} and U_{p+1} in such a way that conditions (1)–(6) will be satisfied. If $B_{p+1} \cap \mathcal{K}_p = \emptyset$, then we set $i_{p+1} = i_p$, $F_{p+1} = F_p$ and $U_{p+1} = U_p$. If $B_{p+1} \cap \mathcal{K}_p \neq \emptyset$, then $i_{p+1} = i_p + 1$. Set $r_0 = \text{dist}(B(x, 2r), \mathcal{K}_p)$. Obviously $r_0 > r$. Choose $\eta \in (0, r_0 - r)$

and set $\varepsilon = (1 - c_1)\eta$. $y \in \text{int}(B(x, a_p r) \setminus B(x, a_{p+1} r))$. Choose $\mathbf{i}$ to be a finite word in the alphabet $\{1, \dots, i_p\}$ such that $f_{\mathbf{i}}(y) \in B(x_0, \varepsilon)$, and let $\kappa > 0$ be such that $B(y, \kappa) \subset \text{int}(B(x, a_p r) \setminus B(x, a_{p+1} r))$ and $f_{\mathbf{i}}(B(y, \kappa)) \subset B(x_0, \varepsilon)$. Since $\mathcal{L}^s(f_{\mathbf{i}}(B(y, \kappa))) > 0$, $\mathcal{L}^s(\mathcal{K}_p) = 0$ and $\mathcal{K}_p$ is compact, there exist $y_0 \in B(y, \kappa)$ and $\kappa_0 < \kappa$ such that $\overline{B}(y_0, \kappa_0) \subset B(y, \kappa)$ and $f_{\mathbf{i}}(\overline{B}(y_0, \kappa_0)) \cap \mathcal{K}_p = \emptyset$, where $\overline{B}(y_0, \kappa_0)$ denotes the closed κ_0 -ball centered at y_0 . Set $W = f_{\mathbf{i}}(\overline{B}(y_0, \kappa_0))$. We see at once that

$$\#\{\mathbf{j} : \mathbf{j} \text{ is a finite word in the alphabet } \{1, \dots, i_p\} \text{ and } f_{\mathbf{j}}^{-1}(W) \cap \text{cl } U \neq \emptyset\} < \infty.$$

Moreover since $f_{\mathbf{i}}(y_0)$ does not belong to $\mathcal{K}_p$, it is not a fixed point for $f_{\mathbf{j}}$, where $\mathbf{j}$ is an arbitrary finite word in the alphabet $\{1, \dots, i_p\}$. Then, if we choose κ_0 small enough, we can also assume that $W \cap f_{\mathbf{j}}^{-1}(W) = \emptyset$ for all finite words $\mathbf{j}$. Define

$$F_{p+1} = \bigcup \left(f_{\mathbf{j}}^{-1}(W) \cap \text{cl } U \right),$$

where the above union is over all finite words in the alphabet $\{1, \dots, i_p\}$, and observe that by the above F_{p+1} is compact. Further, since $\mathcal{K}_p = \text{cl } \bigcup_{\mathbf{j}} \text{Fix}(f_{\mathbf{j}})$, where the union is taken over all finite words in the alphabet $\{1, \dots, i_p\}$ (see [1]), there exists $\mathbf{j}_0$ such that $\text{Fix}(f_{\mathbf{j}_0}) \in B(x_0, \varepsilon)$. Without loss of generality we may assume that $\text{diam } f_{\mathbf{j}_0}(U) < c_1 \kappa_0$ and $\sum_{i=1}^p c_i^s + c_{\mathbf{j}_0}^s < 1$. Define $f_{i_{p+1}} = f_{\mathbf{j}_0} + (f_{\mathbf{i}}(y_0) - \text{Fix}(f_{\mathbf{j}_0}))$ and $U_{p+1} = U_p \setminus F_{p+1}$.

We see at once that conditions (1), (3), (5) and (6) hold. Now we check condition (2). From the construction it follows that $f_i(U_{p+1}) \cap f_j(U_{p+1}) = \emptyset$ for $i, j \in \{1, \dots, i_{p+1}\}$, $i \neq j$. To finish the proof of the OSC we have to show that $f_i(U_{p+1}) \subset U_{p+1}$ for $i \in \{1, \dots, i_{p+1}\}$. We see at once that $f_{i_{p+1}}(U_{p+1}) \subset U_{p+1}$. Let $f_i(z) \notin U_{p+1}$ for some $z \in U_n$ and $i \in \{1, \dots, i_{p+1} - 1\}$. Then $f_i(z) \in F_{p+1}$ and hence $z \in f_i^{-1}(F_{p+1})$. Further, since $z \in f_i^{-1}(F_{p+1}) \cap U_p \subset F_{p+1}$, we obtain $z \notin U_{p+1} = U_p \setminus F_{p+1}$, which is a contradiction. From the construction it follows that

$$\text{dist}(\mathcal{K}_{p+1}, \mathcal{K}_p) \leq \varepsilon(1 - c_1)^{-1}$$

and consequently

$$\begin{aligned} \text{dist}(\mathcal{K}_{p+1}, B(x, 2r)) &\geq \text{dist}(\mathcal{K}_{p+1}, B(x, 2r)) - \text{dist}(\mathcal{K}_{p+1}, \mathcal{K}_p) \\ &\geq \text{dist}(\mathcal{K}_{p+1}, B(x, 2r)) - \varepsilon(1 - c_1)^{-1} \\ &= \text{dist}(\mathcal{K}_{p+1}, B(x, 2r)) - \eta > r, \end{aligned}$$

which finishes the proof of (2). To prove (4) assume that $B(x, a_{p+1} r) \setminus U_{p+1} \neq \emptyset$. We have $B(x, a_{p+1} r) \subset B(x, a_p r) \subset U_p$. Since $U_{p+1} = U_p \setminus F_{p+1}$, we have $B(x, a_{p+1} r) \cap F_{p+1} \neq \emptyset$. By (3) we see that if $f_{\mathbf{i}}(B(x, a_p r)) \cap f_{\mathbf{j}}(B(x, a_p r)) \neq \emptyset$, then $\mathbf{i} = \mathbf{j}$. Therefore if $f_{\mathbf{i}}^{-1}(W) \cap B(x, a_p r) \neq \emptyset$ and $f_{\mathbf{j}}^{-1}(W) \cap B(x, a_p r) \neq \emptyset$, we have $\mathbf{i} = \mathbf{j}$. From this it follows that there is a unique finite word $\mathbf{i}$ in the alphabet $\{1, \dots, i_{p+1}\}$ such that $f_{\mathbf{i}}^{-1}(W) \subset B(x, a_p r)$. Since $f_{\mathbf{i}}^{-1}(W) \subset B(x, a_p r) \setminus B(x, a_{p+1} r)$, we have $B(x, a_{p+1} r) \cap F_{p+1} = \emptyset$, which, in turn, is a contradiction. The proof of (4) is finished. $\square$

Theorem. *Every class of contracting similitudes $\{f_1, \dots, f_N\}$ on R^s satisfying the OSC and such that $\dim_H \mathcal{K}_0 < s$, where $\mathcal{K}_0$ denotes the corresponding fractal, can be extended to an infinite family of contracting similitudes which still satisfies the OSC but the SOSC does not hold.*

Proof. We show that an infinite ifs given by the above lemma will satisfy the OSC but will not satisfy the SOSC. Set $U_0 = U \setminus \text{cl} \bigcup_{i=1}^{\infty} F_i$. From conditions (4) and (5) it follows that $U_0 \neq \emptyset$. Observe that $f_k(U_0) \subset U_0$ for $k \in \mathbb{N}$. Indeed, for every k there exists $n_0 \in \mathbb{N}$ such that $i_{n_0} \geq k$. Then $f_k(U_{n_0}) \subset U_{n_0}$. Let $f_k(z) \notin U_0$ for some $z \in U_0$. Therefore there exists a sequence $(y_n)_{n \geq 1}$, $y_n \in \bigcup_{i=1}^{\infty} F_i$, $n \in \mathbb{N}$, such that $y_n \rightarrow f_k(z) \in U_{n_0} = U \setminus \bigcup_{i=1}^{n_0} F_i$. Hence we may assume that $y_n \notin F_j$ for $j = 1, \dots, n_0$. On the other hand, $f_k^{-1}(F_j) \cap U_0 \subset F_j$ for $j > n_0$ and

$$|f_k^{-1}(y_n) - z| = c_k^{-1}|y_n - f_k(z)| \rightarrow 0$$

as $n \rightarrow \infty$. Since $f_k^{-1}(y_n) \in \bigcup_{i=n_0+1}^{\infty} F_i$ for all large n , we have $z \in \text{cl} \bigcup_{i=n_0+1}^{\infty} F_i \subset \text{cl} \bigcup_{i=1}^{\infty} F_i$, which is a contradiction.

Now we show that $f_p(U_0) \cap f_q(U_0) = \emptyset$ for $p \neq q$. Namely, choose m such that $i_m > \max\{p, q\}$. Then we have $U_0 \subset U_m$ and $\{f_1, \dots, f_{i_m}\}$ satisfies the OSC with U_m .

Finally, we show that $\{f_1, f_2, \dots\}$ does not satisfy the SOSC. Let $\mathcal{K}$ be its attractor. It is obvious that $\bigcup_{i=1}^{\infty} \mathcal{K}_i \subset \mathcal{K} \subset \text{cl} \bigcup_{i=1}^{\infty} \mathcal{K}_i$. Suppose to the contrary that the SOSC holds. Then there exist $z \in \mathcal{K}$ and an open set V such that $z \in V$ and $f_i(V) \cap f_j(V) = \emptyset$ for $i \neq j$. However, since V is open and $z \in \mathcal{K}$, there exists $m \in \mathbb{N}$ such that $\mathcal{K}_m \cap V \neq \emptyset$. On the other hand, since $\mathcal{K}_m = \text{cl} \bigcup \text{Fix } f_{\mathbf{i}}$, where the union is over all finite words $\mathbf{i}$ in the alphabet $\{1, \dots, i_m\}$ (see [1]), without loss of generality we can assume that z is a fixed point of $f_{\mathbf{i}}$ for some finite word $\mathbf{i}$ in the alphabet $\{1, \dots, i_m\}$. Let $\tau > 0$ be such that $B(z, \tau) \subset V$. Then $f_{\mathbf{i}}(B(z, \tau)) = B(z, c_{\mathbf{i}}\tau)$. Further let $n \in \mathbb{N}$ be such that $i_{n+1} \geq i_m$ and $B_{n+1} = B(x_{n+1}, r_{n+1})$ satisfies $B_{n+1} \cap \mathcal{K}_m \neq \emptyset$ and $B_{n+1} \subset B(z, c_{\mathbf{i}}\tau)$. Then $B(z, c_{\mathbf{i}}\tau) \subset f_{\mathbf{i}}(V)$ and $f_{i_{n+1}}(\mathcal{K}_m) \subset B(z, c_{\mathbf{i}}\tau)$. Hence $f_{\mathbf{i}}(V) \cap f_{i_{n+1}}(V) \neq \emptyset$, which is a contradiction. $\square$

ACKNOWLEDGEMENT

We are grateful to an anonymous referee for remarks and suggestions which have improved the presentation of this paper.

REFERENCES

- [1] Hata M., *On the structure of self-similar sets*, Japan J. Appl. Math. 2 (1985), 381–414. MR 0839336 (87g:58080)
- [2] Hutchinson J., *Fractals and self-similarity*, Indiana Univ. Math. J. 30 (1981), 713–747. MR 0625600 (82h:49026)
- [3] Mauldin R. D. and Urbański M., *Dimension and measures in infinite iterated function systems*, Proc. London Math. Soc. (3) 73 (1996), 105–154. MR 1387085 (97c:28020)
- [4] Moran P. A. P., *Additive functions of intervals and Hausdorff measure*, Proc. Cambridge Philos. Soc. 42 (1946), 15–23. MR 0014397 (7:278f)
- [5] Peres Y., Rams M., Simon K., and Solomyak B., *Equivalence of positive Hausdorff measure and the open set condition for self-conformal sets*, Proc. Amer. Math. Soc. 129 (2001), 2689–2699. MR 1838793 (2002d:28004)
- [6] Schief A., *Separation properties for self-similar sets*, Proc. Amer. Math. Soc. 122 (1994), 111–115. MR 1191872 (94k:28012)

INSTITUTE OF MATHEMATICS, SILESIAN UNIVERSITY – AND – INSTITUTE OF MATHEMATICS OF THE POLISH ACADEMY OF SCIENCES, BANKOWA 14, 40–007 KATOWICE, POLAND

E-mail address: szarek@ux2.math.us.edu.pl

DEPARTMENT OF MATHEMATICS, TECHNICAL UNIVERSITY OF RZESZÓW, W. POLA 6, 35–959 RZESZÓW, POLAND