ON THE EVALUATION OF SALIÉ SUMS

ÁRPÁD TÓTH

(Communicated by David E. Rohrlich)

ABSTRACT. The Salié sum $S(m, n; c)$ can be evaluated as the product of a Gauss sum and an exponential sum involving square roots of $mn \bmod c$. We give a new proof of this fact that can simultaneously handle a twisted version of these sums that arise in the theory of half-integral weight modular forms.

The exponential sum

$$K(m, n; c) = \sum_{a\bar{a} \equiv 1(c)} \varepsilon_a \left(\frac{c}{a} \right) e((ma + n\bar{a})/c)$$

arises in the theory of modular forms of half-integral weight. $K(m, n; c)$ is only defined when $4|c$, and then $\varepsilon_a = 1$ or i according to whether $a \equiv 1$ or $3 \pmod{4}$, $(-)$ is the extension of the Legendre-Jacobi symbol as in [5], and $e(z) = \exp(2\pi iz)$. In Iwaniec's celebrated estimates for the Fourier coefficients of said forms (see [2]), an essential role is played by the identity

$$(1) \quad K(D, 1; c) = \frac{G(1, 0; c)}{2} \sum_{x^2 \equiv D(c)} e(2x/c)$$

valid when $8|c$ and the analogous formula

$$(2) \quad S(D, 1; c) = \sum_{a\bar{a} \equiv 1(c)} \left(\frac{a}{c} \right) e((Da + \bar{a})/c) = G(1, 0; c) \sum_{x^2 \equiv D(c)} e(2x/c),$$

which is valid whenever c is odd. In the formula above $G(a, b; c)$ is the Gauss sum

$$G(a, b; c) = \sum_{x(c)} e((ax^2 + bx)/c).$$

When the modulus is prime this identity was first proved by Salié [3]; see also [6]. Iwaniec derived a formula for the general modulus by pasting the local results together using quadratic reciprocity, while Sarnak gave a “global” proof of (2) that works when $(2D, c) = 1$; see [4].

The purpose of this note is to give a simple argument that leads to (1) when c is even and to (2) when c is odd, without any assumption on D . It is based on a very simple idea: we start by the sum

$$A = \sum_{x^2 \equiv D(c)} e(2x/c)$$

Received by the editors October 6, 2003.

2000 *Mathematics Subject Classification*. Primary 11L05; Secondary 11F37.

©2004 American Mathematical Society
 Reverts to public domain 28 years from publication

and sieve out the support of the sum by

$$A = \frac{1}{c} \sum_{x(c)} e(2x/c) \sum_{a(c)} e(a(x^2 - D)/c).$$

Interchanging the two sums, we are led to

$$A = \frac{1}{c} \sum_{a(c)} G(a, 2; c) e(-aD/c) = \sum_{d|c} A_d$$

where

$$A_d = \frac{1}{c} \sum_{\substack{a(c) \\ (a,c)=d}} G(a, 2; c) e(-aD/c).$$

It is well known that $G(a, b; c) = dG(a/d, b/d; c/d)$ if $d = (a, c)|b$ and is zero otherwise. This shows that $A_d = 0$ for all $d > 1$, when c is odd or $8|c$, because we even have $G(a', 1; c') = 0$ if $4|c'$.

Now, when $(a, c) = 1$,

$$G(a, 2; c) = e(-\bar{a}/c) G(a, 0; c),$$

and so

$$A = A_1 = \frac{G(1, 0; c)}{c} \sum_{\substack{a(c) \\ (a,c)=1}} \frac{G(a, 0; c)}{G(1, 0; c)} e(-\bar{a}/c) e(-aD/c).$$

This proves (1) and (2), since

$$(3) \quad G(a, 0; c)/G(1, 0; c) = \begin{cases} \left(\frac{a}{c}\right) & \text{if } c \text{ is odd,} \\ \left(\frac{c}{a}\right) \varepsilon_a^{-1} & \text{if } 4|c. \end{cases}$$

Since the explicit form of the θ -multiplier arises from $G(a, 0; c)/G(1, 0; c)$, see e.g. [4], identity (3) can be eliminated from the argument. In this respect, note that in [1], using the Davenport-Hasse relation, Duke evaluated generalized Salié sums that bear the same relation to metaplectic forms on GL_n as $K(m, n; c)$ to half-integral weight forms. Although the method presented here does not seem to be applicable in a direct fashion, this fact still suggests that Duke's theorem can be generalized to non-prime arguments, without any recourse to the explicit evaluation of generalized Gauss sums.

REFERENCES

- [1] Duke, W. On multiple Salié sums. *Proc. Amer. Math. Soc.* **114** (1992), no. 3, 623–625. MR1077785 (92f:11113)
- [2] Iwaniec, Henryk. Fourier coefficients of modular forms of half-integral weight. *Invent. Math.* **87** (1987), no. 2, 385–401. MR0870736 (88b:11024)
- [3] Salié, Hans. Über die Kloostermanschen Summen $S(u, v; q)$. (German) *Math. Z.* **34**, 91–109, (1931).
- [4] Sarnak, P. Some applications of modular forms. *Cambridge Tracts in Mathematics*, 99. *Cambridge University Press, Cambridge*, 1990. MR1102679 (92k:11045)

- [5] Shimura, Goro. On modular forms of half integral weight. *Ann. of Math. (2)* **97**, (1973), 440–481. MR0332663 (48:10989)
- [6] Williams, Kenneth S. Note on Salié's sum. *Proc. Amer. Math. Soc.* **30**, (1971), 393–394. MR0284408 (44:1635)

DEPARTMENT OF ANALYSIS, EÖTVÖS LÓRAND UNIVERSITY, PÁZMÁNY PÉTER SÉTÁNY 1/C,
H-1117 BUDAPEST, HUNGARY

E-mail address: toth@cs.elte.hu