COVERINGS BY CONVEX BODIES AND INSCRIBED BALLS

VLADIMIR KADETS

(Communicated by N. Tomczak-Jaegermann)

ABSTRACT. Let H be a Hilbert space. For a closed convex body A denote by $r(A)$ the supremum of the radii of balls contained in A . We prove that $\sum_{n=1}^{\infty} r(A_n) \geq r(A)$ for every covering of a convex closed body $A \subset H$ by a sequence of convex closed bodies A_n , $n \in \mathbb{N}$. It looks like this fact is new even for triangles in a 2-dimensional space.

1. INTRODUCTION

Recall that by a plank of width w in a Hilbert space one means a set of the form

$$P = \{h \in H : |\langle h - h_0, e \rangle| \leq \frac{w}{2}\},$$

where $\|e\| = 1$. According to T. Bang's theorem [3], if a sequence P_n of planks of widths w_n covers a ball of diameter w , then $\sum w_n \geq w$. (To be more precise this is a particular case of Bang's theorem. The complete statement includes a convex set P of minimal width w instead of a ball.)

K. Ball generalized Bang's theorem to coverings of a ball in a Banach space, for planks defined with the help of linear functionals instead of inner product.

Our work is inspired by Bang's and Ball's theorems. The idea was to find a "symmetric" generalization of these theorems, where the elements of the covering and the covered set are of the same nature, and they are measured "from inside". In this paper we prove such a generalization for sets in a Hilbert space. An analogous question for general Banach spaces remains open. In the proof we use ideas from Ball's exposition of Bang's theorem.

2. THE MAIN RESULT

For simplicity throughout the paper we consider real Hilbert spaces (finite- or infinite-dimensional). All the results are true for complex spaces too: the only thing which one must change in the proofs is the equation of a real hyperplane: in the complex case such an equation uses not the inner product itself, but the real part of the inner product.

Received by the editors November 6, 2003 and, in revised form, January 7, 2004.

2000 *Mathematics Subject Classification*. Primary 52A37; Secondary 46C05.

Key words and phrases. Hilbert space, convex sets, inscribed ball.

The author expresses thanks to the Department of Mathematics, University of Missouri-Columbia, and especially to Professor Nigel Kalton for hospitality and a fruitful working atmosphere.

©2004 American Mathematical Society
 Reverts to public domain 28 years from publication

Let H be a Hilbert space, and let B and S be its unit ball and its unit sphere respectively. By “body” in H we mean a closed subset with non-empty interior. For a convex body A denote by $r(A)$ the supremum of the radii of balls $x + aB$ contained in A .

If A is a ball of radius r , then $r(A) = r$. If A is a plank of width h , then $r(A) = \frac{h}{2}$.

Theorem 2.1. *If a convex body $A \subset H$ is covered by a sequence of convex bodies A_n , $n \in \mathbb{N}$, then $\sum_{n=1}^{\infty} r(A_n) \geq r(A)$.*

To prove the theorem we need first some lemmas. The goal of the lemmas is to prove that a convex bounded body in a Hilbert space can be approximated in some sense by a polytope with a finite number of faces.

Lemma 2.2. *Let $A \subset H$ be a bounded convex body. Then there is a ball $U \subset A$ with $r(U) = r(A)$.*

Proof. Denote $r(A) = r$. For every $p \in [0, r)$ consider the set A_p , consisting of all those $x \in A$ for which $x + pB \subset A$. Each A_p is a bounded convex closed set, and hence it is a weak compact set. Since $A_p \neq \emptyset$ and decreases as $p \rightarrow r$, there is a point $o_1 \in \bigcap_{p \in [0, r)} A_p$. Then $U = o_1 + rB$ will be the ball we need. $\square$

Lemma 2.3. *Let $A \subset H$ be a bounded convex body, $r(A) = r$, and let $U = o_1 + rB$ be the ball from the previous lemma. Then for every $\varepsilon > 0$ there is a finite subset $G = \{g_1, \dots, g_n\} \subset rB$ such that $(1 + \varepsilon)g_i + o_1 \in H \setminus A$ and*

$$(1) \quad \text{dist}(\text{conv}\{g_i\}_{i=1}^n, 0) \leq \varepsilon.$$

Proof. Without loss of generality, we may assume $o_1 = 0$ (otherwise shift the picture). We must prove that the closed convex hull of $(1 + \varepsilon)rB \setminus A$ contains 0.

Assume it is not so. By the Hahn-Banach theorem, there is a hyperplane P , strictly separating $(1 + \varepsilon)rB \setminus A$ from 0. This means that the set F - the bigger part of the ball $(1 + \varepsilon)rB$, lying on the same side of P as 0, is included in A . The subset $F \cup rB$ of A evidently contains a ball of radius bigger than r , which contradicts the definition of r . $\square$

Lemma 2.4. *Let $x, y \in H$, $\|x\| = 1 + \varepsilon$, and suppose the hyperplane*

$$\{h \in H : \langle h, y \rangle = \|y\|^2\}$$

generated by y separates x from the unit ball. Then $\|x - y\| \leq \delta(\varepsilon)$, where $\delta(\varepsilon)$ tends to 0 as ε tends to 0.

Proof. Our conditions mean that

$$\langle x, y \rangle \geq \|y\|^2$$

and

$$\|y\| = \sup_{h \in B} \langle h, y \rangle \leq \|y\|^2,$$

i.e. $\|y\| \geq 1$. We have

$$\|x - y\|^2 = \|x\|^2 - \|y\|^2 + 2(\|y\|^2 - \langle x, y \rangle) \leq \|x\|^2 - \|y\|^2 \leq (1 + \varepsilon)^2 - 1.$$

$\square$

Lemma 2.5. *Let $A \subset H$ be a bounded convex body, $r(A) = r$, and let $U = o_1 + rB$ be the ball from Lemma 2.2. Then for every $\varepsilon > 0$ there is a finite subset $V = \{v_1, \dots, v_n\} \subset rB$ (n depends on ε) such that*

$$(2) \quad A \subset W := \bigcap_{v \in V} \{h \in H : \langle h - o_1, v \rangle < r^2 + \varepsilon\}, \text{ and}$$

$$(3) \quad \text{dist}(\text{conv}V, 0) \leq \varepsilon.$$

Proof. Apply Lemma 2.3 for a small ε_1 to obtain the corresponding $G = \{g_1, \dots, g_n\} \subset rB$. Since $(1 + \varepsilon_1)g_i + o_1 \in H \setminus A$, one can separate $(1 + \varepsilon_1)g_i + o_1$ from A by a hyperplane

$$P_i = \{h \in H : \langle h - o_1, w_i \rangle = \|w_i\|^2\}.$$

Since $U = o_1 + rB \subset A$, P_i separates U from $(1 + \varepsilon_1)g_i + o_1$. This means that the hyperplane

$$\{h \in H : \langle h, w_i \rangle = \|w_i\|^2\}$$

separates rB from $(1 + \varepsilon_1)g_i$, and by Lemma 2.4, w_i is “very close” to g_i , and $\|w_i\|$ is “very close” to r . This means in turn that when ε_1 is small enough, the elements $v_k = r \frac{w_k}{\|w_k\|}$, $k = 1, 2, \dots, n$ fulfill the demands of this lemma. $\square$

Lemma 2.6. *For a convex body $A \subset B$ and for arbitrary $\delta > 0$ there is a polytope W of the form*

$$\bigcap_{k=1}^n \{h \in H : \langle h - o_1, v_k \rangle < a_k\},$$

such that $W \supset A$ and $r(W \cap B) \leq r(A) + \delta$.

Proof. The polytope W can be taken from (2) of the previous Lemma 2.5, with ε small enough. Let us show this. Consider $U = o_1 + rB$ from Lemma 2.2. For simplicity assume $o_1 = 0$ (the general case differs not too much from this one). Assume, contrary to our statement, that $W \cap B$ contains a ball of the form $U_0 = o_0 + (r + \delta)B$ (and automatically $\|o_0\| \leq 1$). Then $W \supset \text{conv}\{U_0, U\}$. According to condition (3), since $(1 + \varepsilon)V$ lies outside W ,

$$\text{dist}(\text{conv}((1 + \varepsilon)U \setminus W), 0) \leq (1 + \varepsilon)\varepsilon,$$

and hence

$$\text{dist}(\text{conv}((1 + \varepsilon)U \setminus \text{conv}\{U_0, U\}), 0) \leq (1 + \varepsilon)\varepsilon.$$

But for fixed δ and $\varepsilon \rightarrow 0$ the last inequality cannot be true, since the radius of U_0 equals $(r + \delta)$ and the distance between the centers of U_0 and U is bounded by a number, independent of ε . $\square$

Proof of Theorem 2.1. Denote $r(A_n) = r_n$. By the definition of $r(A)$ we must prove that $\sum_{n=1}^{\infty} r_n \geq r(U)$ for every ball $U \subset A$. By homogeneity this means that it is sufficient to prove the theorem for the case of A being the unit ball B of H . Also without loss of generality one may assume $A_n \subset B$: otherwise consider $A_n \cap B$ instead of A_n .

So the theorem can be reformulated as follows:

Let $\sum_{n=1}^{\infty} r_n < 1$, and let $A_n \subset B$ be convex bodies with $r(A_n) = r_n$. Then

$$B \setminus \bigcup_{n=1}^{\infty} A_n \neq \emptyset.$$

According to Lemma 2.6 every A_n may be included in the interior of a polytope (say, C_n), in such a way that the condition $\sum_{n=1}^{\infty} r(C_n \cap B) < 1$ still takes place. So, in fact, we may assume that the interiors of all of the A_n are relatively weakly open in B , and it is enough to prove that B cannot be covered by the interiors of the A_n . But in this case, since B is a weak compact set, it is sufficient to prove that B cannot be covered by the union of finitely many interiors of the A_n . So we reduced our theorem to the case when the number of sets is finite (say, equals N).

Let us fix a sequence of positive δ_n with

$$(4) \quad \sum_{n=1}^N (1 + \delta_n) r_n < 1,$$

and select $\varepsilon_n > 0$ with

$$(5) \quad (1 + \delta_n)(r_n^2 - 6N\varepsilon_n) \geq r_n^2 + \varepsilon_n.$$

For every A_n apply Lemma 2.5 with $\varepsilon = \varepsilon_n$ to get corresponding vectors $o_n \in A_n$, finite sets $V_n \subset r_n B$ and open polytopes

$$W_n := \bigcap_{v \in V_n} \{h \in H : \langle h - o_n, v \rangle < r_n^2 + \varepsilon_n\},$$

such that $A_n \subset W_n$ and

$$(6) \quad \text{dist}(\text{conv} V_n, 0) \leq \varepsilon_n.$$

We are going to prove that

$$(7) \quad B \setminus \bigcup_{n=1}^N W_n \neq \emptyset.$$

Let us introduce an auxiliary space $X = H \oplus H_1 \oplus H_2 \oplus \dots \oplus H_N$, the Hilbertian orthogonal direct sum of isometric copies of the original space H . Let $U_n : H \rightarrow H_n$ be the corresponding isometries.

Consider $K = V_1 \times V_2 \times \dots \times V_N$, and for every $\bar{g} = (g_1, g_2, \dots, g_N) \in K$ introduce

$$f(\bar{g}) = \sum_{n=1}^N ((1 + \delta_n)(g_n + U_n g_n) - U_n o_n).$$

Fix $\bar{x} = (x_1, x_2, \dots, x_N) \in K$ for which $\|f(\bar{x})\|$ is maximal. We are going to prove that

$$x = \sum_{n=1}^N (1 + \delta_n) x_n$$

is the element we need: namely, x belongs B but does not belong to any of W_j .

First of all, due to (4), $\|\sum_{n=1}^N (1 + \delta_n) x_n\| \leq \sum_{n=1}^N (1 + \delta_n) r_n < 1$, i.e. $x \in B$. Now for a fixed $j \in \{1, 2, \dots, N\}$ consider

$$y_j = f(\bar{x}) - (1 + \delta_j)(x_j + U_j x_j).$$

Taking into account that, according to our construction,

$$\|y_j + (1 + \delta_j)(x_j + U_j x_j)\| = \|f(\bar{x})\| \geq \|y_j + (1 + \delta_j)(v + U_j v)\|$$

for all $v \in V_j$, we obtain for all $v \in V_j$,

$$(8) \quad \langle y_j, (1 + \delta_j)(x_j + U_j x_j) \rangle \geq \langle y_j, (1 + \delta_j)(v + U_j v) \rangle.$$

Due to condition (6), there is a convex combination $\sum_{v \in V_j} \alpha_v v$, having norm less than ε_j . Multiplying (8) by α_v and adding for all $v \in V_j$ we obtain

$$\langle y_j, (1 + \delta_j)(x_j + U_j x_j) \rangle \geq -2\varepsilon_j(1 + \delta_j)\|y_j\| \geq -6N\varepsilon_j(1 + \delta_j).$$

Let us transform the last inequality using the definition of y_j and the pairwise orthogonality of the subspaces $H, H_1, \dots, H_N$:

$$\begin{aligned} & \left\langle x - (1 + \delta_j)x_j + (1 + \delta_j) \sum_{n \neq j} U_n x_n - \sum_{n=1}^N U_n o_n, (1 + \delta_j)(x_j + U_j x_j) \right\rangle \\ & \geq -6N\varepsilon_j(1 + \delta_j), \\ & (1 + \delta_j)\langle x, x_j \rangle - (1 + \delta_j)^2 r_j^2 - (1 + \delta_j)\langle U_j o_j, U_j x_j \rangle \geq -6N\varepsilon_j(1 + \delta_j), \\ & (1 + \delta_j)\langle x, x_j \rangle - (1 + \delta_j)\langle o_j, x_j \rangle \geq (1 + \delta_j)^2 r_j^2 - 6N\varepsilon_j(1 + \delta_j). \end{aligned}$$

By (5) we have

$$\langle x - o_j, x_j \rangle \geq r_j^2 + \varepsilon_j,$$

and this by definition of W_j means that x does not belong to W_j . $\square$

REFERENCES

- [1] Keith Ball, *Convex Geometry and Functional Analysis*, in W.B.Johnson and J.Lindenstrauss (editors) Handbook of the geometry of Banach spaces, vol. 1 (2001), 161 - 194. MR1863692 (2003c:52001)
- [2] Keith Ball, *The plank problem for symmetric bodies*, Invent. Math. **104** (1991), 535 - 543. MR1106748 (92c:52003)
- [3] T. Bang, *A solution of the "Plank problem"*, Proc. Amer. Math. Soc. **2** (1951), 990 - 993. MR0046672 (13:769a)

DEPARTMENT OF MECHANICS AND MATHEMATICS, KHARKOV NATIONAL UNIVERSITY, PL. SVOBODY 4, 61077 KHARKOV, UKRAINE

E-mail address: vova1kadets@yahoo.com

Current address: Department of Mathematics, University of Missouri, Columbia, Missouri 65211