SOME REMARKS ON AN EXISTENCE PROBLEM FOR DEGENERATE ELLIPTIC SYSTEMS

OLLI MARTIO, VLADIMIR MIKLYUKOV, AND MATTI VUORINEN

(Communicated by Richard A. Wentworth)

ABSTRACT. The system $au_x + bu_y = v_y$, $cu_x + du_y = -v_x$, which yields Beltrami's system if $b = c$, is considered. Under a condition for the coefficients a, b, c, d a non-existence theorem is proved.

1. MAIN RESULTS

Let $D, \mathcal{D} \subset \mathbf{R}^2$ be simply connected domains of the $z = (x, y)$ and the $w = (u, v)$ planes, respectively. Below we consider the problem of existence of homeomorphic ACL_{loc}^2 mappings $w = w(z)$ from D onto $\mathcal{D}$ satisfying the system

$$(1.1) \quad au_x + bu_y = v_y, \quad cu_x + du_y = -v_x.$$

A function h is, by definition, in the class ACL_{loc}^2 iff h is absolutely continuous along a.e. horizontal and vertical lines and its partial derivatives h_x, h_y belong to L_{loc}^2 . Here the coefficients a, b, c, d are measurable functions in D . We set

$$\delta \equiv ad - (b + c)^2/4.$$

In what follows we assume that $a > 0$ and

$$(1.2) \quad \text{ess inf}_{D'} (\delta) > 0 \quad \text{for every } D' \subset\subset D.$$

In the particular case $a = d = 1$ and $b = c = 0$, (1.1) is the classic Cauchy-Riemann system and the solution of the existence problem is given by the Riemann mapping theorem. In the case $b = c$, we have the well-known Beltrami system. For more on the existence theorems given here see, for example, G. David [5], M.A. Brakalova and J.A. Jenkins [2], U. Srebro and E. Yakubov [14], V. Gutlyanskii, O. Martio, T. Sugawa and M. Vuorinen [8], O. Martio and V. M. Miklyukov [10].

In the general case, the global mapping problem is much more complicated than in the aforementioned particular cases. Under the assumption

$$(1.3) \quad \text{ess inf}_D \delta > 0,$$

the existence problem was solved by B. Bojarski [1]. If this condition is not fulfilled, then there are isolated results only. The case in which (1.3) is violated at a finite number of boundary points was considered by A. Džuraev [6] and E.A. Chtcherbakov [4]. On the other hand, I.S. Ovchinnikov [13] and A.P. Mikhailov [12] proved some results pertaining to the solvability problem of the system

Received by the editors June 1, 2003 and, in revised form, January 22, 2004.
 2000 *Mathematics Subject Classification*. Primary 30C62; Secondary 35J70.

(1.1) with (1.2) under some special conditions, which allow degeneration close to a boundary arc.

Here we give a condition for the coefficients of (1.1) under which the coordinate function $u(z)$ of $w(z)$ is monotone close to the boundary [11]. We use this condition for a non-existence theorem.

Let $D \subset \mathbf{R}^2$ be a domain. By $\tilde{\partial}D$ we denote the boundary of D in the extended plane $\tilde{\mathbf{R}}^2 = \mathbf{R}^2 \cup \{\infty\}$. For an arbitrary subdomain $\Delta \subset D$, we set

$$\partial'\Delta = \tilde{\partial}\Delta \setminus \tilde{\partial}D \quad \text{and} \quad \partial''\Delta = \tilde{\partial}\Delta \cap \tilde{\partial}D.$$

Let Γ be a subset of $\tilde{\partial}D$. A continuous function $f : D \rightarrow \mathbf{R}$ is called *monotone close to* Γ if for every subdomain $\Delta \subset D$ with $\partial'\Delta \subset \Gamma$,

$$(1.4) \quad \text{osc}(f, \Delta) \leq \text{osc}(f, \partial'\Delta).$$

Here the symbol $\text{osc}(f, E)$ means the oscillation of f along the set E .

Every function f monotone close to Γ is monotone in the classical sense of Lebesgue since for domains $\Delta \subset\subset D$, (1.4) reduces to $\text{osc}(f, \Delta) \leq \text{osc}(f, \partial\Delta)$. On the other hand, if $\Gamma = \tilde{\partial}D$, then every function monotone close to Γ is a constant function. This is obvious since choosing $\Delta = D \setminus \{z_0\}$, where $z_0 \in D$ is an arbitrary point, by (1.4) we obtain

$$\text{osc}(f, \Delta) \leq \text{osc}(f, \{z_0\}) = 0.$$

We shall describe the behaviour of the coefficients of the system close to a set of degeneracy by means of a special exhaustion function h . Fix a set $\Gamma \subset \tilde{\partial}D$ and a positive locally Lipschitz function $h : D \rightarrow \mathbf{R}$ such that $\lim_{z \rightarrow \Gamma} h(z) = 0$ and

$$(1.5) \quad 0 < h_0 \leq \text{ess inf}_D |\nabla h(z)| \leq \text{ess sup}_D |\nabla h(z)| \leq h_1 < \infty,$$

where h_0, h_1 are some constants.

Below we let $E_t = \{z \in D : h(z) = t\}$ denote the level curve of h .

Theorem 1.6. *Let $\Gamma \subset \tilde{\partial}D$ be an arbitrary set, and let $w = (u, v)$ be an ACL_{loc}^2 homeomorphic solution of (1.1) from D onto $\mathcal{D}$ satisfying (1.2) such that $|u| < M$ in D . If*

$$(1.7) \quad \int_0^\infty dt \left(\int_{D \cap E_t} (a + d) \frac{ad - bc}{\delta} d\mathcal{H}^1(E_t) \right)^{-1} = \infty,$$

then u is monotone close to Γ .

For a homeomorphism $w : D \rightarrow \mathcal{D}$ and for an arbitrary $\Gamma \subset \tilde{\partial}D$ we set

$$w(\Gamma) = \{y \in \tilde{\partial}\mathcal{D} : \exists \text{ a sequence } z_n \in D, z_n \rightarrow \Gamma, \text{ such that } w(z_n) \rightarrow y\}.$$

Clearly, if $\Gamma \subset \partial D$ is connected and ∂D is a simple Jordan curve, then $w(\Gamma)$ is also connected.

A set $L \subset \tilde{\partial}\mathcal{D}$ is called *u -forked* if there are at least two points $w' = (u', v')$, $w'' = (u', v'') \in L$ such that the segment

$$l = \{(u, v) \in \mathbf{R}^2 : u = u', v' < v < v''\} \quad \text{lies in } \mathcal{D}$$

and separates from $\tilde{\partial}\mathcal{D} \setminus L$ some subdomain $U \subset \mathcal{D}$ with $\tilde{\partial}U \subset \bar{l} \cup L$.

For example, let $\mathcal{D}$ be a disk and let $\partial\mathcal{D}$ be its boundary circle. Then the right and left semicircles are *u -forked*; however, the upper and lower semicircles are not *u -forked*.

Theorem 1.8. *Let $M > 0$ be a constant. Suppose that $\mathcal{D}$ is a subdomain of $\{(u, v) : |u| < M\}$, the coefficients of (1.1) satisfy (1.7) and $L \subset \tilde{\partial}\mathcal{D}$ is u -forked. Then there is no ACL_{loc}^2 homeomorphic solution $w = w(z)$ of (1.1) from D onto $\mathcal{D}$ such that $w(\Gamma) \supset L$.*

2. PROOF OF THEOREM 1.6

Let Δ be a subdomain of D with $\partial''\Delta \subset \Gamma$. We shall prove that

$$(2.1) \quad \sup_{\Delta} u(z) = \sup_{\partial'\Delta} u(z).$$

Assume the contrary, that is, there exists a point $z_0 \in \Delta$ such that

$$u(z_0) > \sup_{\partial'\Delta} u(z) = A.$$

Choose $\epsilon > A$ such that $u(z_0) > \epsilon$. Fix the connected component U of the set $\{z \in \Delta : u(z) > \epsilon\}$ containing z_0 . By [15, Theorem 5.4.4] for almost all $\epsilon > A$, the sets $\{z \in \Delta : u(z) = \epsilon\}$ are locally rectifiable. Therefore, without loss of generality, we may assume that $\partial'U$ is locally rectifiable.

Fix numbers $0 < \delta' < \delta'' < h(z_0)$ and an absolutely continuous function $\psi_0 : [\delta', \delta''] \rightarrow [0, 1]$ such that $\psi_0(\delta'') = 1$ and $\psi_0(\delta') = 0$. We shall specify ψ_0 later. Define $\psi : (0, \infty) \rightarrow \mathbf{R}$ as

$$\psi(\tau) = \begin{cases} 1 & \text{for } \delta'' < \tau < \infty, \\ \psi_0(\tau) & \text{for } \delta' \leq \tau \leq \delta'', \\ 0 & \text{for } 0 < \tau < \delta'. \end{cases}$$

Then ψ is an absolutely continuous function. Write $\phi(z) = \psi(h(z))^2(u(z) - \epsilon)$ for $z \in U$ and $\phi \equiv 0$ for $z \in D \setminus U$. Using [9, Theorem 1.20] we conclude that $\phi \in ACL_{\text{loc}}^2(D)$. Because $\text{supp } \phi \subset \subset D$ we have by Green's formula

$$\int_U d(\phi dv) = 0$$

and hence

$$\int_U d\phi \wedge dv = 0.$$

Because

$$d\phi \wedge dv = \psi^2 du \wedge dv + 2\psi(u - \epsilon) d\psi \wedge dv,$$

we obtain

$$\int_U \psi^2(u_x v_y - u_y v_x) dx dy = -2 \int_U \psi(u - \epsilon)(\psi_x v_y - \psi_y v_x) dx dy.$$

This is clear for $u, v \in C^2(D)$. In the general case we can easily prove it if we use a standard approximation procedure (see, for example, [9, Lemma 14.15]).

From here

$$\int_U \psi^2(u_x v_y - u_y v_x) dx dy \leq 2 \int_U \psi |u - \epsilon| |\nabla \psi| |\nabla v| dx dy.$$

Using (1.1) we find

$$\begin{aligned}
& \int_U \psi^2 (au_x^2 + (b+c)u_xu_y + du_y^2) \, dxdy \\
& \leq 4M \int_U \psi |\nabla \psi| \sqrt{(au_x + bu_y)^2 + (cu_x + du_y)^2} \, dxdy \\
& = 4M \int_U \psi |\nabla \psi| \sqrt{(a^2 + c^2)u_x^2 + 2(ab + cd)u_xu_y + (b^2 + d^2)u_y^2} \, dxdy \\
& \leq 4M \left(\int_U |\nabla \psi|^2 \frac{(a^2 + c^2)u_x^2 + 2(ab + cd)u_xu_y + (b^2 + d^2)u_y^2}{au_x^2 + (b+c)u_xu_y + du_y^2} \, dxdy \right)^{1/2} \\
& \quad \times \left(\int_U \psi^2 (au_x^2 + (b+c)u_xu_y + du_y^2) \, dxdy \right)^{1/2}.
\end{aligned}$$

Thus we obtain

$$\begin{aligned}
(2.2) \quad & \int_U \psi^2 (au_x^2 + (b+c)u_xu_y + du_y^2) \, dxdy \\
& \leq 16M^2 \int_U |\nabla \psi|^2 \frac{(a^2 + c^2)u_x^2 + 2(ab + cd)u_xu_y + (b^2 + d^2)u_y^2}{au_x^2 + (b+c)u_xu_y + du_y^2} \, dxdy.
\end{aligned}$$

We write

$$I(\xi, \eta) = (a^2 + c^2)\xi^2 + 2(ab + cd)\xi\eta + (b^2 + d^2)\eta^2$$

and

$$II(\xi, \eta) = (a\xi^2 + (b+c)\xi\eta + d\eta^2).$$

From (1.2) it follows that the quadratic form $II(\xi, \eta)$ is positive definite, and hence the bundle of quadratic forms

$$H(\xi, \eta; \lambda) = I(\xi, \eta) - \lambda II(\xi, \eta)$$

is regular in the sense of [7, Chapter X].

Let $\lambda_{\max}$ be a maximal eigenvalue of $H(\xi, \eta; \lambda)$. Then using well-known properties of quadratic forms (see [7, Theorem 13, Chapter X]) for every $\xi, \eta \neq 0$, we have

$$(2.3) \quad \frac{I(\xi, \eta)}{II(\xi, \eta)} \leq \lambda_{\max}.$$

The characteristic equation of $H(\xi, \eta; \lambda)$ has the form

$$\begin{vmatrix} a^2 + c^2 - \lambda a & ab + cd - \lambda(b+c)/2 \\ ab + cd - \lambda(b+c)/2 & b^2 + d^2 - \lambda d \end{vmatrix} = 0,$$

i.e.,

$$\delta \lambda^2 - (a+d)(ad-bc)\lambda + (ad-bc)^2 = 0.$$

Solving this equation for λ and using (1.2) we find that

$$\lambda_{\max} \leq \mu \equiv \frac{(ad - bc)(a + d)}{\delta}.$$

Clearly, $ad - bc > 0$ since from (1.2) it follows that

$$0 < 4ad - (b + c)^2 \leq 4ad - 4bc.$$

By (2.3) for every $(x, y) \in U$, we obtain

$$(2.4) \quad \frac{(a^2 + c^2)u_x^2 + 2(ab + cd)u_x u_y + (b^2 + d^2)u_y^2}{au_x^2 + (b + c)u_x u_y + du_y^2} \leq \mu.$$

Thus from (2.2) we arrive at the estimate

$$(2.5) \quad \int_U \psi^2 (au_x^2 + (b + c)u_x u_y + du_y^2) \, dxdy \leq 16M^2 \int_U \mu |\nabla \phi|^2 \, dxdy.$$

Now we shall estimate the integral in the right side of (2.5). We have

$$I \equiv \int_U \mu |\nabla \phi|^2 \, dxdy = \int_{\substack{\delta' < h(z) < \delta'' \\ z \in U}} \mu \psi_0'^2(h(z)) |\nabla h|^2 \, dxdy$$

and next, by the well-known Kronrod-Federer coarea formula [3, Chapter 3, §2.4],

$$I = \int_{\delta'}^{\delta''} \psi_0'^2(t) \, dt \int_{E_t \cap U} \mu |\nabla h| \, d\mathcal{H}^1(E_t).$$

Using (1.5) we find

$$(2.6) \quad I \leq h_1 \int_{\delta'}^{\delta''} \psi_0'^2(t) \, dt \int_{E_t \cap U} \mu \, d\mathcal{H}^1(E_t).$$

Since $\psi_0(\delta') = 1$ and $\psi_0(\delta'') = 0$, we have

$$\begin{aligned} 1 &\leq \left(\int_{\delta'}^{\delta''} |\psi_0'(t)| \, dt \right)^2 \leq \int_{\delta'}^{\delta''} |\psi_0'|^2 \, dt \int_{E_t \cap U} \mu \, d\mathcal{H}^1(E_t) \\ &\quad \times \int_{\delta'}^{\delta''} dt \left(\int_{E_t \cap U} \mu \, d\mathcal{H}^1(E_t) \right)^{-1}. \end{aligned}$$

Therefore, for every absolutely continuous function $\psi_0 : [\delta', \delta''] \rightarrow [0, 1]$ such that $\psi_0(\delta'') = 1$ and $\psi_0(\delta') = 0$, we have

$$\left(\int_{\delta'}^{\delta''} dt \left(\int_{E_t \cap U} \mu \, d\mathcal{H}^1(E_t) \right)^{-1} \right)^{-1} \leq \int_{\delta'}^{\delta''} |\psi_0'|^2 \, dt \int_{E_t \cap U} \mu \, d\mathcal{H}^1(E_t).$$

Choosing for $\delta' \leq \tau \leq \delta''$,

$$\psi_0(\tau) = \int_{\delta'}^{\tau} dt \left(\int_{E_t \cap U} \mu d\mathcal{H}^1(E_t) \right)^{-1} \bigg/ \int_{\delta'}^{\delta''} dt \left(\int_{E_t \cap U} \mu d\mathcal{H}^1(E_t) \right)^{-1},$$

we have

$$\min_{\psi_0} \int_{\delta'}^{\delta''} |\psi'_0|^2 dt \int_{E_t \cap U} \mu d\mathcal{H}^1(E_t) = \left(\int_{\delta'}^{\delta''} dt \left(\int_{E_t \cap U} \mu d\mathcal{H}^1(E_t) \right)^{-1} \right)^{-1}.$$

Thus from (2.5) and (2.6) we see that

$$\begin{aligned} & \int_{\substack{\delta'' < h(z) \\ z \in U}} (au_x^2 + (b+c)u_xu_y + du_y^2) dx dy \\ & \leq 16M^2h_1 \left(\int_{\delta'}^{\delta''} dt \left(\int_{E_t \cap U} \mu d\mathcal{H}^1(E_t) \right)^{-1} \right)^{-1}. \end{aligned}$$

Setting here $\delta' \rightarrow 0$ we obtain

$$\begin{aligned} & \int_{\substack{\delta'' < h(z) \\ z \in U}} (au_x^2 + (b+c)u_xu_y + du_y^2) dx dy \\ & \leq 16M^2h_1 \left(\int_0^{\delta''} dt \left(\int_{E_t \cap U} \mu d\mathcal{H}^1(E_t) \right)^{-1} \right)^{-1}. \end{aligned}$$

Previously we chose $\delta'' < h(z_0)$ where $z_0 \in U$. Thus the open set $U_1 = \{z \in U : \delta'' < h(z_0)\}$ is not empty. Now from (1.7) we conclude that

$$\int_{U_1} (au_x^2 + (b+c)u_xu_y + du_y^2) dx dy = 0,$$

that is,

$$au_x^2 + (b+c)u_xu_y + du_y^2 = 0 \quad \text{a.e. in } U_1.$$

The assumption (1.2) implies that $\nabla u = 0$ a.e. in U_1 and hence $u \equiv \text{const}$ in U_1 . But $\delta'' > \delta' > 0$ can be arbitrarily small, and hence $\nabla u = 0$ a.e. in U . There is a contradiction with the definition of the connected component U and (2.1) is true.

Analogously, we obtain

$$\inf_{\Delta} u(z) = \inf_{\partial' \Delta} u(z)$$

and next, we arrive at (1.4). □

3. PROOF OF THEOREM 1.8

Suppose that there exists a homeomorphic solution $w(z)$ of (1.1), (1.2) such that $w(D) = \mathcal{D}$ and $L \subset w(\Gamma)$. Since L is u -forked, it follows that there is a vertical segment $l \subset \mathcal{D}$ with the end points in L and moreover, there exists a subdomain

U of $\mathcal{D}$ with the boundary $\tilde{\partial}U \subset \bar{l} \cup L$. Consider its preimage $\Delta = w^{-1}(U)$. By Theorem 1.6 the function u is monotone close to Γ . Hence,

$$\begin{aligned} \operatorname{osc}(u, \Delta) &\leq \operatorname{osc}(u, \tilde{\partial}'\Delta) \\ &= \operatorname{osc}(u, w^{-1}(l)) = 0. \end{aligned}$$

Thus, $u \equiv \text{const}$ in Δ , which is impossible. $\square$

4. AN EXAMPLE

Let γ be a simple open Jordan arc lying in the upper half-plane with end points $(0, 0)$ and $(1, 0)$ on the x -axis. We set

$$\Gamma = \{(x, y) : 0 \leq x \leq 1, y = 0\},$$

and denote by D the subdomain of $\mathbf{R}^2$ enclosed by $\gamma \cup \Gamma$.

Choose in Theorem 1.8 the function $h(z) = y$. Evidently, (1.5) is fulfilled.

Consider the case of (1.1) in which the functions $a = a(y)$, $d = d(y)$ are positive and $b = c = 0$. Then (1.1) admits the form

$$(4.1) \quad a(y)u_x = v_y, \quad d(y)u_y = -v_x.$$

The assumption (1.2) is fulfilled too.

The assumption (1.7) takes the form

$$(4.2) \quad \int_0^1 \frac{dy}{a(y) + d(y)} = \infty.$$

We obtain:

Corollary 4.3. *Let $\mathcal{D}$ be a simply connected subdomain of $\{(u, v) : |u| < M\}$, where $0 < M < \infty$ is a constant. Assume that (4.2) is fulfilled and $L \subset \tilde{\partial}\mathcal{D}$ is u -forked. Then there are no ACL_{loc}^2 solutions $w = w(z)$ of (4.1) mapping D homeomorphically onto $\mathcal{D}$ such that $L \subset w(\Gamma)$.*

ACKNOWLEDGEMENTS

The authors are indebted to Prof. V.A. Klyachin and to the anonymous referee for a number of useful remarks and corrections.

REFERENCES

- [1] B.V. BOJARSKI: Generalized solutions of an elliptic system of first order with discontinuous coefficients (in Russian), Mat. Sb. **43**, No. 4, 451–503, 1957. MR0106324 (21:5058)
- [2] M.A. BRAKALOVA AND J.A. JENKINS: On solutions of the Beltrami equation. J. Anal. Math. **76** (1998), 67–92. MR1676936 (2000h:30029)
- [3] YU. BURAGO AND V. A. ZALGALLER: Geometric inequalities, Leningrad, "Nauka", 1980 (in Russian). MR0602952 (82d:52009)
- [4] E.A. CHTCHERBAKOV (ŠČERBAKOV): Homeomorphic solutions of degenerate elliptic systems (in Russian), Math. Phys., Kiev, Naukova Dumka **8**, 187–190, 1970. MR0293239 (45:2316)
- [5] G. DAVID: Solutions de l'équation de Beltrami avec $\|\mu\|_\infty = 1$, Ann. Acad. Sci. Fenn. Ser. A I Math. **13**, 25–70, 1988. MR0975566 (90d:30058)
- [6] A. DŽURAEV: On a Beltrami system of equations degenerating on a line, - Soviet Math. Dokl. **10**, No. 2, 1969, 449–452. (Russian) Dokl. Akad. Nauk SSSR **185** 1969, 984–986. MR0241668 (39:3007)

- [7] F.R. GANTMACHER: Theory of Matrices, Chelsea, New York, 1959. MR0107649 (21:6372c)
- [8] V. GUTLYANSKII, O. MARTIO, T. SUGAWA AND M. VUORINEN: On the degenerate Beltrami equation, Reports of the Dept. of Math., Univ. of Helsinki, Preprint 282, 2001, 1-32, Trans. Amer. Math. Soc. (to appear).
- [9] J. HEINONEN, T. KILPELÄINEN AND O. MARTIO: Nonlinear Potential Theory of Degenerate Elliptic Equations, Oxford Mathematical Monographs, Oxford University Press, Oxford, 1993. MR1207810 (94e:31003)
- [10] O. MARTIO AND V.M. MIKLYUKOV: On existence and uniqueness of degenerate Beltrami equation, Reports of the Dept. of Math., Univ. of Helsinki, Preprint 347, 2003, 1-12.
- [11] O. MARTIO, V.M. MIKLYUKOV AND M. VUORINEN: Functions monotone close to boundary, Reports of the Dept. of Math., Univ. of Helsinki, Preprint 330, March 2002, 1-20, Tohoku Math. J. (to appear).
- [12] A.P. MIHAILOV: A problem for a mapping by solutions of elliptic system degenerating on boundary (in Russian), Sibirsk. Mat. Zh. **24**, No. 3, 119–127, 1983. MR0704163 (84m:30075)
- [13] I.S. OVCHINNIKOV: On existence of plane mappings for degenerate elliptic systems of the first order (in Russian), Dokl. AN SSSR **191**, No. 3, 526–529, 1970. MR0259355 (41:3993)
- [14] U. SREBRO AND E. YAKUBOV: Branched folder maps and alternating Beltrami equations, J. Anal. Math. **70**, 65-90, 1996. MR1444258 (98f:30021)
- [15] W.P. ZIEMER: Weakly Differentiable Functions, Graduate Texts in Mathematics 120, Springer-Verlag, New York, 1989. MR1014685 (91e:46046)

DEPARTMENT OF MATHEMATICS AND STATISTICS, P.O. Box 68, FIN-00014, UNIVERSITY OF HELSINKI, FINLAND

E-mail address: `martio@cc.helsinki.fi`

DEPARTMENT OF MATHEMATICS, VOLGOGRAD STATE UNIVERSITY, 2 PRODOLNAYA, 30, VOLGOGRAD, 400062, RUSSIA

E-mail address: `miklyuk@mail.ru`

DEPARTMENT OF MATHEMATICS, FIN-20014, UNIVERSITY OF TURKU, FINLAND

E-mail address: `vuorinen@csc.fi`