VECTOR BUNDLES ON A PRODUCT OF REAL ALGEBRAIC CURVES

J. BOCHNAK AND W. KUCHARZ

(Communicated by Michael Stillman)

ABSTRACT. We study complex vector bundles on a product of nonsingular real algebraic curves.

Throughout this note the term *real algebraic variety* designates a locally ringed space isomorphic to an algebraic subset of $\mathbb{R}^N$, for some N , endowed with the Zariski topology and the sheaf of $\mathbb{R}$ -valued regular functions. Recall that each Zariski locally closed subset of real projective space $\mathbb{P}^N(\mathbb{R})$ can be regarded as a real algebraic variety [3, Propositions 3.2.10, 3.4.3]. Morphisms of real algebraic varieties will be called *regular maps*. Every real algebraic variety carries also the *Euclidean topology*, which is determined by the usual metric topology on $\mathbb{R}$. Unless explicitly stated otherwise, all topological notions related to real algebraic varieties will refer to the Euclidean topology.

Given a real algebraic variety X , an interesting problem is the one of comparison between algebraic and topological vector bundles on X . We deal here with $\mathbb{C}$ -vector bundles, identifying $\mathbb{C}$ with $\mathbb{R}^2$ and viewing it as a real algebraic variety. A topological $\mathbb{C}$ -vector bundle ξ on X is said to *admit an algebraic structure* if it is topologically isomorphic to an algebraic subbundle of the trivial vector bundle with total space $X \times \mathbb{C}^p$, for some p (equivalently, if ξ is topologically isomorphic to the $\mathbb{C}$ -vector bundle corresponding, in the usual way [11, 12], to a finitely generated projective module over the ring of regular functions from X into $\mathbb{C}$) [3, 4]. It is known that every topological $\mathbb{C}$ -vector bundle on the unit sphere

$$S^n = \{(x_0, \dots, x_n) \in \mathbb{R}^{n+1} \mid x_0^2 + \dots + x_n^2 = 1\}$$

or projective space $\mathbb{P}^n(\mathbb{R})$ admits an algebraic structure [9], [3, p. 326], and [4, Example 2.3(a)]. However, the behavior of $\mathbb{C}$ -vector bundles on other “simple” real algebraic varieties can be drastically different. For example, a topological $\mathbb{C}$ -vector bundle on the k -fold product $S^1 \times \dots \times S^1$ admits an algebraic structure if and only if it is stably trivial [3, Corollary 12.6.6, and p. 326]. More generally, for “generic” compact nonsingular real algebraic curves $C_1, \dots, C_k$, a topological $\mathbb{C}$ -vector bundle on $C_1 \times \dots \times C_k$ admits an algebraic structure if and only if it is stably trivial (cf. [5, 6, 7] for the precise meaning of the adjective “generic”).

Received by the editors December 6, 2001 and, in revised form, February 26, 2004.

2000 *Mathematics Subject Classification*. Primary 14P25, 19E99.

Both authors were partially supported by the Volkswagen Stiftung (Research in Pairs at Oberwolfach).

We have the following:

Theorem. *Let M_i be a compact smooth submanifold (of class $\mathcal{C}^\infty$ and without boundary) of $\mathbb{P}^{n_i}(\mathbb{R})$, with $2 \dim M_i + 1 \leq n_i$ for $1 \leq i \leq k$. Then for each i there exists a smooth embedding $e_i : M_i \rightarrow \mathbb{P}^{n_i}(\mathbb{R})$, arbitrarily close in the $\mathcal{C}^\infty$ topology to the inclusion map $M_i \hookrightarrow \mathbb{P}^{n_i}(\mathbb{R})$, such that $X_i = e_i(M_i)$ is a nonsingular Zariski closed subvariety of $\mathbb{P}^{n_i}(\mathbb{R})$ and every topological $\mathbb{C}$ -line bundle on $X_1 \times \cdots \times X_k$ admits an algebraic structure; assuming $\dim M_i = 1$ for $1 \leq i \leq k$, every topological $\mathbb{C}$ -vector bundle on $X_1 \times \cdots \times X_k$ admits an algebraic structure. Furthermore, if $n_1 = \cdots = n_k$ and $M_1 = \cdots = M_k$, then one can take $e_1 = \cdots = e_k$ and hence $X_1 = \cdots = X_k$.*

Let us recall a result of R. Benedetti and A. Tognoli [2, Theorem 4.2]. For any compact smooth submanifold (without boundary) M of $\mathbb{P}^n(\mathbb{R})$, with $2 \dim M + 1 \leq n$, there exists a smooth embedding $e : M \hookrightarrow \mathbb{P}^n(\mathbb{R})$, arbitrarily close in the $\mathcal{C}^\infty$ topology to the inclusion map $M \hookrightarrow \mathbb{P}^n(\mathbb{R})$, such that $X = e(M)$ is a nonsingular Zariski closed subvariety of $\mathbb{P}^n(\mathbb{R})$ and every topological $\mathbb{C}$ -vector bundle on X admits an algebraic structure (strictly speaking, [2] deals with $\mathbb{R}$ -vector bundles and with $\mathbb{R}^n$ instead of $\mathbb{P}^n(\mathbb{R})$, but a straightforward modification of the proof yields the assertion just stated). This does not directly imply our theorem.

In preparation for the proof we shall review a certain construction. Let X be a nonsingular real algebraic variety. A nonsingular complexification of X is a pair (V, j) , where V is a nonsingular complex quasiprojective variety defined over $\mathbb{R}$ and $j : X \rightarrow V$ is an injective map such that the set $V(\mathbb{R})$ of real points of V is Zariski dense in V , $j(X) = V(\mathbb{R})$, and j , viewed as a map from X onto $V(\mathbb{R})$, is a regular isomorphism of real algebraic varieties. Define $H_{\text{alg}}^2(V, \mathbb{Z})$ to be the subgroup of $H^2(V, \mathbb{Z})$ consisting of all cohomology classes of the form $c_1(L)$, where L is a complex algebraic line bundle on V and $c_1(-)$ stands for the first Chern class. Set

$$H_{\mathbb{C}\text{-alg}}^2(X, \mathbb{Z}) = j^*(H_{\text{alg}}^2(V, \mathbb{Z})),$$

where $j^* : H^*(V, \mathbb{Z}) \rightarrow H^*(X, \mathbb{Z})$ is the homomorphism induced by j . One readily checks that $H_{\mathbb{C}\text{-alg}}^2(X, \mathbb{Z})$ does not depend on the choice of (V, j) [4]. Given a topological $\mathbb{C}$ -line bundle λ on X ,

$$(1) \quad \lambda \text{ admits an algebraic structure} \Leftrightarrow c_1(\lambda) \in H_{\mathbb{C}\text{-alg}}^2(X, \mathbb{Z})$$

[4]. For any regular map $f : X \rightarrow Y$ between nonsingular real algebraic varieties, one has (cf. [4])

$$(2) \quad f^*(H_{\mathbb{C}\text{-alg}}^2(Y, \mathbb{Z})) \subseteq H_{\mathbb{C}\text{-alg}}^2(X, \mathbb{Z}).$$

If $Y_1, \dots, Y_n$ are connected nonsingular real algebraic varieties, then

$$H^2(Y_1 \times \cdots \times Y_n, \mathbb{Z}) = \sum_{i < j} p_{ij}^*(H^2(Y_i \times Y_j, \mathbb{Z})),$$

where $p_{ij} : Y_1 \times \cdots \times Y_n \rightarrow Y_i \times Y_j$ is the canonical projection, and hence (2) implies

$$(3) \quad H_{\mathbb{C}\text{-alg}}^2(Y_1 \times \cdots \times Y_n, \mathbb{Z}) = H^2(Y_1 \times \cdots \times Y_n, \mathbb{Z}),$$

provided $H_{\mathbb{C}\text{-alg}}^2(Y_i \times Y_j, \mathbb{Z}) = H^2(Y_i \times Y_j, \mathbb{Z})$ for all $i < j$.

Note that

$$\sigma : \mathbb{P}^n(\mathbb{C}) \times \mathbb{P}^n(\mathbb{C}) \rightarrow \mathbb{P}^n(\mathbb{C}) \times \mathbb{P}^n(\mathbb{C}), \quad \sigma(x, y) = (\bar{y}, \bar{x}),$$

where the bar indicates complex conjugation, is an antiholomorphic involution. Thus $\mathbb{P}^n(\mathbb{C}) \times \mathbb{P}^n(\mathbb{C})$ can be regarded as a complex projective variety defined over $\mathbb{R}$, whose set of real points, denoted here by $\mathbb{P}^n(\mathbb{C})_{\mathbb{R}}$, coincides with the set of fixed points of the involution σ . Clearly, the map $\mathbb{P}^n(\mathbb{C}) \rightarrow \mathbb{P}^n(\mathbb{C})_{\mathbb{R}}$, $x \rightarrow \sigma(x, x) = (x, \bar{x})$, is a homeomorphism. Furthermore, by construction,

$$(4) \quad H_{\mathbb{C}\text{-alg}}^2(\mathbb{P}^n(\mathbb{C})_{\mathbb{R}}, \mathbb{Z}) = H^2(\mathbb{P}^n(\mathbb{C})_{\mathbb{R}}, \mathbb{Z}).$$

Proof of the theorem. By [6, Corollary 1.6, Proposition 2.2], there exists a nonsingular algebraic curve C in $\mathbb{P}^2(\mathbb{R})$ such that C is connected and

$$(5) \quad H_{\mathbb{C}\text{-alg}}^2(C \times C, \mathbb{Z}) = H^2(C \times C, \mathbb{Z}).$$

Choose a positive integer r with the property that the group $H^q(M_i, \mathbb{Z})$ can be generated by r elements for $1 \leq q \leq 2$ and $1 \leq i \leq k$. Let ℓ be a positive integer satisfying $\dim M_i + 1 \leq \ell$ for $1 \leq i \leq k$. Setting

$$B = (\mathbb{P}^{\ell}(\mathbb{C})_{\mathbb{R}})^r \times C^r,$$

we can find a smooth map $\varphi_i : M_i \rightarrow B$ for which the induced homomorphism

$$\varphi_i^* : H^q(B, \mathbb{Z}) \rightarrow H^q(M_i, \mathbb{Z})$$

is surjective for $1 \leq q \leq 2$ and $1 \leq i \leq k$ (see, for example, [8, Chap. V, 11.6, 11.9; Chap. VII, Sect. 12]).

Given a nonnegative integer p , we identify $\mathbb{P}^{n_i}(\mathbb{R})$ with the subset $\mathbb{P}^{n_i}(\mathbb{R}) \times \{0\}$ of $\mathbb{P}^{n_i}(\mathbb{R}) \times \mathbb{R}^p$; thus $\mathbb{P}^{n_i}(\mathbb{R}) \subseteq \mathbb{P}^{n_i}(\mathbb{R}) \times \mathbb{R}^p$. By [1, Theorem 2.8.4, Lemma 2.7.1], there exists a nonnegative integer p , a smooth embedding $\epsilon_i : M_i \rightarrow \mathbb{P}^{n_i}(\mathbb{R}) \times \mathbb{R}^p$, a nonsingular Zariski closed subvariety Y_i of $\mathbb{P}^{n_i} \times \mathbb{R}^p$, and a regular map $\psi_i : Y_i \rightarrow B$ such that $Y_i = \epsilon_i(M_i)$, ϵ_i is close in the $\mathcal{C}^{\infty}$ topology to the inclusion map $M_i \hookrightarrow \mathbb{P}^{n_i}(\mathbb{R}) \times \mathbb{R}^p$, and φ_i is homotopic to $\psi_i \circ \epsilon_i$ (the cited results of [1] are applicable since every homology class in $H_j(B, \mathbb{Z}/2)$, for $0 \leq j \leq \dim B$, can be represented by a Zariski closed subvariety of B). In particular, the induced homomorphism

$$\psi_i^* : H^q(B, \mathbb{Z}) \rightarrow H^q(Y_i, \mathbb{Z})$$

is surjective for $1 \leq q \leq 2$. Since $2 \dim Y_i + 1 \leq n_i$, one can find an algebraic embedding $\eta_i : Y_i \rightarrow \mathbb{P}^{n_i}(\mathbb{R})$ (this means that $\eta_i(Y_i)$ is a Zariski closed subvariety of $\mathbb{P}^{n_i}(\mathbb{R})$ and $\eta_i : Y_i \rightarrow \eta_i(Y_i)$ is a regular isomorphism), close to $\pi_i|_{Y_i}$, where $\pi_i : \mathbb{P}^{n_i}(\mathbb{R}) \times \mathbb{R}^p \rightarrow \mathbb{P}^{n_i}(\mathbb{R})$ is the canonical projection; cf. [10, pp. 21, 22]. By construction, $e_i = \eta_i \circ \epsilon_i : M_i \rightarrow \mathbb{P}^{n_i}(\mathbb{R})$ is a smooth embedding, close in the $\mathcal{C}^{\infty}$ topology to the inclusion map $M_i \hookrightarrow \mathbb{P}^{n_i}(\mathbb{R})$, $X_i = e_i(M_i) = \eta_i(Y_i)$ is a nonsingular Zariski closed subvariety of $\mathbb{P}^{n_i}(\mathbb{R})$, and $f_i = \psi_i \circ \eta_i^{-1} : X_i \rightarrow B$ is a regular map for which the induced homomorphism

$$f_i^* : H^q(B, \mathbb{Z}) \rightarrow H^q(X_i, \mathbb{Z})$$

is surjective for $1 \leq q \leq 2$.

Consider the regular map

$$f = f_1 \times \cdots \times f_k : X_1 \times \cdots \times X_k \rightarrow B^k.$$

Clearly, the induced homomorphism

$$(6) \quad f^* : H^2(B^k, \mathbb{Z}) \rightarrow H^2(X_1 \times \cdots \times X_k, \mathbb{Z})$$

is surjective. Furthermore, it easily follows from (3), (4), (5) that

$$(7) \quad H_{\mathbb{C}\text{-alg}}^2(B^k, \mathbb{Z}) = H^2(B^k, \mathbb{Z}).$$

Combining (2), (6), (7), we obtain

$$(8) \quad H_{\mathbb{C}\text{-alg}}^2(X_1 \times \cdots \times X_k, \mathbb{Z}) = H^2(X_1 \times \cdots \times X_k, \mathbb{Z}),$$

which in view of (1) implies that every topological $\mathbb{C}$ -line bundle on $X_1 \times \cdots \times X_k$ admits an algebraic structure. Thus the first part of the theorem is proved.

Clearly, assuming $n_1 = \cdots = n_k$ and $M_1 = \cdots = M_k$, we can take $e_1 = \cdots = e_k$ and $X_1 = \cdots = X_k$.

Suppose now $\dim M_i = 1$ for $1 \leq i \leq k$. Without loss of generality, we may assume that each M_i is connected. Then the real algebraic variety $X = X_1 \times \cdots \times X_k$ is connected, $H^*(X, \mathbb{Z})$ has no torsion, and the ring $H^{\text{even}}(X, \mathbb{Z}) = \bigoplus_{j \geq 0} H^{2j}(X, \mathbb{Z})$ is generated by $H^0(X, \mathbb{Z})$ and $H^2(X, \mathbb{Z})$. Since, in addition to these facts, condition (8) holds, it follows from [5, Proposition 2.4] that every topological $\mathbb{C}$ -vector bundle on X admits an algebraic structure. This completes the proof.

REFERENCES

- [1] S. Akbulut and H. King, *Topology of Real Algebraic Sets*, Math. Sci. Research Institute Publ. **25**, New York Berlin Heidelberg, Springer, 1992. MR1225577 (94m:57001)
- [2] R. Benedetti and A. Tognoli, *On real algebraic vector bundles*, Bull. Sci. Math. **104** (1980), 89-112. MR0560747 (81e:14009)
- [3] J. Bochnak, M. Coste and M.-F. Roy, *Real Algebraic Geometry*, Ergebnisse der Mathematik und ihrer Grenzgebiete, Folge 3, Vol. **36**, Berlin Heidelberg New York, Springer, 1998. MR1659509 (2000a:14067)
- [4] J. Bochnak, M. Buchner and W. Kucharz, *Vector bundles on real algebraic varieties*, K-Theory **3** (1989), 271-298, Erratum, K-Theory **4** (1990), p. 103. MR1040403 (91b:14075)
- [5] J. Bochnak and W. Kucharz, *Vector bundles on a product of real cubic curves*, K-Theory **6** (1992), 487-497. MR1204824 (94e:14077)
- [6] J. Bochnak and W. Kucharz, *Elliptic curves and real algebraic morphisms*, J. Algebraic Geometry **2** (1993), 635-666. MR1227471 (94e:14072)
- [7] J. Bochnak, W. Kucharz, and R. Silhol, *Morphisms, line bundles and moduli spaces in real algebraic geometry*, Inst. Hautes Etudes Sci. Publ. Math. **86** (1997), 5-65. MR1608561 (99h:14055)
- [8] G. E. Bredon, *Geometry and Topology*, New York Berlin Heidelberg, Springer, 1997. MR1700700 (2000b:55001)
- [9] R. Fossum, *Vector bundles on spheres are algebraic*, Invent. Math. **8** (1969), 222-225. MR0250298 (40:3537)
- [10] N. Ivanov, *Approximation of smooth manifolds by real algebraic sets*, Russian Math. Surveys **37** (1982), 1-59. MR0643764 (84i:57029)
- [11] J.-P. Serre, *Faisceaux algébriques cohérents*, Ann. of Math. **61** (1955), 197-278. MR0068874 (16:953c)
- [12] R. G. Swan, *Vector bundles and projective modules*, Trans. Amer. Math. Soc. **105** (1962), 264-277. MR0143225 (26:785)

DEPARTMENT OF MATHEMATICS, VRIJE UNIVERSITEIT, DE BOELELAAN 1081A, 1081 HV AMSTERDAM, THE NETHERLANDS

E-mail address: bochnak@cs.vu.nl

DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF NEW MEXICO, ALBUQUERQUE, NEW MEXICO 87131-1141

E-mail address: kucharz@math.unm.edu