ON DECOMPOSITIONS OF BANACH SPACES OF CONTINUOUS FUNCTIONS ON MRÓWKA'S SPACES

PIOTR KOSZMIDER

(Communicated by Alan Dow)

ABSTRACT. It is well known that if K is infinite compact Hausdorff and scattered (i.e., with no perfect subsets), then the Banach space $C(K)$ of continuous functions on K has complemented copies of c_0 , i.e., $C(K) \sim c_0 \oplus X \sim c_0 \oplus c_0 \oplus X \sim c_0 \oplus C(K)$. We address the question if this could be the only type of decompositions of $C(K) \not\sim c_0$ into infinite-dimensional summands for K infinite, scattered. Making a special set-theoretic assumption such as the continuum hypothesis or Martin's axiom we construct an example of Mrówka's space (i.e., obtained from an almost disjoint family of sets of positive integers) which answers positively the above question.

1. INTRODUCTION

This paper is concerned with a set-theoretic construction of a compact topological space which has interesting consequences in the isomorphic theory of Banach spaces, of continuous functions with the supremum norm.¹ It was recently proved in [7] that there are Banach spaces $C(K)$ of continuous functions on compact connected K with no non-trivial decomposition $C(K) = A \oplus B$. Actually it was only recently proved that there are at all such Banach spaces (not of continuous functions) [4]. In this paper we consider similar questions about the minimal possible space of operators or collection of projections or complemented subspaces in the context of K compact and scattered.

Infinite scattered compact spaces have non-trivial convergent sequences of isolated points, which easily give rise to complemented copies of c_0 in the function space which implies that there are many operators including projections with ranges

Received by the editors July 24, 2003 and, in revised form, April 15, 2004.

2000 *Mathematics Subject Classification.* Primary 03E50, 46E15, 54G12.

Key words and phrases. Banach spaces of continuous functions, few operators, scattered spaces, almost disjoint families.

The author acknowledges support from CNPQ, Processo Número 300369/01-8, from FAPESP, Processo Número 02/03677-7 and from Centre de Recerca Matemàtica at Universidad Autònoma de Barcelona.

¹ $\sim$ will denote an isomorphism of Banach spaces, i.e., a linear isomorphism which is continuous. A is complemented in X if and only if $X \sim A \oplus B$ for some B . This is equivalent to the existence of a projection $P : X \rightarrow A$, i.e., an onto bounded operator such that $P^2 = P$ ([17], §21). Since finite-dimensional or finite-co-dimensional subspaces are always complemented, by nontrivial decompositions we mean those with two infinite-dimensional factors (see [17], [11]). Unless stated otherwise all compact spaces are infinite and Hausdorff and all Banach spaces are infinite-dimensional.

included in c_0 . We ask in this paper if it is possible that there are no more operators and no more complemented subspaces in a $C(K) \not\sim c_0$ for a scattered K . In section 2, we obtain consistent examples.

Theorem. *Assuming CH or MA there is a compact separable scattered Hausdorff space with $C(K) \not\sim c_0$ such that all operators $T : C(K) \rightarrow C(K)$ are of the form $cI + S$ where c is a real and S has range included in a subspace isomorphic to c_0 . Moreover the only decompositions $C(K) = A \oplus B$ into two infinite-dimensional complemented subspaces are such that $A \sim c_0$ and $B \sim C(K)$ or $A \sim C(K)$ and $B \sim c_0$.*

Note that such a space K must be non-metrizable since for separable $C(K)$'s we have that the space is isomorphic to its square and that operators into c_0 have relatively simple structure, i.e., they correspond to weakly* null sequences in the dual to $C(K)$. Note that the question of the existence of Banach spaces whose only operators are scalar multiples of the identity plus an operator with separable range has been considered by various authors [18], [19], [20], [6]. We do not know if any set-theoretic assumption is needed for our $C(K)$ or even to get an example of any Banach space having the above properties, i.e., $c_0 \not\sim X \sim c_0 \oplus X$ and whether this type of a non-trivial decomposition is the only one. We know, however, that the existence of $C(K) \not\sim c_0$ which is Lindelöf in the weak topology and has the above properties is consistent and independent from the axioms of ZFC. This result will be published elsewhere.

In this paper we only analyze the Banach space of continuous functions on Mrówka's or Isbell's space, also known as the Ψ -space, which was first considered by Alexandroff. If $\mathcal{A} = \{A_\xi : \xi < \kappa\}$ is an infinite almost disjoint (i.e., with finite pairwise intersections) family of subsets of N , then the space $K_{\mathcal{A}}$ is the Stone space of the Boolean algebra generated by $\mathcal{A}$. The Banach space of continuous functions on $K_{\mathcal{A}}$ was considered for example in [5] (example 2) or in [12] (example 8). There are three types of ultrafilters-points of $K_{\mathcal{A}}$: first the isolated points corresponding to principal ultrafilters generated by singletons of $n \in N$; the second level is made up from ultrafilters generated by A_ξ and the cofinite sets (these points will be denoted x_ξ); the third level is represented by the unique ultrafilter which contains all cofinite sets and the complements of A_ξ 's (this point will be denoted by ∞). We will use the same notation A_ξ for the subset of N and the corresponding clopen subset of $K_{\mathcal{A}}$ hoping that the meaning is clear from the context. The sets $A_\xi - \{0, \dots, n-1\}$ will be denoted by A_ξ^n , the characteristic functions of $n \in N$ will be denoted by $e_n \in C(K_{\mathcal{A}})$ and the characteristic functions of the $A_\xi^n \cup \{x_\xi\}$'s will be denoted by $d_\xi^n \in C(K_{\mathcal{A}})$. $C_0(K_{\mathcal{A}})$ denotes the space of functions which vanish at ∞ . The following are two natural questions, which appear in the light of the results of this paper:

Question 1. Is there a separable Banach space $X \not\sim c_0$ whose only non-trivial decompositions are of the form $c_0 \oplus X$?

Question 2. Are there (with no additional set-theoretic assumption) almost disjoint families such that the only decompositions of $C(K_{\mathcal{A}})$ into two factors have one factor separable?

Let us make more comments on the second question. One can prove (Lemma 4) that the property of having few decompositions in the sense of our theorem for a

$C(K)$ where K is Mrówka's space is equivalent to the fact that the only decompositions of $C(K)$ have just one separable factor. Easy examples of decompositions for $A \subseteq N$ are of the form

$$C(K) \sim C_0(\overline{A}) \oplus C_0(\overline{N-A}) \oplus R,$$

where for every $A_\xi \in \mathcal{A}$ one has either $A_\xi - A$ is finite or $A \cap A_\xi$ is finite. Such an A is called a separation of $\mathcal{A}$. The projection on the first factor is given by $P_1(f) = (f - f(\infty))|_{\overline{A}}$, and similarly for the second factor, while the last projection is given by $P_3(f) = f(\infty)$.

Thus the family $\mathcal{A}$ as in the theorem cannot have separations into two uncountable parts. There are two known examples of almost disjoint families with this property which, unlike our theorem, do not require any additional set-theoretic assumptions. One, due to Lusin ([10], [3], 4.1), is of cardinality ω_1 , and the other due to Mrówka [13, 3.11] is maximal and of cardinality 2^ω .

Section 3 is devoted to showing that Lusin's family cannot witness the theorem in ZFC. It turns out that under Martin's axiom all spaces $C(K_{\mathcal{A}})$ for $\mathcal{A}$ of cardinality less than 2^ω have many decompositions into both non-separable factors (even though, as in the case of Lusin's family, there may be no separations of the family into two uncountable parts). On the other hand, Mrówka's families could serve to prove the existence of spaces such as in the theorem without the use of any special set-theoretic assumptions. As Mrówka's argument relies on the perfect set property of Borel sets of the reals (we thank A. Miller for explaining to us this construction in [11]), it is reasonable to expect that the positive answer could be obtained by the methods of descriptive set-theory. In the light of our result from section 3 it is natural to ask the following:

Question 3. Suppose MA. Is it true that if $|\mathcal{A}| = |\mathcal{A}'| < 2^\omega$, then $C(K_{\mathcal{A}})$ is isomorphic to $C(K_{\mathcal{A}'})$?

In the context of the Pełczyński decomposition method ([14], [11], Appendix) one could ask:

Question 4. Suppose MA. Is it true that if $|\mathcal{A}| < 2^\omega$, then $C(K_{\mathcal{A}})$ is isomorphic to its square?

It seems also that it is unknown² in ZFC (of course our result decides it under a special set-theoretic assumption):

Question 5. Are there two almost disjoint families $\mathcal{A}$ and $\mathcal{A}'$ of the same cardinality such that $C(K_{\mathcal{A}})$ is not isomorphic to $C(K_{\mathcal{A}'})$?

Our terminology follows [17], [3], [2]. In particular we consider several inter-related structures such as Boolean algebras, compact spaces, Banach spaces of continuous functions, Radon measures, etc. The main links between them are given by the Stone duality and the Riesz representation theorem. We also use Rosenthal's lemma ([16], [2]), which can be formulated in our context as follows: *given a sequence of absolutely convergent series $\sum_{n=0}^{\infty} m_n^k$ whose sums are bounded by the same number and given an $\varepsilon > 0$, one can find an infinite set $M \subseteq N$ such that $\sum_{n \in M - \{k\}} m_n^k < \varepsilon$ for every $k \in M$.*

²Mrówka shows in [13], 3.5, with no additional set-theoretic assumption, that there are 2^{2^ω} non-homeomorphic spaces $K_{\mathcal{A}}$, and it follows from [13], 3.6 that there is one $K_{\mathcal{A}}$ such that $C(K_{\mathcal{A}})$ has 2^ω decompositions into nonseparable factors.

Section 3, which deals with Martin's axiom, uses the terminology of chapter 2 of [8]. The terminology regarding the combinatorics of sets of integers follows [3]; in particular, $A \subseteq^* B$ means that $A - B$ is finite, and almost disjoint means with finite intersection. All almost disjoint families are assumed to be infinite and consist of infinite sets. A pseudointersection of a family of sets $\mathcal{F}$ is an infinite set A such that $A \subseteq^* B$ for all $B \in \mathcal{F}$. The number p is the minimal cardinal such that there is a centred (intersections of finite subfamilies are infinite) family of sets of integers of cardinality p without a pseudointersection. Martin's axiom (and so the continuum hypothesis) implies $p = 2^\omega$. An ideal of subsets of N is a family of proper subsets of N which contains all finite sets and is closed under subsets and finite unions.

2. SPACES OF CONTINUOUS FUNCTIONS ON MRÓWKA'S SPACES WITH FEW OPERATORS

Lemma 1. *Assume that $p = 2^\omega$. Suppose that $\mathcal{I}$ is an ideal of subsets of N which is generated by $< 2^\omega$ sets. Suppose that $\mathcal{F}$ is a family of finite subsets of N such that for every $I \in \mathcal{I}$ there is an $F \in \mathcal{F}$ such that $I \cap F = \emptyset$. Then there is a sequence $(F_n)_{n \in N}$ of pairwise disjoint elements of $\mathcal{F}$ such that $I \cap \bigcup \{F_n : n \in N\}$ is finite for all $I \in \mathcal{I}$.*

Proof. The assumption $p = 2^\omega$ says that any centred family of subsets of N of cardinality $< 2^\omega$ has a pseudointersection (see [3]), i.e., a set which is almost included in all members of the family. Identify N with the family $[N]^{<\omega}$ of all finite subsets of N and consider a centred family

$$\{[N - I]^{<\omega} \cap \mathcal{F} : I \in \mathcal{I}\}.$$

Let $\mathcal{G}$ be its pseudointersection. It exists since $\mathcal{I}$ is generated by less than p sets. Any infinite subset of $\mathcal{G}$ is another pseudointersection. So, using the fact that ideals contain all finite sets of N , one can choose a pairwise disjoint sequence which is a pseudointersection as well. It satisfies the lemma. $\square$

Proposition 2. *Assume that $p = 2^\omega$. There is an almost disjoint family $\mathcal{A}$ such that every bounded linear operator on $C_0(K_{\mathcal{A}})$ is of the form $cI + S$ where c is a constant and S has separable range.*

Proof. Before the construction, fix an enumeration $(m_n^k(\beta) : n, k \in N, \beta < 2^\omega)$, with cofinally many repetitions, of all real sequences $(m_n^k : n, k \in N)$ such that $\sum_{n=0}^\infty |m_n^k| < \rho$ for any $k \in N$ and some $\rho > 0$. Construct the almost disjoint family from three parts:

$$(A_\beta : \beta < 2^\omega) = (B_\beta : \beta < 2^\omega) \cup (C_\beta : \beta < 2^\omega) \cup (D_\beta : \beta < 2^\omega)$$

by induction in $\beta < 2^\omega$. Let $\mathcal{I}_\beta$ denote the ideal generated by $(B_\alpha, C_\alpha, D_\alpha : \alpha < \beta)$ and the finite subsets of N . Suppose that $B_\alpha, C_\alpha, D_\alpha$ for $\alpha < \beta$ are already constructed. At the stage β we inquire if the following is satisfied for some fixed $\varepsilon > 0$:

$$1) \quad \forall A \in \mathcal{I}_\beta \exists F \in [N - A]^{<\omega} \exists k \in N - (A \cup F) \mid \sum_{n \in F} m_n^k(\beta) > \varepsilon.$$

If 1) is not satisfied, then B_β and C_β are arbitrary as long as they are almost disjoint and almost disjoint from the sets already constructed; the assumption $p = 2^\omega$ implies that we can always find such sets since $p = 2^\omega > \beta$ ($p \leq a$ in [3]).

If 1) is satisfied, then $B_\beta = \bigcup \{F_i : i \in 2N\}$ and $C_\beta = \{k_i : i \in N\}$ where the F_i 's are pairwise disjoint finite sets and the k_i 's are distinct positive integers, such that $k_i \notin F_j$ for $i, j \in N$ and

$$2) \quad \forall i \in N \quad \left| \sum_{n \in F_i} m_n^{k_i}(\beta) \right| > \varepsilon,$$

$$3) \quad \forall i \in N \quad \sum_{j \neq i} \sum_{n \in F_j} |m_n^{k_i}(\beta)| < \varepsilon/3.$$

It follows from Lemma 1 applied to the $F \cup \{k\}$'s such as in 1) and $\mathcal{I}_\beta$ that there exist F_n 's and k_n 's satisfying 2) such that the corresponding B_β, C_β are almost disjoint from the previously constructed sets B_α, C_α and D_α for $\beta < 2^\omega$. Rosenthal's lemma implies that one can thin out the sequence obtaining 3).

Define $\mathcal{J}_\beta$ to be the ideal generated by the sets of $\mathcal{I}_\beta$ and B_β, C_β . Before the construction of D_β we inquire if the following is satisfied for some fixed reals $r_1 < r_2$:

$$4) \quad \forall A \in \mathcal{J}_\beta \quad \exists k, l \in N - A \quad m_k^k < r_1 < r_2 < m_l^l.$$

If 4) is not satisfied, then D_β is arbitrary as long as it is almost disjoint from the sets already constructed. Again, it exists, since $p = 2^\omega > \beta$.

If 4) is satisfied, then pick some $r_1 < r_2$ witnessing it and define $D_\beta = \{k_n : n \in N\} \cup \{l_n : n \in N\}$ where $k_n \neq l_j$ for all $n, j \in N$ and

$$5) \quad \forall n \in N \quad m_{k_n}^{k_n} < r_1 < r_2 < m_{l_n}^{l_n},$$

$$6) \quad \forall n \in N \quad \sum_{j \neq n} |m_{l_j}^{l_n}| < \frac{r_2 - r_1}{6}, \quad \sum_{j \neq n} |m_{k_j}^{k_n}| < \frac{r_2 - r_1}{6},$$

$$7) \quad \forall n \in N \quad \sum_{j \in N} |m_{k_j}^{l_n}| < \frac{r_2 - r_1}{6}, \quad \sum_{j \in N} |m_{l_j}^{k_n}| < \frac{r_2 - r_1}{6}.$$

Let us justify the existence of such a D_β . It follows from Lemma 1 applied to $\{k, l\}$ satisfying 4) that there exist two infinite disjoint sets $K, L \subseteq N$ such that $m_k^k < r_1 < r_2 < m_l^l$ for every $k \in K$ and $l \in L$ and both K and L are almost disjoint from the previously constructed sets B_α, C_α for $\alpha < \beta$ and from B_β, C_β . Thus, any enumeration of $K = \{k'_n : n \in N\}$ and $L = \{l'_n : n \in N\}$ would give 5). To obtain 6) and 7) one needs to apply Rosenthal's lemma to $\frac{r_2 - r_1}{6}$ and the series

$$\sum_{i=0}^{\infty} (|m_{k'_i}^{k'_n}| + |m_{l'_i}^{k'_n}| + |m_{k'_i}^{l'_n}| + |m_{l'_i}^{l'_n}|)$$

for all $n \in N$ obtaining infinite $M \subseteq N$ as in Rosenthal's lemma (see the introduction). Now consider two disjoint infinite sets $M_1, M_2 \subseteq M$ and enumerate $\{k_n : n \in N\} = M_1$ and $\{l_n : n \in N\} = M_2$. 6) and 7) are satisfied by the construction. This completes the construction. $\square$

Let $T : C_0(K_A) \rightarrow C_0(K_A)$. Define $g(n) = T(e_n)(n)$. Define $m_n^k = T(e_n)(k)$, i.e., $g(n) = m_n^n$.

Claim 1. *There is a countable set $X \subseteq 2^\omega$ such that for every $\beta \in 2^\omega - X$, for every $n, k \in N$ we have*

$$T(d_\beta^n)(k) = \sum_{i \in A_\beta^n} T(e_i)(k).$$

Proof of the Claim. Note that $\phi_k(f) = T(f)(k) = T^*(\delta_k)$ for $k \in N$ is a bounded linear functional on $C_0(K_{\mathcal{A}})$. Thus by the Riesz representation theorem (see [17], 18.5.3) there is a Radon measure μ_k on $K_{\mathcal{A}}$ such that $\mu_k(\{\infty\}) = 0$ which satisfies $T(f)(k) = \int f d\mu_k$ for all $f \in C(K_{\mathcal{A}})$. Since $K_{\mathcal{A}}$ is scattered, all Radon measures on $K_{\mathcal{A}}$ are atomic (see [17], 19.7.6); in particular, they have countable carriers. Let X be a countable set which includes the intersections of all the carriers of the μ_k 's with the $\{x_\beta : \beta < 2^\omega\}$. If $\beta \notin X$, then $\mu_k(\{x_\beta\}) = 0$ for all $k \in N$ and so

$$T(d_\beta^n)(k) = \int_{K_{\mathcal{A}} - \{x_\beta\}} \left[\sum_{i \in A_\beta^n} e_i + \chi_{\{x_\beta\}} \right] d\mu_k = \sum_{i \in A_\beta^n} T(e_i)(k),$$

since $\chi_{\{x_\beta\}}$ is Borel and so integrable with respect to any Radon measure, as required. $\square$

Before proceeding to Claim 2, note that $m_n^k = \mu_k(\{n\})$. This implies that $\sum_{n \in N} |m_n^k| \leq \|\mu_k\| \leq \|T\|$ for all $k \in N$. It follows that the sequence of the series $(m_n^k : n, k \in N)$ appears for cofinally many $\beta < 2^\omega$ on our list $(m_n^k(\beta) : n, m \in N, \beta < 2^\omega)$.

Claim 2. *There is a constant c and a countable set $Y \subseteq 2^\omega$ such that for every $\beta \in 2^\omega - Y$ we have*

$$\lim_{n \in A_\beta} g(n) = c.$$

Proof of the Claim. Suppose that the claim is false, i.e., for every countable $Y \subseteq 2^\omega$ there are $\alpha, \beta \in 2^\omega - Y$ and two rationals $q_1 < q_2$ such that

$$\liminf_{n \in A_\alpha} g(n) < q_1^{\alpha, \beta} < q_2^{\alpha, \beta} < \limsup_{n \in A_\beta} g(n).$$

Knowing that every element of the ideal is included in the union of finitely many generators, without loss of generality we may assume that there are two reals $r_1 < r_2$ such that

$$\forall A \in \mathcal{I}_{2^\omega} \exists k, l \in N - A \quad g(k) < r_1 < r_2 < g(l).$$

But $g(k) = m_k^k(\beta)$, $g(l) = m_l^l(\beta)$ for some $\beta < 2^\omega$, i.e., 4) is satisfied. As our initial enumeration has cofinally many repetitions we may assume that $\beta \notin X$ from Claim 1. Now Claim 1, 5), 6) and 7) imply that

$$T(\chi_{D_\beta})(k_n) < r_1 + \frac{r_2 - r_1}{3},$$

$$T(\chi_{D_\beta})(l_n) > r_2 - \frac{r_2 - r_1}{3},$$

which means that $T(\chi_{D_\beta})$ is discontinuous at the point of $K_{\mathcal{A}}$ corresponding to D_β , a contradiction. $\square$

Claim 3. *$T - cI$ has separable range.*

Proof of the Claim. Suppose the claim is false. Then there is an $\varepsilon > 0$ such that for every countable $Z \subseteq 2^\omega$ and every $l' \in N$ there are $\alpha, \beta \in 2^\omega - Z$ and $l \geq l'$ such that $|(T - cI)(d_\beta^l)(x_\alpha)| > \varepsilon$. By the continuity of the functions $(T - cI)(d_\beta^l)$, this means that $|(T - cI)(d_\beta^l)(k)| > \varepsilon$ for $k \in A_\alpha$ for sufficiently large k . Let X be

as in Claim 1 and suppose that $X \subseteq Z$. Let Y be as in Claim 2 and suppose that $Y \subseteq Z$. Claims 1 and 2 imply that

$$|T(d_\beta^l)(k) - cd_\beta^l(k)| = \left| \sum_{n \in A_\beta^l} [T(e_n)(k) - cd_\beta^l e_k] \right| = \left| \sum_{n \in A_\beta^l - \{k\}} m_n^k \right| > \varepsilon$$

if k is sufficiently large in A_α , since by Claim 2, then c is close to $g(k) = T(e_k)(k) = m_k^k$ and $d_\beta^l(k) = 0$ if $k \notin A_\beta^l$ and $d_\beta^l(k) = 1$ if $k \in A_\beta^l$ (note that we need to consider the case of $k \in A_\beta$ only because it could happen that $\alpha = \beta$). It follows that 1) is satisfied at stage β such that $m_n^k = m_n^k(\beta)$ for all $n, k \in N$. Since our enumeration has cofinally many repetitions and $cf(2^\omega) > \omega$ we may assume that $\beta \notin X, Y$ where X is as in Claim 1 and Y is as in Claim 2.

Thus 2) and 3) is satisfied for B_β and C_β . But this implies that $|T(\chi_{B_\beta})(k_i)| > 2\varepsilon/3$ if i is even and $|T(\chi_{B_\beta})(k_i)| < \varepsilon/3$ if i is odd, which contradicts the fact that $T(\chi_{B_\beta})$ must be continuous at the point corresponding to C_β . This completes the proof of the proposition. $\square$

Lemma 3 ([5], Example 2c). *Suppose that $A \subseteq C_0(K_A)$ is separable for some almost disjoint A . Then there is a closed subspace B such that $A \subseteq B$ and $B \sim c_0$.*

Lemma 4. *Suppose that $C_0(K_A) = A \oplus B$ where A is separable and B is not separable. Then B has a complemented copy of c_0 . In particular $A \sim c_0$ and $B \sim C_0(K_A)$.*

Proof. By Lemma 3, A is complemented in an isomorphic copy of c_0 ; thus by a result of [14] (Theorem 1), $A \sim c_0$. Well order $\mathcal{A}$ as $(A_\xi : \xi < \kappa)$ for some uncountable cardinal κ so that A is included in the closure of $\{\chi_{A_n} : n \in \omega\} \cup \{e_n : n \in N\}$. Let P be the projection on B and Q the projection on A . By induction construct $\omega < \beta_i < \beta'_i < \beta_{i+1} < \dots < \omega_1$ and $n_i \in N$ such that

$$8) \quad \|Q(d_{\beta_i}^{n_i}) - Q(d_{\beta'_i}^{n_i})\| \leq 1/2^i,$$

$$9) \quad (A_{\beta_i}^{n_i} \cup A_{\beta'_i}^{n_i}) \cap (A_{\beta_j}^{n_j} \cup A_{\beta'_j}^{n_j}) = \emptyset$$

for all $i < j < \omega$. This can be accomplished since $(A_\xi : \xi < \kappa)$ is almost disjoint and A is separable. We claim that

$$X = \left\{ \sum_{i \in N} a_i P(d_{\beta_i}^{n_i} - d_{\beta'_i}^{n_i}) : (a_i)_{i \in N} \in c_0 \right\}$$

is included in B , that it is isomorphic to c_0 and that it is complemented in $C_0(K_A)$ and hence in B .

First note that each element of X is a well-defined element of $C_0(K_A)$ because the series $\sum_{i \in N} a_i P(d_{\beta_i}^{n_i} - d_{\beta'_i}^{n_i})$ is norm convergent. For this we use the fact that $P + Q$ is the identity to write

$$10) \quad P(d_{\beta_i}^{n_i} - d_{\beta'_i}^{n_i}) = (d_{\beta_i}^{n_i} - d_{\beta'_i}^{n_i}) - [Q(d_{\beta_i}^{n_i}) - Q(d_{\beta'_i}^{n_i})]$$

and we note that $\sum_{i \in N} a_i (d_{\beta_i}^{n_i} - d_{\beta'_i}^{n_i})$ norm converges by 9) and $\sum_{i \in N} a_i Q(d_{\beta_i}^{n_i} - d_{\beta'_i}^{n_i})$ norm converges by 8). X is included in B because B is closed. The projection on X is

$$R(f) = \sum_{i \in N} f(x_{\beta_i}) \cdot P(d_{\beta_i}^{n_i} - d_{\beta'_i}^{n_i}).$$

It is linear and bounded and hence continuous by 10). By the fact that all elements of the space A are zero on x_{β_i} (since $\beta_i > \omega$), R is zero on A , and so it is true that

$$R(P(d_{\beta_i}^{n_i} - d_{\beta_i}^{m_i})) = R(d_{\beta_i}^{n_i} - d_{\beta_i}^{m_i}) = P(d_{\beta_i}^{n_i} - d_{\beta_i}^{m_i}),$$

i.e., R is the identity on X . To prove the last part of the lemma let $B \sim c_0 \oplus C$ (since B by the above has a complemented copy of c_0) and note that

$$C_0(K_{\mathcal{A}}) \sim A \oplus B \sim c_0 \oplus c_0 \oplus C \sim c_0 \oplus C \sim B.$$

□

Proof of the Theorem. Let $\mathcal{A}$ be as in Proposition 2. Since every operator on $C(K_{\mathcal{A}})$ is an operator on $C_0(K_{\mathcal{A}})$ plus an operator with finite-dimensional range (restriction projected on $C_0(K_{\mathcal{A}})$ by subtracting the value at ∞), the operators on $C(K_{\mathcal{A}})$ are of the desired form by Proposition 2.

Suppose that P is any projection on $C(K_{\mathcal{A}})$, in particular $P = P^2$, so if $P = cI + S$ where S has a separable range, we get $cI + S = c^2I + 2cS + S^2$, that is, $(c - c^2)I$ has separable range, which means that $c = c^2$. If $c = 0$, P has separable range; otherwise the projection on the complement has separable range since the sum of the projections is I . The conclusion is that the only decompositions of $C(K_{\mathcal{A}})$ are into one separable and one nonseparable summand. Lemma 4 implies that the separable summand is a copy of c_0 and the nonseparable summand is a copy of $C(K_{\mathcal{A}})$ as required. □

3. DECOMPOSITIONS OF $C(K_{\mathcal{A}})$ FOR SMALL $\mathcal{A}$ UNDER MARTIN'S AXIOM

Lemma 5. *Suppose that κ is an infinite cardinal. Assume MA_{κ} and suppose that an almost disjoint family $\mathcal{A} = \{A_{\xi} : \xi < \kappa\}$ is of cardinality κ . For every $X \subseteq \kappa$ there is a finite-to-one function $f : N \rightarrow N$ such that $f \circ f = f$ and 1) $(f[A_{\xi}] \cup f^{-1}[A_{\xi}]) \subseteq^* A_{\xi}$ for all $\xi \in X$, 2) $f[N] \cap A_{\eta}$ is finite for all $\eta \in \kappa - X$.*

Proof. First we define a partial order P to which we will be applying Martin's axiom. The conditions of P are of the form $p = (a_p, n_p, f_p)$, where $a_p \in [\kappa]^{<\omega}$, $n_p \in N$, $f_p : n_p \rightarrow n_p$ such that $f_p \circ f_p = f_p$ and

- a) for all distinct $\xi, \eta \in a_p$ we have $A_{\xi} \cap A_{\eta} \subseteq n_p$. The order on P is defined by $p \leq q$ if and only if $a_p \supseteq a_q$, $n_p \geq n_q$, $f_p \supseteq f_q$ and
- b) $f_p^{-1}(\{n\}) = f_q^{-1}(\{n\})$ for all $n < n_q$,
- c) for $\xi \in a_q \cap X$ we have $n \in A_{\xi} \cap [n_q, n_p]$ iff $f(n) \in A_{\xi} \cap [n_q, n_p]$,
- d) $f[[n_q, n_p]] \cap A_{\eta} = \emptyset$ for $\eta \in a_q - X$.

Note that the above relation is a partial order. To be able to take advantage of Martin's axiom, we must prove that P satisfies the countable chain condition. Suppose that an uncountable family of conditions of P is given. We may assume without loss of generality that their second coordinates are all equal to some integer n and that all third coordinates are equal to some $f : n \rightarrow n$. So it is enough to prove that any two conditions q, r with $n = n_q = n_r$ and $f = f_q = f_r$ are compatible.

Find $n_p > n$ so that for all $\xi \in (a_q \cup a_r) \cap X$ there is an $n_{\xi} \in [n, n_p]$ which does not belong to any A_{η} for η distinct from ξ and $\eta \in a_q \cup a_r$; moreover, $[n, n_p]$ should contain an m which does not belong to any of the sets A_{ξ} for $\xi \in a_q \cup a_r$, and we also require that $A_{\xi} \cap A_{\eta} \subseteq n_p$ for distinct $\xi, \eta \in a_q \cup a_r$. This can be accomplished because the sets are almost disjoint. Put $a_p = a_q \cup a_r$. For $k \in n$ put $f_p(k) = f_q(k) = f_r(k)$ and for $k \in [n, n_p]$ define $f_p(k)$ as follows:

- i) if $k \in A_{\xi} \cap A_{\xi'}$ for distinct $\xi, \xi' \in (a_q \cup a_r) \cap X$, then $f_p(k) = k$;

- ii) if $n \in A_\xi$ for $\xi \in (a_q \cup a_r) \cap X$, but i) doesn't hold, then $f(k) = n_\xi$;
- iii) if neither i) nor ii) holds for k , then $f(k) = m$.

It is clear that $p = (a_p, n, f)$ is a condition of P . Let us check that $p \leq q, r$. Conditions b) and c) are clear from the construction.

The key observation needed to prove d) is that if i) holds for k , then $\xi \in a_q - a_r$ and $\xi' \in a_r - a_q$. This is because condition a) for q or r implies that the intersections $A_\xi \cap A_{\xi'}$ for $\xi, \xi' \in a_q$ or $\xi, \xi' \in a_r$ are included in n . Consequently, again applying a) in a similar way for q or r , already knowing that $k \in A_\xi \cap A_{\xi'}$, we conclude that if i) holds for k , then k does not belong to any A_η for $\eta \in a_p - X$, which implies d).

A similar but simpler argument shows that the sets $D_n = \{p \in P : n_p \geq n\}$ are dense in P for $n \in N$. The sets $E_\xi = \{p : \xi \in a_p\}$ are dense in P for each $\xi \in \kappa$ because by the density of D_n 's one can get n_p large enough so that adding ξ to a_p does not violate a).

By Martin's axiom there is a filter $G \subseteq P$ which intersects all sets E_ξ and D_n for $n \in N$ and $\xi \in \kappa$. Then $\bigcup \{f_p : p \in G\}$ is a function that satisfies the lemma. Indeed, c) implies 1) and d) implies 2). $\square$

Proposition 6. *Assume MA_κ and suppose that $\mathcal{A} = \{A_\xi : \xi < \kappa\}$ is an almost disjoint family. For any $X \subseteq \kappa$ there is a decomposition $C(K_\mathcal{A}) = A \oplus B$ such that for suitable $\{n_\xi : \xi < \kappa\} \subseteq N$ we have $d_\xi^{n_\xi} \in A$ for $\xi \in X$ and $d_\xi^{n_\xi} \in B$ for $\xi \in \kappa - X$. In particular if both X and $\kappa - X$ are uncountable, then both of the factors are non-separable.*

Proof. Let f be as in Lemma 5 for X . Define $r : K_\mathcal{A} \rightarrow K_\mathcal{A}$ in the following way: $r(n) = f(n)$ for $n \in N$, $r(x_\xi) = x_\xi$ for $\xi \in X$, $r(x_\xi) = \infty$ for $\xi \in \kappa - X$ and $r(\infty) = \infty$.

Let us check that r is a retraction. It is clear that $r \circ r = r$, so we need to check that it is continuous. It is vacuously true at points of N . If $n_i \rightarrow x_\xi$ for $\xi \in X$, then $f(n_i) \rightarrow x_\xi = f(x_\xi)$ by 1) of Lemma 5 and by the fact that f is finite-to-one. If $n_i \rightarrow x_\xi$ for $\xi \in \kappa - X$, then $f(n_i) \rightarrow \infty = f(x_\xi)$ by 2) of Lemma 5, by the part $f^{-1}[A_\xi] \subseteq^* A_\xi$ for $\xi \in X$ of 1) of Lemma 5 and by the fact that f is finite-to-one. Finally if $M \subseteq N$ has ∞ in its closure, then its image cannot be included in finitely many elements of the almost disjoint family by Lemma 5, i.e. the image has ∞ in its closure as well, i.e., f is continuous at ∞ as well.

Let $K = f[N] \cup \{x_\xi : \xi \in X\} \cup \{\infty\} \subseteq K_\mathcal{A}$ be the image of r . It is well known ([17], §21) that $C(K)$ is isomorphic to a subspace A of $C(K_\mathcal{A})$ defined by $\{f \circ r : f \in C(K)\}$ and that $P(f) = (f|K) \circ r = f \circ r$ is a projection witnessing the fact that A is complemented in $C(K_\mathcal{A})$. Note that $P(d_\xi^{n_\xi}) = d_\xi^{n_\xi}$ for some $n_\xi \in N$ if $\xi \in X$ by 1) of Lemma 5) and that $P(d_\xi^{n_\xi}) = 0$ if $A_\xi - n_\xi$ is disjoint from $f[N]$ which can be accomplished by 2) of Lemma 5) for $\xi \in \kappa - X$. $\square$

ACKNOWLEDGMENT

We are thankful to Professor Joan Bagaria for organizing the visit at CRM.

REFERENCES

- [1] D. Alspach, Y. Benyamini; Primariness of spaces of continuous functions on ordinals; Israel J. Math. vol 27, No 1, 1977, pp. 64 - 92. MR0440349 (55:13224)
- [2] J. Diestel; Sequences and series in Banach spaces; Springer-Verlag 1984. MR0737004 (85i:46020)

- [3] E. van Douwen; The Integers and Topology; Handbook of Set-theoretic Topology; eds K. Kunen, J. Vaughan, North Holland 1984; pp. 111 - 167. MR0776622 (87f:54008)
- [4] W. T. Gowers, B. Maurey; The unconditional basic sequence problem; Journal A. M. S. 6 (1993), pp. 851-874. MR1201238 (94k:46021)
- [5] W. Johnson, J. Lindenstrauss; Some remarks on weakly compactly generated Banach spaces; Israel J. Math. 17, 1974, pp. 219 - 230. and Israel J. Math. 32 (1979), no. 4, pp. 382 - 383. MR0417760 (54:5808); MR0571092 (81g:46015)
- [6] P. Koszmider; Banach spaces of large densities but few operators. Preprint.
- [7] P. Koszmider; Banach spaces of continuous functions with few operators; Math. Annalen. 330, 2004, pp. 151 - 183. MR2091683
- [8] K. Kunen; Set Theory. An Introduction to Independence Proofs; North Holland, 1980. MR0597342 (82f:03001)
- [9] Y. Lindenstrauss, A. Pełczyński; Contributions to the theory of the classical Banach spaces; J. Funct. Anal. 8, 1971, pp. 225 - 249. MR0291772 (45:863)
- [10] N. Luzin; On subsets of the series of natural numbers; Izv. Akad. Nauk SSSR, Ser. Mat., 11, pp. 403 - 411. MR0021576 (9:82c)
- [11] A. Miller; private notes, 2003.
- [12] A. Moltó; On a theorem of Sobczyk; Bull. Austral. Math. Soc. 43, 1991, 123 - 130. MR1086724 (92d:46043)
- [13] S. Mrówka; Some set-theoretic constructions in topology; Fund. Math. vol. XCIV, 1977, pp. 83 - 92. MR0433388 (55:6364)
- [14] A. Pełczyński; Projections in certain Banach spaces; Studia Math. vol. XIX, 1960, pp. 209 - 228. MR0126145 (23:A3441)
- [15] A. Pełczyński, Z. Semadeni; Spaces of continuous functions (III) (Spaces $C(\Omega)$ for Ω without perfect subsets); Studia Math. 18, 1959, pp. 211 - 222. MR0107806 (21:6528)
- [16] H. Rosenthal; On relatively disjoint families of measures with some applications to Banach space theory; Studia Math. 37, (1970), pp. 13-36. MR0270122 (42:5015)
- [17] Z. Semadeni; Banach spaces of continuous functions; Państwowe Wydawnictwo Naukowe, 1971. MR0296671 (45:5730)
- [18] S. Shelah; A Banach space with few operators; Israel J. Math. 30 (1978), pp. 181-191. MR0508262 (80b:46033)
- [19] S. Shelah, J. Steprans; A Banach space on which there are few operators; Proc. Amer. Math. Soc. 104 (1988), pp. 101-105. MR0958051 (90a:46047)
- [20] H. Wark; A non-separable reflexive Banach space on which there are few operators. J. London Math. Soc. (2) 64 (2001), no. 3, pp. 675 - 689. MR1865556 (2003a:46031)

DEPARTAMENTO DE MATEMÁTICA, UNIVERSIDADE DE SÃO PAULO, CAIXA POSTAL: 66281, SÃO PAULO, SP CEP: 05315-970, BRAZIL

E-mail address: `piotr@ime.usp.br`