A NOTE ON EXPONENTIAL DECAY PROPERTIES OF GROUND STATES FOR QUASILINEAR ELLIPTIC EQUATIONS

YI LI AND CHUNSHAN ZHAO

(Communicated by David S. Tartakoff)

ABSTRACT. We give an explicit formula for exponential decay properties of ground states for a class of quasilinear elliptic equations in the whole space $\mathbb{R}^N$.

1. INTRODUCTION

We consider exponential decay properties of ground states of the quasilinear elliptic equation

$$(1.1) \quad \Delta_m u + f(u) = 0 \quad \text{in } \mathbb{R}^N, \quad N > m > 1,$$

where $\Delta_m u = \operatorname{div}(|Du|^{m-2} Du)$ is the degenerate m -Laplace operator. Here by a ground state we mean a non-negative non-trivial C^1 distribution solution of (1.1) which tends to zero at ∞ .

Remark 1. There are some sufficient conditions which guarantee the existence of a positive radial ground state $u(x)$ satisfying $u(x) = u(r)$ with $r = |x|$, $u(0) = \max_{x \in \mathbb{R}^N} u(x)$, $u'(0) = 0$ and $u'(r) < 0$ for $r > 0$, where $u'(r) = \frac{du(r)}{dr}$. See, for example, (H1)—(H3) in reference [6]. A good example for $f(u)$ which satisfies (A1) is:

$$f(u) = -u^{m-1} + u^q \quad \text{with } m-1 < q < \frac{N(m-1)+m}{N-m}.$$

From now on we make the following assumption on f :

(A1): $f: \mathbb{R}^+ \rightarrow \mathbb{R}$ is continuous and there exist positive constants k and δ such that $f(z) + kz^{m-1} = O(z^{m-1+\delta})$ as $z \downarrow 0$.

Before stating our results, let us recollect some facts about ground states for (1.1) with $m = 2$, i.e.,

$$(1.2) \quad \Delta u + f(u) = 0 \quad \text{in } \mathbb{R}^N.$$

Received by the editors February 25, 2004.

2000 *Mathematics Subject Classification.* Primary 35B40, 35J70.

Key words and phrases. m -Laplace operator, ground states, exponential decay.

The first author was supported in part by the NSFC (10471052) and the Xiao-Xiang Funds of Hunan Normal University.

Under suitable conditions on f (e.g. [2]) it is well known that the ground state for (1.2) satisfies

$$u(x) = u(|x|) = u(r) > 0, \quad u(0) = \max_{x \in \mathbb{R}^N} u(x),$$

$$u'(0) = 0 \quad \text{and} \quad u'(r) < 0 \quad \text{for} \quad r \in (0, \infty).$$

Moreover,

$$(1.3) \quad \lim_{r \rightarrow \infty} u(r) r^{\frac{N-1}{2}} e^r = C_1$$

where $0 < C_1 < \infty$ is a constant, and such precise estimates of asymptotes have been proved to be very useful. For example, for applications of such estimates for $m = 2$, readers can refer to [1], [2], [3], [4], [5], etc. In this note we show that a similar estimate to (1.3) also holds for radial ground states of (1.1) if assumption (A1) is satisfied. Our main result is as follows.

Theorem 1.1. *Let $u(x) = u(r)$ be a positive radial ground state for (1.1) and let $f(z)$ satisfy assumptions (A1). Then there exists a sequence of constants $\{c_i\}$ ($i = 1, 2, \dots$) such that*

$$\left(-\frac{u'}{u}\right)^{m-1} = \varphi_\infty + \frac{c_1}{r} + \frac{c_2}{r^2} + \dots + \frac{c_l}{r^l} + \dots \quad \text{as } r \rightarrow \infty$$

where $\{c_i\}$ ($i = 1, 2, \dots$) are determined by

$$c_1 = \frac{N-1}{m} \left(\frac{k}{m-1}\right)^{\frac{m-2}{m}}, \quad c_2 = \frac{(N-2)c_1 - \frac{m}{2(m-1)}\varphi_\infty^{\frac{2-m}{m-1}}c_1^2}{m\varphi_\infty^{\frac{1}{m}}},$$

and for $l > 2$, c_l can be uniquely determined by

$$(N-l)c_{l-1} - m\varphi_\infty^{\frac{1}{m-1}}c_l = \sum_{i=2}^l \frac{F^{(i)}(0)}{i!} \left(\sum_{i_1+\dots+i_i=l, i_1, \dots, i_i > 0} c_{i_1}c_{i_2}\dots c_{i_i} \right),$$

where $\varphi_\infty = \left(\frac{k}{m-1}\right)^{\frac{m-1}{m}}$ and $F(\rho) = (m-1)(\varphi_\infty + \rho)^{\frac{m}{m-1}}$. In particular we have

$$\lim_{r \rightarrow \infty} u(r) r^{\frac{N-1}{m(m-1)}} e^{\left(\frac{k}{m-1}\right)^{\frac{1}{m}} r} = C_2$$

for some $0 < C_2 < \infty$.

2. PROOF OF THEOREM 1.1

In this section, we will present the proof of Theorem 1.1. First let $u(x) = u(r)$ be a radial ground state as stated in Theorem 1.1. It follows from (1.1) that

$$(2.1) \quad \left(|u'|^{m-2} u'\right)' + \frac{N-1}{r} |u'|^{m-2} u' + f(u) = 0.$$

Let $\varphi(r) = -\frac{|u'|^{m-2} u'}{u^{m-1}}$. Then $|u'|^{m-2} u' = -\varphi u^{m-1}$ and $\frac{u'}{u} = -\varphi^{\frac{1}{m-1}}$. Substituting them into (2.1) yields

$$(2.2) \quad \varphi' - (m-1)\varphi^{\frac{m}{m-1}} + \frac{N-1}{r}\varphi - \frac{f(u)}{u^{m-1}} = 0.$$

Lemma 2.1. $\limsup_{r \rightarrow \infty} \varphi(r) < \infty$.

Proof. It follows from assumption (A1) and $\lim_{r \rightarrow \infty} u(r) = 0$ that

$$K = \sup_r \frac{|f(u(r))|}{u^{m-1}(r)} < \infty.$$

Noticing from $\varphi(r) \geq 0$ and (2.2) we obtain that as long as $\varphi(r) \geq \left(\frac{4K}{m-1}\right)^{\frac{m-1}{m}}$ and $r \geq \frac{4(N-1)}{m-1} \left(\frac{m-1}{4K}\right)^{\frac{m-1}{m}}$,

$$(2.3) \quad \varphi' = (m-1)\varphi^{\frac{m}{m-1}} - \frac{N-1}{r}\varphi + \frac{f(u)}{u^{m-1}} \geq \frac{m-1}{2}\varphi^{\frac{m}{m-1}}.$$

Now, suppose to the contrary that $\limsup_{r \rightarrow \infty} \varphi(r) = \infty$. Let

$$r_1 = \inf\left\{r \geq \frac{4(N-1)}{m-1} \left(\frac{m-1}{4K}\right)^{\frac{m-1}{m}} : \varphi(r) \geq \left(\frac{4K}{m-1}\right)^{\frac{m-1}{m}}\right\}.$$

Since $\varphi(0) = 0$ and $\varphi(r) > 0$ for $r > 0$, it follows that $0 < r_1 \leq \infty$. If $r_1 = \infty$ we are done. Next we suppose $r_1 < \infty$. Then $\varphi(r_1) = \left(\frac{4K}{m-1}\right)^{\frac{m-1}{m}}$ and $\varphi'(r) \geq \frac{m-1}{2}\varphi(r)^{\frac{m}{m-1}}$ for all $r \geq r_1$, which blows up before or at $r_2 = r_1 + 2\left(\frac{m-1}{4K}\right)^{\frac{1}{m}}$. This contradicts the well-definedness of φ . The proof of this lemma is completed. $\square$

Lemma 2.2.

$$(2.4) \quad \lim_{r \rightarrow \infty} \varphi = \varphi_\infty = \left(\frac{k}{m-1}\right)^{\frac{m-1}{m}}.$$

Proof. Since $\varphi(0) = 0$ and $\varphi(r) > 0$ for $r > 0$ it follows from (2.2) and Lemma 2.1 that

$$0 \leq \liminf_{r \rightarrow \infty} \varphi(r) = \alpha < \infty$$

and

$$0 < \limsup_{r \rightarrow \infty} \varphi(r) = \beta < \infty.$$

Next we use contradiction arguments to prove (2.4). Suppose

$$\alpha = \liminf_{r \rightarrow \infty} \varphi(r) < \limsup_{r \rightarrow \infty} \varphi(r) = \beta.$$

Then we may choose two sequences $\{\eta_i\}$ and $\{\zeta_i\}$ going to ∞ as $i \rightarrow \infty$ such that

$$\{\eta_i\} \text{ are local minima of } \varphi$$

and

$$\{\zeta_i\} \text{ are local maxima of } \varphi$$

and

$$\eta_i < \zeta_i < \eta_{i+1} \text{ and } \liminf_{r \rightarrow \infty} \varphi(r) = \lim_{i \rightarrow \infty} \varphi(\eta_i) = \alpha, \limsup_{r \rightarrow \infty} \varphi(r) = \lim_{i \rightarrow \infty} \varphi(\zeta_i) = \beta.$$

Then at η_i ($i = 1, 2, \dots$) we know $u'(\eta_i) = 0$ and thus

$$(2.5) \quad -(m-1)\varphi(\eta_i)^{\frac{m}{m-1}} + \frac{N-1}{\eta_i}\varphi(\eta_i) - \frac{f(u(\eta_i))}{u(\eta_i)^{m-1}} = 0.$$

Similarly at ζ_i ($i = 1, 2, \dots$) we get

$$(2.6) \quad -(m-1)\varphi(\zeta_i)^{\frac{m}{m-1}} + \frac{N-1}{\zeta_i}\varphi(\zeta_i) - \frac{f(u(\zeta_i))}{u(\zeta_i)^{m-1}} = 0,$$

$\forall \epsilon > 0$. Since $\lim_{z \downarrow 0} \frac{f(z)}{z^{\frac{m}{m-1}}} = -k$ and $\lim_{r \rightarrow \infty} u(r) = 0$, we can take r_3 sufficiently large such that

$$\left| \frac{f(u(r))}{u(r)^{\frac{m}{m-1}}} + k \right| < \epsilon \quad \text{for all } r > r_3.$$

Next we take i_0 sufficiently large such that

$$\eta_i > r_3 \text{ for } i > i_0.$$

From (2.5) and (2.6) it follows that for all $i > i_0$:

$$k - \epsilon < (m-1)\varphi(\eta_i)^{\frac{m}{m-1}} - \frac{N-1}{\eta_i}\varphi(\eta_i) < k + \epsilon$$

and

$$k - \epsilon < (m-1)\varphi(\zeta_i)^{\frac{m}{m-1}} - \frac{N-1}{\zeta_i}\varphi(\zeta_i) < k + \epsilon.$$

Letting $i \rightarrow \infty$ and noticing the arbitrariness of ϵ , we get

$$(m-1)\alpha^{\frac{m}{m-1}} = k, \quad (m-1)\beta^{\frac{m}{m-1}} = k,$$

which yields $\alpha = \beta$, contradicting $\alpha < \beta$.

Therefore

$$\lim_{r \rightarrow \infty} \varphi(r) = \varphi_\infty = \left(\frac{k}{m-1} \right)^{\frac{m-1}{m}}.$$

The proof of this lemma is complete. $\square$

After presenting the above two lemmas we can now give a proof of our main result.

Proof of Theorem 1.1. We deduce from Lemma 2.2 that

$$\lim_{r \rightarrow \infty} \frac{u'}{u} = \lim_{r \rightarrow \infty} -\varphi^{\frac{1}{m-1}} = -\left(\frac{k}{m-1} \right)^{\frac{1}{m}}.$$

Therefore $\forall \epsilon > 0$, there exists a constant $0 < c(\epsilon) < \infty$ such that

$$u(r) \leq c(\epsilon)e^{-\left(\frac{k-\epsilon}{m-1}\right)^{\frac{1}{m}}r}.$$

In particular taking $\epsilon = \frac{k}{2}$ we have

$$u(r) \leq c(k)e^{-\left(\frac{k}{2(m-1)}\right)^{\frac{1}{m}}r}.$$

For convenience, let $\left(\frac{k}{2(m-1)}\right)^{\frac{1}{m}} = \mu$. We know

$$(2.7) \quad u(r) \leq ce^{-\mu r}.$$

Next we give a more precise expansion of $\varphi(r)$ at ∞ . Let $\varphi = \varphi_\infty + \varphi_1$. We know from (2.2) and Lemma 2.2 that $\lim_{r \rightarrow \infty} \varphi_1(r) = 0$ and $\varphi_1(r)$ satisfies

$$\varphi_1' - (m-1)(\varphi_\infty + \varphi_1)^{\frac{m}{m-1}} + \frac{N-1}{r}(\varphi_\infty + \varphi_1) - \frac{f(u)}{u^{m-1}} = 0,$$

i.e.,

$$(2.8) \quad \begin{aligned} & \varphi_1' - m\varphi_\infty^{\frac{1}{m-1}}\varphi_1 + \frac{N-1}{r}\varphi_1 \\ &= \frac{f(u)}{u^{m-1}} + (m-1)(\varphi_\infty + \varphi_1)^{\frac{m}{m-1}} - m\varphi_\infty^{\frac{1}{m-1}}\varphi_1 - \frac{N-1}{r}\varphi_\infty. \end{aligned}$$

Noticing that $\lim_{r \rightarrow \infty} u(r) = 0$ and $\frac{f(z) + kz^{m-1}}{z^{m-1}} = O(z^\delta)$ as $z \downarrow 0$, we get for r sufficiently large that

$$(2.9) \quad \frac{f(u)}{u^{m-1}} = -k + O(u^\delta).$$

At the same time for r sufficiently large we have

$$(2.10) \quad (m-1)(\varphi_\infty + \varphi_1)^{\frac{m}{m-1}} = (m-1)\varphi_\infty^{\frac{m}{m-1}} + m\varphi_\infty^{\frac{1}{m-1}}\varphi_1 + O(\varphi_1^2).$$

Thus from (2.8)–(2.10) it follows that for r sufficiently large,

$$(2.11) \quad \varphi_1' - m\varphi_\infty^{\frac{1}{m-1}}\varphi_1 + \frac{N-1}{r}\varphi_1 = O(\varphi_1^2) - \frac{N-1}{r}\varphi_\infty + O(u^\delta).$$

Multiplying each side of (2.11) by φ_1 and integrating from r to ∞ for r sufficiently large, we get

$$(2.12) \quad \begin{aligned} & \frac{1}{2}\varphi_1^2(r) + \int_r^\infty \left(m\varphi_\infty^{\frac{1}{m-1}} - \frac{N-1}{s} + O(\varphi_1) \right) \varphi_1^2 ds \\ &= \int_r^\infty \frac{N-1}{s} \varphi_\infty \varphi_1 ds - \int_r^\infty O(u^\delta) \varphi_1 ds. \end{aligned}$$

We can take r sufficiently large such that

$$m\varphi_\infty^{\frac{1}{m-1}} - \frac{N-1}{s} + O(\varphi_1) \geq \frac{1}{2}m\varphi_\infty^{\frac{1}{m-1}} \text{ for } s > r.$$

Therefore for such large r it follows that

$$\varphi_1^2(r) + \int_r^\infty \left(m\varphi_\infty^{\frac{1}{m-1}} \right) \varphi_1^2 ds \leq 2 \int_r^\infty \frac{N-1}{s} \varphi_\infty \varphi_1 ds - 2 \int_r^\infty O(u^\delta) \varphi_1 ds.$$

Notice that

$$2 \int_r^\infty \frac{N-1}{s} \varphi_\infty \varphi_1 ds \leq \frac{1}{4} \left(m\varphi_\infty^{\frac{1}{m-1}} \right) \int_r^\infty \varphi_1^2 ds + \frac{4(N-1)^2 \varphi_\infty^2}{m\varphi_\infty^{\frac{1}{m-1}}} \int_r^\infty \frac{1}{s^2} ds$$

and

$$2 \int_r^\infty O(u^\delta) \varphi_1 ds \leq \frac{1}{4} \left(m\varphi_\infty^{\frac{1}{m-1}} \right) \int_r^\infty \varphi_1^2 ds + \frac{4C}{m\varphi_\infty^{\frac{1}{m-1}}} \int_r^\infty O(u^{2\delta}) ds.$$

By virtue of the above estimates and (2.7) we obtain for r sufficiently large,

$$(2.13) \quad \begin{aligned} & \varphi_1^2(r) + \frac{1}{2} \left(m\varphi_\infty^{\frac{1}{m-1}} \right) \int_r^\infty \varphi_1^2(s) ds \\ & \leq \frac{4(N-1)^2 \varphi_\infty^2}{m\varphi_\infty^{\frac{1}{m-1}}} \frac{1}{r} + \frac{4c}{2\delta m\varphi_\infty^{\frac{1}{m-1}}} e^{-2\delta\mu r} \\ & \leq \frac{8(N-1)^2 \varphi_\infty^2}{m\varphi_\infty^{\frac{1}{m-1}}} \frac{1}{r}. \end{aligned}$$

Thus we have

$$(2.14) \quad \varphi_1^2(r) = O\left(\frac{1}{r}\right) \text{ as } r \rightarrow \infty.$$

By this estimate and (2.11) it follows that as $r \rightarrow \infty$,

$$(2.15) \quad \varphi_1' - m\varphi_\infty^{\frac{1}{m-1}}\varphi_1 + \frac{N-1}{r}\varphi_1 = O\left(\frac{1}{r}\right).$$

For convenience let $\alpha_0 = m\varphi_\infty^{\frac{1}{m-1}}$. Then we get from (2.15) as $r \rightarrow +\infty$,

$$(2.16) \quad (r^{N-1}e^{-\alpha_0 r}\varphi_1)' = r^{N-1}e^{-\alpha_0 r}O\left(\frac{1}{r}\right).$$

Integrating both sides of (2.16) from r to ∞ yields as $r \rightarrow \infty$,

$$(2.17) \quad \begin{aligned} \varphi_1(r) &= \frac{e^{\alpha_0 r}}{r^{N-1}} \int_r^\infty s^{N-1} O\left(\frac{1}{s}\right) e^{-\alpha_0 s} ds \\ &= O\left(\frac{e^{\alpha_0 r}}{r^{N-1}} \int_r^\infty s^{N-2} e^{-\alpha_0 s} ds\right). \end{aligned}$$

Applying integration by parts as many steps as we want we arrive that there exists a sequence of constants $\{a_i\}$ ($i = 1, 2, \dots$) such that

$$(2.18) \quad \begin{aligned} \int_r^\infty s^{N-2} e^{-\alpha_0 s} ds &= a_1 r^{N-2} e^{-\alpha_0 r} + a_2 r^{N-3} e^{-\alpha_0 r} + \dots \\ &= e^{-\alpha_0 r} (a_1 r^{N-2} + a_2 r^{N-3} + \dots + a_l r^{N-l-1} + \dots) \end{aligned}$$

where $a_1 = \frac{1}{\alpha_0}$. Thus it follows from (2.16) that

$$(2.19) \quad \varphi_1(r) = O\left(\frac{1}{r}\right), \text{ as } r \rightarrow \infty,$$

which is an improvement of (2.14). Using (2.19) and (2.11) we obtain

$$(2.20) \quad (r^{N-1}e^{-\alpha_0 r}\varphi_1)' = -r^{N-1}e^{-\alpha_0 r} \left(\frac{N-1}{r} \varphi_\infty + O\left(\frac{1}{r^2}\right) \right).$$

Similar to (2.16) and (2.17), we arrive at

$$\begin{aligned} \varphi_1(r) &= \frac{e^{\alpha_0 r}}{r^{N-1}} \int_r^\infty s^{N-1} e^{-\alpha_0 s} \left(\frac{N-1}{s} \varphi_\infty + O\left(\frac{1}{s^2}\right) \right) ds \\ &= \frac{(N-1)\varphi_\infty a_1}{r} + O\left(\frac{1}{r^2}\right). \end{aligned}$$

Again if we let $\varphi_1 = \frac{(N-1)\varphi_\infty a_1}{r} + \varphi_2$ such that $\varphi_2(r) = O\left(\frac{1}{r^2}\right)$, we obtain from (2.10) and (2.11) that

$$\varphi_2' - \alpha_0 \varphi_2 + \frac{N-1}{r} \varphi_2 = \varphi_\infty^{\frac{m-2}{m-1}} \left[\frac{N-1}{2m(m-1)} - \frac{N-2}{m} \right] \frac{N-1}{r^2} + O\left(\frac{1}{r^2}\right).$$

We can then repeat the same process to obtain the expansion as stated in Theorem 1.1 to any polynomial order as we want. Next we need to determine c_i ($i = 1, 2, \dots$) in Theorem 1.1. Let $F(\rho) = (m-1)(\varphi_\infty + \rho)^{\frac{m}{m-1}}$. Then the Taylor expansion of $F(\rho)$ at $\rho = 0$ is as follows:

$$(2.21) \quad \begin{aligned} F(\rho) &= (m-1)\varphi_\infty^{\frac{m}{m-1}} + \alpha_0 \rho + \frac{m}{2(m-1)} \varphi_\infty^{\frac{m-2}{m-1}} \rho^2 \\ &\quad - \frac{m(m-2)}{3!(m-1)^2} \varphi_\infty^{\frac{-2m-3}{m-1}} \rho^3 + \dots + \frac{F^{(n)}(0)}{n!} \rho^n + \dots \end{aligned}$$

where

$$F^{(n)}(0) = \frac{(-1)^n m(m-2)(2m-3) \cdots (lm-l-1) \cdots [(n-2)m-n+1]}{(m-1)^{n-1}}$$

for $n \geq 4$. Thus from (2.8), (2.9) and (2.21) we get

$$(2.22) \quad \begin{aligned} \varphi_1' - \alpha_0 \varphi_1 + \frac{N-1}{r} \varphi_1 &= O(u^\delta) - \frac{N-1}{r} \varphi_\infty \\ &+ \frac{m}{2(m-1)} \varphi_\infty^{-\frac{m-2}{m-1}} \varphi_1^2 - \frac{m(m-2)}{3!(m-1)^2} \varphi_\infty^{-\frac{2m}{m-1}} \varphi_1^3 + \cdots + \frac{F^{(n)}(0)}{n!} \varphi_1^n + \cdots. \end{aligned}$$

Substituting

$$\varphi_1(r) = \frac{c_1}{r} + \frac{c_2}{r^2} + \cdots + \frac{c_l}{r^l} + \cdots$$

into (2.22) we get by comparing the coefficients of $\frac{1}{r^n}$ ($n = 1, 2, \dots$) that

$$c_1 = \frac{N-1}{m} \left(\frac{k}{m-1} \right)^{\frac{m-2}{m}}$$

and

$$c_2 = \frac{(N-2)c_1 - \frac{m}{2(m-1)} \varphi_\infty^{-\frac{m-2}{m-1}} c_1^2}{\alpha_0}.$$

c_l ($l > 2$) is determined by

$$(N-l)c_{l-1} - \alpha_0 c_l = \sum_{i=2}^l \frac{F^{(i)}(0)}{i!} \left(\sum_{i_1+i_2+\cdots+i_i=l} c_{i_1} c_{i_2} \cdots c_{i_i} \right).$$

Notice that

$$\frac{u'}{u} = -\varphi^{\frac{1}{m-1}}.$$

We know that as $r \rightarrow \infty$,

$$\begin{aligned} \frac{u'}{u} &= - \left(\varphi_\infty^{\frac{1}{m-1}} + \frac{1}{m-1} \varphi_\infty^{\frac{2-m}{m-1}} c_1 \cdot \frac{1}{r} + O\left(\frac{1}{r^2}\right) \right) \\ &= - \left(\frac{k}{m-1} \right)^{\frac{1}{m}} - \frac{N-1}{m(m-1)} \cdot \frac{1}{r} + O\left(\frac{1}{r^2}\right), \end{aligned}$$

which yields that

$$u(r) = O \left(r^{-\frac{(N-1)}{m(m-1)}} e^{-\left(\frac{k}{m-1}\right)^{\frac{1}{m}} r} \right) \quad \text{as } r \rightarrow \infty,$$

i.e.,

$$\lim_{r \rightarrow \infty} u(r) r^{\frac{N-1}{m(m-1)}} e^{\left(\frac{k}{m-1}\right)^{\frac{1}{m}} r} = C_2,$$

for some $0 < C_2 < \infty$. The proof of Theorem 1.1 is complete. $\square$

REFERENCES

- [1] S. Agmon, Lectures on exponential decay of solutions of second-order elliptic equations: bounds on eigenfunctions of N -body Schrödinger operators. Mathematical Notes, 29. Princeton University Press, Princeton, NJ; University of Tokyo Press, Tokyo, 1982. 118 pp. MR0745286 (85f:35019)
- [2] B. Gidas, W. M. Ni, L. Nirenberg, *Symmetry of positive solutions of nonlinear elliptic equations in R^n* , Mathematical analysis and applications, Part A, pp. 369–402, Academic Press, New York-London, 1981. MR0634248 (84a:35083)
- [3] Y. Li, *Asymptotic behavior of positive solutions of equations $\Delta u + K(x)u^p = 0$ in R^n* , J. Differential Equations 95 (1992), No. 2, pp. 304–330. MR1165425 (93k:35048)
- [4] W. M. Ni, I. Takagi, On the shape of least energy solutions to a semi-linear Neumann problem, Comm. Pure Appl. Math 44 (1991), No. 7, 819–851. MR1115095 (92i:35052)

- [5] W. M. Ni, J. Wei, On the location and profile of spike-layer solutions to singularly perturbed semi-linear Dirichlet problems, *Comm. Pure Appl. Math.* 48 (1995), No. 7, 731–768. MR1342381 (96g:35077)
- [6] J. Serrin, M. Tang, *Uniqueness of ground states for quasilinear elliptic equations*, *Indiana Univ. Math. J.* 49 (2000) No. 3, pp. 897–923. MR1803216 (2002d:35072)

DEPARTMENT OF MATHEMATICS, HUNAN NORMAL UNIVERSITY, CHANGSHA, PEOPLE'S REPUBLIC OF CHINA – AND – DEPARTMENT OF MATHEMATICS, THE UNIVERSITY OF IOWA, IOWA CITY, IOWA 52242

E-mail address: yli@math.uiowa.edu

DEPARTMENT OF MATHEMATICS, THE UNIVERSITY OF IOWA, IOWA CITY, IOWA 52242

E-mail address: chuzhao@math.uiowa.edu