ON FINITE-SHEETED COVERING MAPPINGS ONTO SOLENOIDS

R. N. GUMEROV

(Communicated by Alan Dow)

ABSTRACT. We study limit mappings from a solenoid onto itself. It is shown that each equivalence class of finite-sheeted covering mappings from connected topological spaces onto a solenoid is determined by a limit mapping. Properties of periodic points of limit mappings are also studied.

INTRODUCTION

There are various ways one can view solenoids (see, e.g., [4], [11, (10.12)]). A solenoid may be defined as follows. Let $P = (p_1, p_2, \dots)$ be a sequence of prime numbers (1 not being included as a prime). The P -adic solenoid Σ_P is the inverse limit of the inverse sequence

$$(1) \quad \mathbb{S}^1 \xleftarrow{f_1^2} \mathbb{S}^1 \xleftarrow{f_2^3} \mathbb{S}^1 \xleftarrow{f_3^4} \dots,$$

where $\mathbb{S}^1$ is the unit circle considered as a subspace of the space $\mathbb{C}$ of all complex numbers endowed with the natural topology, and every bonding mapping f_n^{n+1} is given by $f_n^{n+1}(z) = z^{p_n}$ for each $z \in \mathbb{S}^1$, $n \in \mathbb{N}$. If $p_n = 2$ for all $n \in \mathbb{N}$, then the solenoid Σ_P is said to be *dyadic*. As is well known, the P -adic solenoid is a metric continuum which is not locally connected at any point. Sequences P and Q of prime numbers are said to be *equivalent* (written $P \sim Q$) if a finite number of terms can be deleted from each sequence so that every prime number occurs the same number of times in the deleted sequences. The solenoids Σ_P and Σ_Q are homeomorphic iff $P \sim Q$ [2], [14] (see also [1]). The solenoid is a compact abelian group under the coordinatewise multiplication with the identity $(1, 1, \dots)$. The condition $P \sim Q$ is also a criterion for the existence of a topological isomorphism between the topological groups Σ_P and Σ_Q [14].

For given $k \in \mathbb{N}$ let us consider the limit mapping $h_P^k : \Sigma_P \rightarrow \Sigma_P$ induced by the mapping $\{h_n^k : n \in \mathbb{N}\}$ between two copies of (1) :

$$(2) \quad \begin{array}{ccccccc} \mathbb{S}^1 & \xleftarrow{f_1^2} & \mathbb{S}^1 & \xleftarrow{f_2^3} & \mathbb{S}^1 & \xleftarrow{f_3^4} & \dots & \Sigma_P \\ h_1^k \downarrow & & \downarrow h_2^k & & \downarrow h_3^k & & & \downarrow h_P^k \\ \mathbb{S}^1 & \xleftarrow{f_1^2} & \mathbb{S}^1 & \xleftarrow{f_2^3} & \mathbb{S}^1 & \xleftarrow{f_3^4} & \dots & \Sigma_P, \end{array}$$

Received by the editors December 16, 2003 and, in revised form, April 19, 2004.

2000 *Mathematics Subject Classification.* Primary 54F15.

Key words and phrases. Covering mapping, limit mapping, periodic point, solenoid.

©2005 American Mathematical Society
 Reverts to public domain 28 years from publication

where h_n^k is the k -th potency mapping for each $n \in \mathbb{N}$, that is, $h_n^k(z) = z^k$ for every $z \in \mathbb{S}^1$.

This note deals with the limit mappings h_P^k . Note that, for each $k \in \mathbb{N}$, the limit mapping $h_{(2,2,\dots)}^k$ of the dyadic solenoid is a covering mapping [17], [3]. Theorem 19 of [3] states that the dyadic solenoid admits any odd degree k covering mapping of the form $h_{(2,2,\dots)}^k$, while it does not admit any covering mapping $h_{(2,2,\dots)}^k$ of an even degree k . For each $k \in \mathbb{N}$ the set of periodic points of $h_{(2,2,\dots)}^k$ is dense in $\Sigma_{(2,2,\dots)}$ [17], [3].

The purpose of this note is to extend the above results concerning the mappings $h_{(2,2,\dots)}^k$ to the limit mappings h_P^k of an arbitrary P -adic solenoid.

1. PRELIMINARIES

In this section we establish some notation that is used throughout. As usual, we denote by $\mathbb{N}$ the set of all positive integers. For any $n \in \mathbb{N}$ we use $\sqrt[n]{1}$ to denote the set of all values of the n -th root of 1.

It is well known that in studying finite-sheeted covering mappings from connected topological spaces onto solenoids there is no loss of generality in assuming that covering spaces are metric continua, i.e., metric compact connected spaces. So all topological spaces are supposed to be metric. A *mapping* between two spaces always means a continuous function. For basic notions of the theory of inverse limit spaces we refer the reader to [5, Chapter VIII] and [6, Chapter 2]. Let $\{X_n, \pi_n^{n+1}\}$ and $\{Y_n, \rho_n^{n+1}\}$ be inverse sequences of spaces X_n and Y_n with bonding mappings $\pi_n^{n+1} : X_{n+1} \rightarrow X_n$ and $\rho_n^{n+1} : Y_{n+1} \rightarrow Y_n$, where $n \in \mathbb{N}$. We denote by X_∞ and Y_∞ the inverse limit spaces of $\{X_n, \pi_n^{n+1}\}$ and $\{Y_n, \rho_n^{n+1}\}$ respectively. Recall that a sequence $\{\sigma_n : X_n \rightarrow Y_n \mid n \in \mathbb{N}\}$ of mappings is called a *mapping* between the inverse sequences $\{X_n, \pi_n^{n+1}\}$ and $\{Y_n, \rho_n^{n+1}\}$ if $\sigma_n \circ \pi_n^{n+1} = \rho_n^{n+1} \circ \sigma_{n+1}$ for all $n \in \mathbb{N}$. Then there exists a mapping $\sigma_\infty : X_\infty \rightarrow Y_\infty : (x_1, x_2, \dots) \mapsto (\sigma_1(x_1), \sigma_2(x_2), \dots)$. It is called the *limit mapping* induced by $\{\sigma_n : n \in \mathbb{N}\}$.

For any $m, n \in \mathbb{N}$ satisfying $m \leq n$, we denote by f_m^n the bonding mapping of the inverse sequence (1). Thus, for each $n \in \mathbb{N}$, the mapping f_n^n is the identity on $\mathbb{S}^1$ and $f_l^n = f_l^m \circ f_m^n$ for all $l, m, n \in \mathbb{N}$ satisfying $l \leq m \leq n$.

The identity $(1, 1, \dots)$ of a solenoid is denoted by e .

Recall that a surjective mapping $f : X \rightarrow Y$ between spaces X and Y is called:

- *k -to-1*, where $k \in \mathbb{N}$, provided that $\text{card } f^{-1}(y) = k$ for each $y \in Y$;
- a *finite-sheeted covering mapping* if it is a *k -sheeted* (*k -fold*) *covering mapping* for some $k \in \mathbb{N}$; that is, every point $y \in Y$ has an open neighborhood W in Y such that the inverse image $f^{-1}(W)$ can be written as the union of k disjoint open subsets of X each of which is mapped homeomorphically onto W under f .

The number k in the above definitions is called a *degree* of the mapping. Finite-sheeted covering mappings $f_1 : X_1 \rightarrow Y$ and $f_2 : X_2 \rightarrow Y$ are said to be *isomorphic* (or *equivalent*) if there is a homeomorphism $g : X_1 \rightarrow X_2$ such that $f_1 = f_2 \circ g$.

Throughout this note $P = (p_1, p_2, \dots)$ denotes a sequence of prime numbers. We say that a prime number p *occurs infinitely often* in P if $p = p_n$ for infinitely many terms p_n , $n \in \mathbb{N}$. We denote by $S(P)$ a subset of $\mathbb{N} \setminus \{1\}$ consisting of all prime numbers which do not occur infinitely often in P . In other words, $q \in S(P)$ iff $q > 1$ is a prime number such that, for some $m \in \mathbb{N}$, we have $q \neq p_n$ for all $n \geq m$.

2. COVERING MAPPINGS

The following proposition is an analog of [17, Lemma 1] and [3, Proposition 8].

Proposition 1. *For each $k \in \mathbb{N}$ the limit mapping $h_P^k : \Sigma_P \rightarrow \Sigma_P$ is a finite-sheeted covering mapping, and its degree is at most k .*

Proof. Let us fix $k \in \mathbb{N}$. Clearly, it suffices to show that h_P^k is an m -to-1 surjective open mapping for some $m \leq k$ (cf. [16, Chapter X, §6], [3, Proposition 1]).

Since $\mathbb{S}^1$ is compact and each mapping h_n^k in (2) is surjective, the limit mapping h_P^k is surjective too [6, Theorem 3.2.14]. Because h_P^k is a continuous homomorphism from a compact group onto itself, h_P^k is an open mapping [11, (5.29)] and the equality $\text{card}(h_P^k)^{-1}(y) = \text{card}(h_P^k)^{-1}(e)$ holds for each $y \in \Sigma_P$. Note, if $(x_1, x_2, \dots) \in (h_P^k)^{-1}(e)$, then $x_n \in \sqrt[k]{1}$ for all $n \in \mathbb{N}$. Using these observations, one can easily see that h_P^k is an m -to-1 mapping and $m \leq k$. $\square$

We shall determine degrees of the limit mappings h_P^k . In order to do this we find out cardinalities of fibers $(h_P^k)^{-1}(e)$, $k \in \mathbb{N}$. (See also [3, Statements 15, 17, 18]).

Proposition 2. *If k is a prime number which occurs infinitely often in the sequence P , then the limit mapping $h_P^k : \Sigma_P \rightarrow \Sigma_P$ is a homeomorphism.*

Proof. We claim that the set $(h_P^k)^{-1}(e)$ consists of only one point e . To show this we suppose that $z = (z_1, z_2, \dots) \in (h_P^k)^{-1}(e)$. Then $z_n^k = 1$ for all $n \in \mathbb{N}$. By assumption, for given $n \in \mathbb{N}$, there is an integer $m \geq n$ such that f_m^{m+1} in (1) is the k -th potency mapping. Therefore, we have

$$z_n = f_n^m(f_m^{m+1}(z_{m+1})) = f_n^m(z_{m+1}^k) = f_n^m(1) = 1.$$

Since these equalities hold for each $n \in \mathbb{N}$, it follows that $z = e$, as claimed.

Thus $\text{card}(h_P^k)^{-1}(e) = 1$ and, by Proposition 1, the limit mapping h_P^k is a homeomorphism. $\square$

The verification of the following lemma is straightforward (cf. [3, Fact 16]).

Lemma. *If $k = l \cdot m$ for some $l, m \in \mathbb{N}$, then $h_P^k = h_P^l \circ h_P^m$.*

Combining the Lemma and Proposition 2, we have:

Proposition 3. *If each prime divisor of $k \in \mathbb{N}$ occurs infinitely often in the sequence P , then the limit mapping $h_P^k : \Sigma_P \rightarrow \Sigma_P$ is a homeomorphism.*

Let $n \in \mathbb{N}$ and $\xi_n = \cos \frac{2\pi}{n} + i \sin \frac{2\pi}{n}$, where i is the imaginary unit, i.e., $i^2 = -1$. The set $\sqrt[n]{1}$ is a multiplicative cyclic group $\{1, \xi_n, \xi_n^2, \dots, \xi_n^{n-1}\}$ generated by ξ_n . For $m \in \mathbb{N}$, we define a homomorphism $\psi_m : \sqrt[n]{1} \rightarrow \sqrt[n]{1}$ by setting $\psi_m(\xi_n^j) = \xi_n^{jm}$, where $j \in \{1, 2, \dots, n\}$. Let n and m be relatively prime. In this case, one can easily see that ψ_m is injective. This implies that ψ_m is an automorphism, i.e., a bijective homomorphism from a group onto itself. Therefore, there exists an automorphism $\phi_m : \sqrt[n]{1} \rightarrow \sqrt[n]{1}$, which is the inverse of ψ_m . In other words, for each $a \in \sqrt[n]{1}$ we have exactly one element b of the group $\sqrt[n]{1}$ whose m -th power is equal to a . The value of the automorphism ϕ_m at a is just the element b .

Proposition 4. *If $k \in S(P)$, then the limit mapping $h_P^k : \Sigma_P \rightarrow \Sigma_P$ is a k -fold covering mapping.*

Proof. Denote by $m \in \mathbb{N}$ the least number l such that $k \neq p_n$ for all $n \geq l$.

If $m = 1$, i.e., k is not a term of the sequence P , then for each $p_n \in P$, the numbers p_n and k are relatively prime. Let ϕ_{p_n} be the inverse of the automorphism $\psi_{p_n} : \sqrt[k]{1} \rightarrow \sqrt[k]{1} : z \mapsto z^{p_n}$. We have $(\phi_{p_n}(z))^{p_n} = z$ for every $z \in \mathbb{S}^1$. It is easy to see that for each k -th root of unity ξ_k^j , where $j \in \{1, 2, \dots, k\}$, the element

$$(\xi_k^j, \phi_{p_1}(\xi_k^j), \phi_{p_2} \circ \phi_{p_1}(\xi_k^j), \dots)$$

lies in the set $(h_P^k)^{-1}(e)$. Clearly, in this way one obtains k distinct points of the fiber $(h_P^k)^{-1}(e)$. Hence $\text{card}((h_P^k)^{-1}(e)) \geq k$. On the other hand, by Proposition 1, $\text{card}((h_P^k)^{-1}(e)) \leq k$ and the desired conclusion follows.

If $m > 1$, then we consider the sequence $Q = (p_m, p_{m+1}, \dots)$ which is equivalent to P . As mentioned in the Introduction, there exists a topological isomorphism $\rho : \Sigma_P \rightarrow \Sigma_Q$ of topological groups (take ρ to be the shift of the form $\rho((x_1, x_2, \dots)) = (x_m, x_{m+1}, \dots)$, where $(x_1, x_2, \dots) \in \Sigma_P$). The inverse of ρ is denoted by τ .

One can readily verify that the following diagram

$$(3) \quad \begin{array}{ccc} \Sigma_P & \xrightarrow{\rho} & \Sigma_Q \\ h_P^k \downarrow & & \downarrow h_Q^k \\ \Sigma_P & \xrightarrow{\rho} & \Sigma_Q \end{array}$$

is commutative, i.e., $\rho \circ h_P^k = h_Q^k \circ \rho$. According to the first part of this proof, the limit mapping h_Q^k is a k -fold covering mapping.

It follows from the commutativity of the diagram (3) that

$$(h_P^k)^{-1}(e) = (h_P^k)^{-1}(\tau(e)) = \tau((h_Q^k)^{-1}(e)).$$

But the mapping τ is injective and $\text{card}((h_Q^k)^{-1}(e)) = k$. Thus the equality $\text{card}((h_P^k)^{-1}(e)) = k$ holds. Therefore, by Proposition 1, h_P^k is a k -fold covering mapping. $\square$

As an immediate consequence of the above results we have the following theorem.

Theorem 1. *Let $k \in \mathbb{N}$ be given. The P -adic solenoid admits a k -fold covering mapping of the form h_P^k if and only if k has no prime divisor which occurs infinitely often in the sequence P .*

It is interesting to restate Theorem 1. To do this we recall some facts. We refer to [11] for basic material on topological groups. The character group of Σ_P , consisting of all continuous homomorphisms from Σ_P into $\mathbb{S}^1$, is topologically isomorphic to the discrete group F_P of P -adic rationals (all rationals of the form $m/(p_1 p_2 \dots p_n)$, where m is an integer and $n \in \mathbb{N}$) [11, (25.3)]. It follows from the Pontrjagin duality that two solenoids Σ_P and Σ_Q are topologically isomorphic if and only if the additive groups F_P and F_Q are isomorphic.

For $n \in \mathbb{N}$, an additive abelian group G is said to be n -divisible provided that for each element $g \in G$ there is an element $g' \in G$ such that $ng' = g$. By number-theoretic considerations one can see that, for a prime number p , the group F_P is p -divisible if and only if p occurs infinitely often in P .

Theorem 1'. *Let $k \in \mathbb{N}$ be given. The P -adic solenoid admits a k -fold covering mapping of the form h_P^k if and only if k has no prime divisor q such that the group of P -adic rationals is q -divisible.*

Remark 1. Let $k \geq 2$ and let G be a compact connected abelian group. Suppose that the character group of G is k -divisible. Using Theorem 1 of [9] and [10], it can be shown that there is no k -sheeted covering mapping from a connected Hausdorff topological space onto G [10, Corollary 1].

In what follows, a *finite-sheeted connected covering of a solenoid* means a finite-sheeted covering mapping from a connected space onto a solenoid. We conclude this section with a theorem which summarizes the results concerning finite-sheeted connected coverings of solenoids. For each $k \in \mathbb{N}$ that is not divisible by any of the primes that occur infinitely often in the sequence P there is just one (up to equivalence) k -fold connected covering of Σ_P , and these are the only finite-sheeted connected coverings of Σ_P (see [7, Example 2], [8, p. 82, Theorem 3] and [15, Proposition 2.2]). This fact is a corollary of the theory of overlays (we refer to [7], [8] and [12] for this theory). According to the above results we have the following theorem (with the same dichotomy of positive integers k as in Theorems 1 and 1').

Theorem 2. *Let P be a sequence of prime numbers and $k \in \mathbb{N}$. If k is a multiple of some prime number which occurs infinitely often in P , then there is no k -fold connected covering of the P -adic solenoid Σ_P . Otherwise, the limit mapping h_P^k is a k -fold connected covering and, moreover, each k -fold connected covering of Σ_P is equivalent to h_P^k .*

Remark 2. The fact that each finite-sheeted connected covering of Σ_P is isomorphic to some limit mapping h_P^k (and, as a consequence, Theorem 2) can be proved without using the theory of overlays. In order to show this we make use of the approximate construction sketched in [9]. (A detailed account of this construction for a finite-sheeted covering mapping from a Hausdorff topological space onto an arbitrary compact connected group is contained in [10].) For the sake of completeness, we outline the proof of the above-mentioned fact as follows.

Let $f : X \rightarrow \Sigma_P$ be a k -fold connected covering. There exists an inverse sequence $\{X_n, g_n^{n+1}\}$ and a mapping $\{g_n : X_n \rightarrow S^1 \mid n \in \mathbb{N}\}$ between $\{X_n, g_n^{n+1}\}$ and the inverse sequence (1) such that the properties listed below are fulfilled: 1) for each $n \in \mathbb{N}$ the space X_n is connected and locally pathwise connected and $g_n : X_n \rightarrow S^1$ is a k -fold covering mapping; 2) the k -fold covering mapping f is isomorphic to the limit mapping $g_\infty : X_\infty \rightarrow \Sigma_P$ induced by $\{g_n : X_n \rightarrow S^1\}$; 3) for each $n \in \mathbb{N}$ there is a point $x_n \in X_n$ such that $g_n^{n+1}(x_{n+1}) = x_n$ and $g_n(x_n) = 1$. Using the classical covering space theory (see, e.g., [13, Chapter V, Corollary 6.4]), one constructs a sequence $\{\phi_n : S^1 \rightarrow X_n\}$ of homeomorphisms such that $\phi_n(1) = x_n$ and $g_n \circ \phi_n = h_n^k$, where h_n^k is the k -fold covering mapping from (2). In view of uniqueness of liftings [13, Chapter V, Lemma 3.2], it is easy to see that $\{\phi_n : n \in \mathbb{N}\}$ is a mapping between the inverse sequences (1) and $\{X_n, g_n^{n+1}\}$. Since each mapping $\phi_n : S^1 \rightarrow X_n$ is a homeomorphism, the limit mapping $\phi_\infty : \Sigma_P \rightarrow X_\infty$ induced by $\{\phi_n : n \in \mathbb{N}\}$ is a homeomorphism too [6, Proposition 2.5.10]. It is clear that $g_\infty \circ \phi_\infty = h_P^k$. In other words, the limit mappings g_∞ and h_P^k are isomorphic. Hence the covering mapping $f : X \rightarrow \Sigma_P$, being isomorphic to $g_\infty : X_\infty \rightarrow \Sigma_P$, is isomorphic to the limit mapping $h_P^k : \Sigma_P \rightarrow \Sigma_P$ as well.

3. PERIODIC POINTS

Let X be a space and let $f : X \rightarrow X$ be a mapping. For $n \in \mathbb{N}$, the composite $f \circ f \circ \dots \circ f$ (n times) is called the *n -th iteration* of f and is denoted by f^n ($f^1 = f$).

A point $x \in X$ is said to be *periodic* if there exists $n \in \mathbb{N}$ such that $f^n(x) = x$. In this case, n is called *the period* of x under f .

In this section we shall prove three propositions which correspond to the following cases: 1) $S(P) = \emptyset$; 2) $S(P)$ is an infinite set; 3) $S(P)$ is a nonempty finite set.

We first remark that for any sequence of prime numbers P the limit mapping h_P^1 is the identity on the P -adic solenoid Σ_P . Therefore, the set of periodic points of h_P^1 coincides with the whole space Σ_P .

Proposition 5. *Let $P = (p_1, p_2, \dots)$ be a sequence of prime numbers such that each prime number from $\mathbb{N}$ occurs infinitely often in P . Then, for each $k \geq 2$, the identity e of the P -adic solenoid is the only periodic point of the limit mapping h_P^k .*

Proof. Fix $k \geq 2$ and suppose that $z = (z_1, z_2, \dots)$ is a point of Σ_P such that $(h_P^k)^m(z) = h_P^{km}(z) = z$ for some $m \in \mathbb{N}$. Then, for each $n \in \mathbb{N}$, we have $z_n^{km-1} = 1$ with $km \geq 2$. If $km = 2$, then $z = e$. Let $km \geq 3$. For given $n \in \mathbb{N}$, we choose $l > n$ such that the product $p_n p_{n+1} \dots p_{l-1}$ of the terms of P is a multiple of $km - 1$. Then

$$(4) \quad z_n = f_n^l(z_l) = z_l^{p_n p_{n+1} \dots p_{l-1}} = 1.$$

Since this is valid for each $n \in \mathbb{N}$, we have $z = e$, as required. $\square$

The next proposition is based on [3, Proposition 40] (cf. [17, Proposition 9]). Note that, in the case of the dyadic solenoid, the set $S((2, 2, \dots))$ is infinite. Before coming to Proposition 6 we recall Euler's Theorem. It states that for relatively prime positive integers a and m the number $a^{\varphi(m)} - 1$ is divisible by m . Here, $\varphi(m)$ is a value of the Euler function at m . That is, $\varphi(1) = 1$ and, for $m > 1$, $\varphi(m) = q_1^{l_1-1} q_2^{l_2-1} \dots q_n^{l_n-1} (q_1 - 1)(q_2 - 1) \dots (q_n - 1)$, where $m = q_1^{l_1} q_2^{l_2} \dots q_n^{l_n}$ is the canonical factorization of m , i.e., $q_1, q_2, \dots, q_n \geq 2$ are distinct prime numbers and $l_1, l_2, \dots, l_n \in \mathbb{N}$.

Proposition 6. *Let $P = (p_1, p_2, \dots)$ be a sequence of prime numbers such that the set $S(P)$ is infinite. Then, for each $k \in \mathbb{N}$, the set of all periodic points of the limit mapping h_P^k is dense in Σ_P .*

Proof. Given $k > 1$, we choose $q \in S(P)$ such that $q > k$. Let $\pi_n : \Sigma_P \rightarrow \mathbb{S}^1 : (z_1, z_2, \dots) \mapsto z_n$ be the n -th projection mapping of the inverse sequence (1), $n \in \mathbb{N}$.

Take a basic open subset U of Σ_P . That is, $U = \pi_l^{-1}(V_l)$ for some $l \in \mathbb{N}$ and some open subset V_l of $\mathbb{S}^1$. Recall that for any $n \in \mathbb{N}$ such that $n \geq l$, the projection mappings π_n and π_l satisfy the equality $\pi_l = f_l^n \circ \pi_n$. Therefore, for each $n \geq l$, we have $U = \pi_l^{-1}(V_l) = \pi_n^{-1}(V_n)$, where $V_n = (f_l^n)^{-1}(V_l)$ is an open subset of $\mathbb{S}^1$. Since $q \in S(P)$ we can choose $n \geq l$ such that $p_j \neq q$ for every $j \geq n$. Fix such n .

Take $m \in \mathbb{N}$ such that the following condition is fulfilled:

$$(5) \quad \text{there exists a point } z_n \in V_n \text{ such that } z_n^{q^m} = 1.$$

For each $j \geq n$ the numbers p_j and q^m are relatively prime. Consider the automorphism $\psi_{p_j} : \sqrt[q^m]{1} \rightarrow \sqrt[q^m]{1} : z \mapsto z^{p_j}$ and its inverse ϕ_{p_j} . Thus we have

$$(6) \quad (\phi_{p_j}(z))^{p_j} = z \quad \text{for all } z \in \sqrt[q^m]{1}.$$

Let $\zeta = (z_n^{p_1 p_2 \dots p_{n-1}}, \dots, z_n^{p_{n-1}}, z_n, \phi_{p_n}(z_n), \phi_{p_{n+1}} \circ \phi_{p_n}(z_n), \dots)$. According to (6), the point ζ belongs to Σ_P . Moreover, by (5),

$$(7) \quad \zeta \in U.$$

We claim that ζ is a periodic point of h_P^k . Indeed, since each term of the sequence ζ is a q^m -th root of 1, we get

$$(8) \quad \zeta^{q^m} = e.$$

Since k and q^m are relatively prime, by Euler's Theorem, $k^{q^{m-1}(q-1)} - 1$ is divisible by q^m . Hence, by (8),

$$\zeta^{k^{q^{m-1}(q-1)}-1} = e \quad \text{and} \quad (h_P^k)^{q^{m-1}(q-1)}(\zeta) = \zeta^{k^{q^{m-1}(q-1)}} = \zeta.$$

In other words, ζ is a point of period $q^{m-1}(q-1)$ under h_P^k .

In view of (7) and since U was chosen as an arbitrary basic open subset of Σ_P , the proof is complete. $\square$

Proposition 7. *Let $P = (p_1, p_2, \dots)$ be a sequence of prime numbers such that the set $S(P)$ is nonempty and finite. Let $k \in \mathbb{N}$ be given. If k is a multiple of the product of all prime numbers from $S(P)$, then the identity e of the P -adic solenoid is the only periodic point of the limit mapping h_P^k . If there exists a prime number from $S(P)$ which is not a divisor of k , then the set of all periodic points of the limit mapping h_P^k is dense in Σ_P .*

Proof. Suppose that $S(P) = \{q_1, \dots, q_t\}$, where $t \in \mathbb{N}$.

First, let k be a multiple of $q_1 \cdot \dots \cdot q_t$, and let $z = (z_1, z_2, \dots) \in \Sigma_P$ be a periodic point of period $m \in \mathbb{N}$ under h_P^k . Thus, for all $n \in \mathbb{N}$, we have $z_n^{k^m-1} = 1$, where $k^m \geq 2$. If $k^m = 2$, then $z = e$. If $k^m \geq 3$, then we choose $n \in \mathbb{N}$ such that

$$(9) \quad p_j \notin S(P) \quad \text{for all } j \geq n.$$

Obviously, for each $q_j \in S(P)$, $j \in \{1, \dots, t\}$, the number $k^m - 1$ is not divisible by q_j . Hence, each prime divisor of $k^m - 1$ occurs infinitely often in the sequence P . Choose $l > n$ such that the number $p_n p_{n+1} \dots p_{l-1}$ is a multiple of $k^m - 1$. Then we have $z_n = 1$ (see (4)). This implies $z_l = f_l^n(z_n) = 1$ for every $l \leq n$. Since this is true for each n satisfying (9), we get $z = e$.

Second, we assume that $k > 1$ is not divisible by some $q \in S(P)$. Then every power of q and k are relatively prime. Repeating the arguments from the proof of Proposition 6 for these k and q , one can see that the set of periodic points of h_P^k is dense in Σ_P . $\square$

The author is much obliged to Professor S. A. Bogatyĭ who called the author's attention to the papers [3] and [17] in connection with the theorems of [9]. The author would like to thank Professor S. A. Grigorian for helpful conversations about the results in this note.

REFERENCES

1. J. M. Aarts and R. J. Fokkink, *The classification of solenoids*, Proc. Amer. Math. Soc. **111** (1991), 1161 – 1163. MR1042260 (91g:54046)
2. R. H. Bing, *A simple closed curve is the only homogeneous bounded plane continuum that contains an arc*, Canad. J. Math. **12** (1960), 209 – 230. MR0111001 (22:1869)
3. J. J. Charatonik and P. P. Covarrubias, *On covering mappings on solenoids*, Proc. Amer. Math. Soc. **130** (2002), 2145–2154. MR1896052 (2003c:54065)
4. D. van Dantzig, *Ueber topologisch homogene Kontinua*, Fund. Math. **15** (1930), 102 – 125.
5. S. Eilenberg and N. Steenrod, *Foundations of algebraic topology*, Princeton Univ. Press, Princeton, N. J., 1952. MR0050886 (14:398b)
6. R. Engelking, *General topology*, Monografie Matematyczne, Vol. 60, Polish Scientific Publishers, Warszawa, 1977. MR0500780 (58:18316b)

7. R. H. Fox, *On shape*, Fund. Math. **74** (1972), 47 – 71. MR0296914 (45:5973)
8. R. H. Fox, *Shape theory and covering spaces*, Lecture Notes in Math., Vol. 375, 71–90, Topology Conference Virginia Polytechnic Institute, 1973 (R.F. Dickman Jr., P. Fletcher, eds.), Springer, Berlin — Heidelberg — New York, 1974. MR0368007 (51:4249)
9. S. A. Grigorian and R. N. Gumerov, *On a covering group theorem and its applications*, Lobachevskii J. Math. **X** (2002), 9–16. MR1903663 (2003c:22011)
10. S. A. Grigorian and R. N. Gumerov, *On the structure of finite-sheeted coverings of compact connected groups*, (manuscript), accessible from: <http://lanl.arxiv.org/abs/math.GN/0403329>.
11. E. Hewitt and K. A. Ross, *Abstract harmonic analysis, Vol. I*, Springer-Verlag, Berlin, 1963. MR0551496 (81k:43001)
12. S. Mardešić and V. Matijević, *Classifying overlay structures of topological spaces*, Topology Appl. **113** (2001), 167 – 209. MR1821853 (2002d:55012)
13. W. S. Massey, *Algebraic topology: An Introduction*, Harcourt Brace Jovanovich, New York, 1967. MR0211390 (35:2271)
14. M. C. McCord, *Inverse limit sequences with covering maps*, Trans. Amer. Math. Soc. **114** (1965), 197 – 209. MR0173237 (30:3450)
15. T. T. Moore, *On Fox's theory of overlays*, Fund. Math. **99** (1978), 205 – 211. MR0482645 (58:2703)
16. G. T. Whyburn, *Analytic topology*, Amer. Math. Soc. Colloq. Publ. 28, Providence, RI, 1942. MR0007095 (4:86b)
17. Zhou Youcheng, *Covering mapping on solenoids and their dynamical properties*, Chinese Sci. Bull. **45** (2000), 1066–1070. MR1777211 (2002b:37022)

DEPARTMENT OF MECHANICS AND MATHEMATICS, KAZAN STATE UNIVERSITY, KREMLEVSKAYA
 18, KAZAN, 420008, TATARSTAN, RUSSIAN FEDERATION
E-mail address: `renat.gumerov@ksu.ru`