DISCRETE SPECTRA OF C^* -ALGEBRAS AND ORTHOGONALLY CLOSED SUBMODULES IN HILBERT C^* -MODULES

MASAHARU KUSUDA

(Communicated by David R. Larson)

ABSTRACT. Let A and B be C^* -algebras and let X be an A - B -imprimitivity bimodule. Then it is shown that if the spectrum $\widehat{A}$ of A (resp. $\widehat{B}$ of B) is discrete, then every closed A - B -submodule of X is orthogonally closed in X , and conversely that if $\widehat{A}$ (resp. $\widehat{B}$) is a T_1 -space and if every closed A - B -submodule of X is orthogonally closed in X , then $\widehat{A}$ (resp. $\widehat{B}$) is discrete.

1. INTRODUCTION

Let A be a C^* -algebra and let X be a Hilbert A -module with an A -valued inner product $\langle \cdot, \cdot \rangle$. For any closed Hilbert A -submodule Y of X , we denote by $Y^\perp$ the orthogonally complemented subspace of Y in X , i.e.,

$$Y^\perp = \{ x \in X \mid \langle x, y \rangle = 0 \text{ for all } y \in Y \}.$$

We say that a closed A -submodule Y of a Hilbert A -module X is *orthogonally complemented* in X if X coincides with $Y \oplus Y^\perp$, and that a closed A -submodule Y of a Hilbert A -module X is *orthogonally closed* in X if $Y^{\perp\perp} = Y$. If Y is orthogonally complemented in X , then it is orthogonally closed in X . But the converse is not necessarily true in general. Suppose that X is a full (right) Hilbert A -module. Then Schweitzer [7, Theorem 1] has shown that every closed right A -submodule of X is orthogonally closed if and only if every closed right A -submodule of X is orthogonally complemented in X . Here we remark that every full right Hilbert A -module is a $\mathcal{K}(X)$ - A -imprimitivity bimodule, where $\mathcal{K}(X)$ is the C^* -algebra generated by those operators $\theta_{x,y}$ with all $x, y \in X$ defined by $\theta_{x,y}(z) = x \cdot \langle y, z \rangle$ for all $z \in X$.

Let A and B be C^* -algebras and let X be an A - B -imprimitivity bimodule. Note that A is isomorphic to $\mathcal{K}(X)$. In this paper, we show that every closed A - B -submodule of X is orthogonally closed if and only if every closed A - B -submodule of X is orthogonally complemented in X . As a corollary, we show that if the spectrum $\widehat{A}$ of A (resp. $\widehat{B}$ of B) is discrete, then every closed A - B -submodule of X is orthogonally closed in X , and conversely that if $\widehat{A}$ (resp. $\widehat{B}$) is a T_1 -space and if every closed A - B -submodule of X is orthogonally closed in X , then $\widehat{A}$ (resp. $\widehat{B}$) is discrete.

Received by the editors December 3, 2003 and, in revised form, June 23, 2004.

2000 *Mathematics Subject Classification.* Primary 46L05, 46L08.

Key words and phrases. Hilbert C^* -modules, orthogonally closed.

©2005 American Mathematical Society
 Reverts to public domain 28 years from publication

2. RESULTS

Let A and B be C^* -algebras and let X be an A - B -imprimitivity bimodule (see [4] or [6] for the definition of an imprimitivity bimodule). We denote by ${}_A\langle \cdot, \cdot \rangle$ the A -valued inner product on X as a left Hilbert A -module and by $\langle \cdot, \cdot \rangle_B$ the B -valued inner product on X as a right Hilbert B -module, respectively.

Throughout this paper, by an A - B -submodule we mean a left A -submodule and right B -submodule.

Two C^* -algebras A and B are said to be *Morita equivalent* if there exists an A - B -imprimitivity bimodule. We remark that in this paper, Morita equivalence means strong Morita equivalence in the sense of Rieffel (cf. [6, Remark 3.15]).

Let A be a C^* -algebra and let I be a closed ideal of A . Throughout this paper, unless otherwise stated, by an ideal of A we always mean a two-sided ideal. Then $I^\perp$ is defined by

$$I^\perp = \{a \in A \mid ax = 0 \text{ for all } x \in I\}.$$

It is easily seen that $I^\perp$ is a closed ideal of A . Note here that A can be regarded as an A - A -imprimitivity bimodule, for the bimodule structure given by the multiplication in A , with ${}_A\langle a, b \rangle = ab^*$ and $\langle a, b \rangle_A = a^*b$ for $a, b \in A$ (see [6, Example 3.5]). Then I is a closed A - A -submodule of A .

Let A and B be C^* -algebras and let X be an A - B -imprimitivity bimodule. For a closed A - B -submodule Y of X , its orthogonally complemented subspace $Y^\perp$ in X is defined by

$$Y^\perp = \{x \in X \mid {}_A\langle x, y \rangle = \langle x, y \rangle_B = 0 \text{ for all } y \in Y\}.$$

Lemma 2.1. *Let A be a C^* -algebra and let I be any closed ideal I of A . If every closed ideal of A is orthogonally closed in A , then we have $A = I \oplus I^\perp$.*

Proof. Using the fact that for an ideal J of A we have $J \cap J^\perp = \{0\}$, it follows that

$$(I \oplus I^\perp)^\perp = I^\perp \cap (I^\perp)^\perp = \{0\}.$$

Hence, using the hypothesis, we have $I \oplus I^\perp = (I \oplus I^\perp)^{\perp\perp} = A$. $\square$

Now we recall the Rieffel correspondence (see [6, Theorem 3.22]). Let two C^* -algebras A and B be Morita equivalent and let X be an A - B -imprimitivity bimodule. We denote by $\mathcal{I}(A)$ (resp. $\mathcal{I}(B)$) the set of all closed (two-sided) ideals of A (resp. B), and by $\mathcal{S}(X)$ the set of closed A - B -submodules of X . Note that $\mathcal{I}(A)$, $\mathcal{I}(B)$ and $\mathcal{S}(X)$ can be partially ordered by inclusion, and are then lattices. Then there are natural lattice isomorphisms among $\mathcal{I}(A)$, $\mathcal{I}(B)$ and $\mathcal{S}(X)$ given by

$$\mathcal{I}(A) \ni I \longmapsto {}_IX \in \mathcal{S}(X), \text{ where } {}_IX = \{y \in X \mid {}_A\langle y, x \rangle \in I \text{ for all } x \in X\};$$

$$\mathcal{S}(X) \ni Y \longmapsto I_Y \in \mathcal{I}(A), \quad {}_YI \in \mathcal{I}(B),$$

where I_Y is the closed linear span of $\{{}_A\langle y, x \rangle \mid y \in Y \text{ and } x \in X\}$

and ${}_YI$ is the closed linear span of $\{\langle x, y \rangle_B \mid y \in Y \text{ and } x \in X\}$;

$$\mathcal{I}(B) \ni J \longmapsto X_J \in \mathcal{S}(X), \text{ where } X_J = \{y \in X \mid \langle x, y \rangle_B \in J \text{ for all } x \in X\}.$$

We refer to such lattice isomorphisms among $\mathcal{I}(A)$, $\mathcal{I}(B)$ and $\mathcal{S}(X)$ as the *Rieffel correspondences* (see [6, 3.3]). Note that ${}_IX$ is the closed linear span of $I \cdot X$ (cf. [6, Lemma 3.23]), and we will employ this fact in the proof of Lemma 2.2.

Lemma 2.2. *Let A and B be C^* -algebras and let X be an A - B -imprimitivity bimodule. For any closed ideal I of A , we have $({}_IX)^\perp = {}_{I^\perp}X$.*

Proof. Since ${}_IX$ is the closed linear span of IX , we see from ${}_A\langle I^\perp X, IX \rangle = I^\perp \cdot {}_A\langle X, X \rangle \cdot I = \{0\}$ that ${}_I^\perp X \subset ({}_IX)^\perp$. For the reverse inclusion, let $Y = (IX)^\perp$ and note that $\{0\} = {}_A\langle Y, IX \rangle = {}_A\langle Y, X \rangle \cdot I$ implies that ${}_A\langle Y, Y \rangle$ is contained in $I^\perp$, hence $Y \subset I^\perp X$ by a well-known argument. $\square$

Now we are in a position to establish the main result.

Theorem 2.3. *Let A and B be C^* -algebras and let X be an A - B -imprimitivity bimodule. Then every closed A - B -submodule of X is orthogonally closed in X if and only if every closed A - B -submodule of X is orthogonally complemented in X .*

Proof. Suppose that every closed A - B -submodule of X is orthogonally closed in X . Let I be any closed ideal of A . We claim that $A = I \oplus I^\perp$. Since it follows from Lemma 2.2 that $({}_IX)^\perp = {}_{I^\perp}X$, we obtain $({}_IX)^{\perp\perp} = ({}_{I^\perp}X)^\perp$, and with I replaced by $I^\perp$ we have $({}_{I^\perp}X)^\perp = {}_{I^{\perp\perp}}X$. Since ${}_IX$ is a closed A - B -submodule, by assumption we see that ${}_{I^{\perp\perp}}X = ({}_{I^\perp}X)^\perp = ({}_IX)^{\perp\perp} = {}_IX$. By the Rieffel correspondence, we then see that $I^{\perp\perp} = I$. Thus we see that every closed ideal of A is orthogonally closed in A . Hence Lemma 2.1 yields that $A = I \oplus I^\perp$ for any closed ideal I .

Let Y be any closed A - B -submodule of X . Since I_Y is a closed ideal of A , it follows from the above claim that $A = I_Y \oplus (I_Y)^\perp$. We remark that $I_{Y^\perp} = (I_Y)^\perp$ (see the proof of [3, Theorem 2.3] for the details). Since the Rieffel correspondences are natural lattice isomorphisms among $\mathcal{I}(A)$, $\mathcal{I}(B)$ and $\mathcal{S}(X)$, a least upper bound $Y \vee Y^\perp (= Y \oplus Y^\perp)$ of Y and $Y^\perp$ corresponds to a least upper bound $I_Y \vee I_{Y^\perp} (= I_Y \vee (I_Y)^\perp = I_Y \oplus (I_Y)^\perp = A)$ of I_Y and $I_{Y^\perp}$. Since A and X correspond by the Rieffel correspondence, we conclude that $X = Y \oplus Y^\perp$. $\square$

We denote by $\widehat{A}$ the spectrum of A , that is, the set of (unitary) equivalence classes of nonzero irreducible representations of A equipped with the Jacobson topology. We note that $\widehat{A}$ is a locally compact space, not necessarily a T_0 -space. The reader is referred to [1] for the spectrum of a C^* -algebra.

Corollary 2.4. *Let two C^* -algebras A and B be Morita equivalent and let X be an A - B -imprimitivity bimodule. Consider the following conditions:*

- (1) *The spectrum $\widehat{A}$ of A is discrete in the Jacobson topology.*
- (2) *The spectrum $\widehat{B}$ of B is discrete in the Jacobson topology.*
- (3) *Every closed A - B -submodule of X is complemented in X .*
- (4) *Every closed A - B -submodule of X is orthogonally closed in X .*

Then we have (1) $\iff$ (2) $\implies$ (3) $\iff$ (4). If either $\widehat{A}$ or $\widehat{B}$ is a T_1 -space, then conditions (1) – (4) are equivalent.

Proof. This easily follows from [3, Theorem 2.3] and Theorem 2.3 above. $\square$

Note that in the implication (4) $\implies$ (1) above, the assumption that either $\widehat{A}$ or $\widehat{B}$ be a T_1 -space is necessary (see [3, Remark 2.4 (2)]). We end this paper by stating a remark.

Remark 2.5. Let A and B be C^* -algebras and let X be an A - B -imprimitivity bimodule. Consider the following conditions:

- (1) *Every closed right B -submodule of X is orthogonally closed in X .*
- (2) *Every closed right B -submodule of X is orthogonally complemented in X .*
- (3) *Every closed A - B -submodule of X is orthogonally closed in X .*

(4) Every closed A - B -submodule of X is orthogonally complemented in X .

Then the equivalence of (1) and (2) follows from [7], the implication (2) $\implies$ (4) is obvious, and the equivalence of (3) and (4) is nothing but Theorem 2.3. When B is simple, it follows from an easy application of the Rieffel correspondence that condition (4) is true. In this case, if B is not an elementary C^* -algebra, both conditions (1) and (2) are false (see [5] and [7]), that is, condition (4) does not necessarily imply condition (2). Thus we see that (1) $\iff$ (2) $\not\Rightarrow$ (3) $\iff$ (4).

ACKNOWLEDGEMENT

This research was financially supported by the Kansai University Research Grants: Grant-in Aid for Encouragement for Scientists 2003.

REFERENCES

1. J. Dixmier, *C^* -algebras*, North Holland, New York, 1982. MR0458185 (56:16388)
2. M. Kusuda, *Three-space problems in discrete spectra of C^* -algebras and dual C^* -algebras*, Proc. Royal Soc. Edinburgh **131A** (2001), 701–707. MR1838507 (2002m:46087)
3. M. Kusuda, *Discrete spectra of C^* -algebras and complemented submodules in Hilbert C^* -modules*, Proc. Amer. Math. Soc. **131** (2003), 3075–3081. MR1993216 (2004d:46067)
4. E. C. Lance, *Hilbert C^* -modules*, London Math. Soc. Lecture Note Series Vol. 210, Cambridge Univ. Press, Cambridge, 1994. MR1325694 (96k:46100)
5. B. Magajna, *Hilbert C^* -modules in which all closed submodules are complemented*, Proc. Amer. Math. Soc. **125** (1997), 849–852. MR1346981 (97e:46079)
6. I. Raeburn and D. P. Williams, *Morita equivalence and continuous-trace C^* -algebras*, Math. Surveys Monographs Vol. 60, Amer. Math. Soc., Providence, 1998. MR1634408 (2000c:46108)
7. J. Schweitzer, *A description of Hilbert C^* -modules in which all closed submodules are orthogonally closed*, Proc. Amer. Math. Soc. **127** (1999), 2123–2125. MR1646207 (99j:46068)

DEPARTMENT OF MATHEMATICS, FACULTY OF ENGINEERING, KANSAI UNIVERSITY, YAMATE-CHO 3-3-35, SUITA, OSAKA 564-8680, JAPAN

E-mail address: kusuda@ipcku.kansai-u.ac.jp