ADDENDUM TO “DENSE SUBSETS OF THE BOUNDARY OF A COXETER SYSTEM”

TETSUYA HOSAKA

(Communicated by Alexander N. Dranishnikov)

ABSTRACT. In this paper, we investigate boundaries of parabolic subgroups of Coxeter groups. Let (W, S) be a Coxeter system and let T be a subset of S such that the parabolic subgroup W_T is infinite. Then we show that if a certain set is quasi-dense in W , then $W\partial\Sigma(W_T, T)$ is dense in the boundary $\partial\Sigma(W, S)$ of the Coxeter system (W, S) , where $\partial\Sigma(W_T, T)$ is the boundary of (W_T, T) .

1. INTRODUCTION AND PRELIMINARIES

The purpose of this paper is to study boundaries of parabolic subgroups of Coxeter groups. In this paper, we use the same notation as [5] and [6]. Every Coxeter system (W, S) determines a *Davis-Moussong complex* $\Sigma(W, S)$ which is a CAT(0) geodesic space ([2], [3], [4], [7]). If W is infinite, then $\Sigma(W, S)$ is noncompact and $\Sigma(W, S)$ can be compactified by adding its ideal boundary $\partial\Sigma(W, S)$ ([1], [3, §4]). For each subset $T \subset S$, we consider the parabolic subgroup W_T generated by T . Then $\Sigma(W_T, T)$ is a subcomplex of $\Sigma(W, S)$ and the boundary $\partial\Sigma(W_T, T)$ of (W_T, T) is a subspace of $\partial\Sigma(W, S)$.

The purpose of this paper is to prove the following theorem.

Theorem 1.1. *Let (W, S) be a Coxeter system and let T be a subset of S such that W_T is infinite. If the set*

$$\bigcup \{W^{\{s\}} \mid s \in S \text{ such that } o(ss_0) = \infty \text{ and } s_0t \neq ts_0 \\ \text{for some } s_0 \in S \setminus T \text{ and } t \in \tilde{T}\}$$

is quasi-dense in W with respect to the word metric, then $W\partial\Sigma(W_T, T)$ is dense in $\partial\Sigma(W, S)$, where $W_{\tilde{T}}$ is the essential parabolic subgroup of (W_T, T) .

Remark. For a Gromov hyperbolic group G and the boundary ∂G of G , we can show that $G\alpha$ is dense in ∂G for any $\alpha \in \partial G$ by an easy argument. Hence if W is a hyperbolic Coxeter group, then $W\partial\Sigma(W_T, T)$ is dense in $\partial\Sigma(W, S)$ for any $T \subset S$ such that W_T is infinite.

Received by the editors July 5, 2004 and, in revised form, September 12, 2004 and October 5, 2004.

2000 *Mathematics Subject Classification.* Primary 57M07, 20F65, 20F55.

Key words and phrases. Boundaries of Coxeter groups.

The author was partly supported by the Grant-in-Aid for Scientific Research, The Ministry of Education, Culture, Sports, Science and Technology, Japan (No. 15740029).

©2005 American Mathematical Society
 Reverts to public domain 28 years from publication

As an application of Theorem 1.1, we obtain the following corollary.

Corollary 1.2. *Let (W, S) be a Coxeter system and let T be a subset of S such that W_T is infinite. Suppose that there exist a maximal spherical subset U of S and an element $s \in S$ such that $o(su) \geq 3$ for every $u \in U$ and $o(su_0) = \infty$ for some $u_0 \in U$. If*

- (1) $s \notin T$ and $u_0 \in \tilde{T}$, or
- (2) $u_0 \notin T$ and $s \in \tilde{T}$,

then $W\partial\Sigma(W_T, T)$ is dense in $\partial\Sigma(W, S)$.

Here the following problem is open.

Problem. Let (W, S) be a Coxeter system and let T be a subset of S such that W_T is infinite. Is it the case that if $\partial\Sigma(W_T, T)$ is not W -invariant, then $W\partial\Sigma(W_T, T)$ is dense in $\partial\Sigma(W, S)$? Particularly, is it the case that if (W, S) is an irreducible Coxeter system, then $W\partial\Sigma(W_T, T)$ is dense in $\partial\Sigma(W, S)$ for any subset T of S such that W_T is infinite?

2. PROOF OF THE MAIN RESULTS

Using some results in [5] and [6], we first prove the following lemma.

Lemma 2.1. *Let (W, S) be a Coxeter system, let T be a proper subset of S such that W_T is infinite, and let*

$$U = \{s \in S \setminus T \mid W^{\{s\}}s \cap W_T \text{ is finite}\}.$$

Then $W_{\tilde{T} \cup U} = W_{\tilde{T}} \times W_U$.

Proof. We note that $S(w) \subset T$ for $w \in W_T$. Let $u_0 \in U$ and let $T(u_0) = \{t \in T \mid tu_0 \neq u_0t\}$. We first show that $W_{T \setminus T(u_0)}$ is a subgroup of finite index in W_T . Here we note that $[W_T : W_{T \setminus T(u_0)}] = |A_{T \setminus T(u_0)} \cap W_T|$ by [5, Lemma 2.4]. Then

$$\begin{aligned} \bigcup_{T' \subset T(u_0)} (W_T)^{T'} &= \{w \in W_T \mid S(w) \subset T(u_0)\} \\ &= \{w \in W_T \mid T \setminus T(u_0) \subset T \setminus S(w)\} \\ &= A_{T \setminus T(u_0)} \cap W_T. \end{aligned}$$

We show that $(W_T)^{T'}$ is finite for any $T' \subset T(u_0)$. Let $T' \subset T(u_0)$. Since $tu_0 \neq u_0t$ for any $t \in T'$, $(W_T)^{T'}u_0 \subset W^{\{u_0\}}$ by [5, Lemma 2.7]. Hence $(W_T)^{T'} \subset W^{\{u_0\}}u_0 \cap W_T$, which is finite because $u_0 \in U$. Thus $(W_T)^{T'}$ is finite for any $T' \subset T(u_0)$, and $[W_T : W_{T \setminus T(u_0)}] = |A_{T \setminus T(u_0)} \cap W_T|$ is finite. By [5, Corollary 3.4], $\tilde{T} \subset T \setminus T(u_0)$. Hence $T(u_0) \subset T \setminus \tilde{T}$ for any $u_0 \in U$. Let $A = \{t \in T \mid tu_0 \neq u_0t \text{ for some } u_0 \in U\}$. Then $A = \bigcup_{u_0 \in U} T(u_0) \subset T \setminus \tilde{T}$ and

$$\tilde{T} \subset T \setminus A = \{t \in T \mid tu = ut \text{ for every } u \in U\}.$$

Thus $tu = ut$ for any $t \in \tilde{T}$ and $u \in U$. This means that $W_{\tilde{T} \cup U} = W_{\tilde{T}} \times W_U$. $\square$

Using the above lemma, we prove the main results.

Proof of Theorem 1.1. Suppose that

$$\begin{aligned} A := \bigcup \{W^{\{s\}} \mid s \in S \text{ such that } o(ss_0) = \infty \text{ and } s_0t \neq ts_0 \\ \text{for some } s_0 \in S \setminus T \text{ and } t \in \tilde{T}\} \end{aligned}$$

is quasi-dense in W .

We first show that for each $w \in A$, there exists $v \in W$ and $\alpha \in \partial\Sigma(W_T, T)$ such that $d(w, \text{Im } \xi_{v\alpha}) \leq N$, where N is the diameter of $K(W, S)$ in $\Sigma(W, S)$ and $\xi_{v\alpha}$ is the geodesic ray issuing from 1 such that $\xi_{v\alpha}(\infty) = v\alpha$.

Let $w \in A$. Then $w \in W^{\{s\}}$, $o(ss_0) = \infty$ and $s_0t \neq ts_0$ for some $s \in S$, $s_0 \in S \setminus T$ and $t \in \tilde{T}$. By Lemma 2.1, $W^{\{s_0\}}s_0 \cap W_T$ is infinite. Hence there exists a sequence $\{x_i\} \subset (W^{\{s_0\}}s_0 \cap W_T)^{-1}$ which converges to some point $\alpha \in \partial\Sigma(W_T, T)$. Since $x_i \in (W^{\{s_0\}}s_0)^{-1}$, $(s_0x_i)^{-1} = x_i^{-1}s_0 \in W^{\{s_0\}}$. By [6, Lemma 3.3], $d(w, \text{Im } \xi_{ws_0x_i}) \leq N$ for any i because $w \in W^{\{s\}}$, $s_0x_i \in (W^{\{s_0\}})^{-1}$ and $o(ss_0) = \infty$. Hence $d(w, \text{Im } \xi_{ws_0\alpha}) \leq N$.

For each $\beta \in \partial\Sigma(W, S)$, there exists a sequence $\{w_i\} \subset A$ which converges to β , because A is quasi-dense in W . By the above argument, there exist sequences $\{v_i\} \subset W$ and $\{\alpha_i\} \subset \partial\Sigma(W_T, T)$ such that $d(w_i, \text{Im } \xi_{v_i\alpha_i}) \leq N$ for each i . Then the sequence $\{v_i\alpha_i\}$ converges to β in $\partial\Sigma(W, S)$ because $\{w_i\}$ converges to β . Therefore $W\partial\Sigma(W_T, T)$ is dense in $\partial\Sigma(W, S)$. $\square$

Proof of Corollary 1.2. Suppose that there exist a maximal spherical subset U of S and an element $s \in S$ such that $o(su) \geq 3$ for any $u \in U$ and $o(su_0) = \infty$ for some $u_0 \in U$. Then $W^{\{s\}}$ is quasi-dense in W by [6, Lemma 2.5].

(1) If $s \notin T$ and $u_0 \in \tilde{T}$, then $W^{\{u_0\}}$ is quasi-dense in W because $W^{\{s\}}u_0 \subset W^{\{u_0\}}$ by [6, Lemma 2.4]. Hence $W\partial\Sigma(W_T, T)$ is dense in $\partial\Sigma(W, S)$ by Theorem 1.1.

(2) If $u_0 \notin T$ and $s \in \tilde{T}$, then by Theorem 1.1, $W\partial\Sigma(W_T, T)$ is dense in $\partial\Sigma(W, S)$, because $o(su_0) = \infty$, $u_0 \in S \setminus T$ and $s \in \tilde{T}$. $\square$

ACKNOWLEDGMENTS

The author would like to thank the referee for helpful advice.

REFERENCES

- [1] M. R. Bridson and A. Haefliger, *Metric spaces of non-positive curvature*, Springer-Verlag, Berlin, 1999. MR1744486 (2000k:53038)
- [2] M. W. Davis, *Groups generated by reflections and aspherical manifolds not covered by Euclidean space*, Ann. of Math. **117** (1983), 293–324. MR0690848 (86d:57025)
- [3] ———, *Nonpositive curvature and reflection groups*, in Handbook of Geometric Topology (Edited by R. J. Daverman and R. B. Sher), pp. 373–422, North-Holland, Amsterdam, 2002. MR1886674 (2002m:53061)
- [4] ———, *The cohomology of a Coxeter group with group ring coefficients*, Duke Math. J. **91** (no.2) (1998), 297–314. MR1600586 (99b:20067)
- [5] T. Hosaka, *Parabolic subgroups of finite index in Coxeter groups*, J. Pure Appl. Algebra **169** (2002), 215–227. MR1897344 (2003c:20041)
- [6] ———, *Dense subsets of the boundary of a Coxeter system*, Proc. Amer. Math. Soc. **132** (2004), 3441–3448. MR2073322
- [7] G. Moussong, *Hyperbolic Coxeter groups*, Ph.D. thesis, The Ohio State University, 1988.

DEPARTMENT OF MATHEMATICS, UTSUNOMIYA UNIVERSITY, UTSUNOMIYA, 321-8505, JAPAN
E-mail address: hosaka@cc.utsunomiya-u.ac.jp