LIFTS OF C_∞ - AND L_∞ -MORPHISMS TO G_∞ -MORPHISMS

GRÉGORY GINOT AND GILLES HALBOUT

(Communicated by Paul Goerss)

ABSTRACT. Let $\mathfrak{g}_2$ be the Hochschild complex of cochains on $C^\infty(\mathbb{R}^n)$ and let $\mathfrak{g}_1$ be the space of multivector fields on $\mathbb{R}^n$. In this paper we prove that given any G_∞ -structure (i.e. Gerstenhaber algebra up to homotopy structure) on $\mathfrak{g}_2$, and any C_∞ -morphism φ (i.e. morphism of a commutative, associative algebra up to homotopy) between $\mathfrak{g}_1$ and $\mathfrak{g}_2$, there exists a G_∞ -morphism Φ between $\mathfrak{g}_1$ and $\mathfrak{g}_2$ that restricts to φ . We also show that any L_∞ -morphism (i.e. morphism of a Lie algebra up to homotopy), in particular the one constructed by Kontsevich, can be deformed into a G_∞ -morphism, using Tamarkin's method for any G_∞ -structure on $\mathfrak{g}_2$. We also show that any two of such G_∞ -morphisms are homotopic.

0. INTRODUCTION

Let M be a differential manifold and let $\mathfrak{g}_2 = (C^\bullet(A, A), b)$ be the Hochschild cochain complex on $A = C^\infty(M)$. The classical Hochschild-Kostant-Rosenberg theorem states that the cohomology of $\mathfrak{g}_2$ is the graded Lie algebra $\mathfrak{g}_1 = \Gamma(M, \wedge^\bullet TM)$ of multivector fields on M . There is also a graded Lie algebra structure on $\mathfrak{g}_2$ given by the Gerstenhaber bracket. In particular $\mathfrak{g}_1$ and $\mathfrak{g}_2$ are also Lie algebras up to homotopy (L_∞ -algebra for short). In the case $M = \mathbb{R}^n$, using different methods, Kontsevich ([Ko1] and [Ko2]) and Tamarkin ([Ta]) have proved the existence of Lie homomorphisms “up to homotopy” (L_∞ -morphisms) from $\mathfrak{g}_1$ to $\mathfrak{g}_2$. Kontsevich's proof uses graph complex and is related to multizeta functions, whereas Tamarkin's construction uses the existence of Drinfeld's associators. In fact Tamarkin's L_∞ -morphism comes from the restriction of a Gerstenhaber algebra up to homotopy homomorphism (G_∞ -morphism) from $\mathfrak{g}_1$ to $\mathfrak{g}_2$. The G_∞ -algebra structure on $\mathfrak{g}_1$ is induced by its classical Gerstenhaber algebra structure and a far less trivial G_∞ -structure on $\mathfrak{g}_2$ was proved to exist by Tamarkin [Ta] and relies on Drinfeld's associator. Tamarkin's G_∞ -morphism also restricts into a commutative, associative up to homotopy morphism (C_∞ -morphism for short). The C_∞ -structure on $\mathfrak{g}_2$ (given by restriction of the G_∞ -one) highly depends on Drinfeld's associator, and any two choices of a Drinfeld associator yields *a priori* different C_∞ -structures. When M is a Poisson manifold, Kontsevich and Tamarkin homomorphisms imply the existence of a star-product (see [BFFLS1] and [BFFLS2] for a definition). A connection between the two approaches has been given in [KS] but the morphisms

Received by the editors April 1, 2003 and, in revised form, September 30, 2004.

2000 *Mathematics Subject Classification.* Primary 16E40, 53D55; Secondary 18D50, 16S80.

Key words and phrases. Deformation quantization, star-product, homotopy formulas, homological methods.

given by Kontsevich and Tamarkin are not the same. The aim of this paper is to show that, given any G_∞ -structure on $\mathfrak{g}_2$ and any C_∞ -morphism φ between $\mathfrak{g}_1$ and $\mathfrak{g}_2$, there exists a G_∞ -morphism Φ between $\mathfrak{g}_1$ and $\mathfrak{g}_2$ that restricts to φ . We also show that any L_∞ -morphism can be deformed into a G_∞ -one.

In the first section, we fix notation and recall the definitions of L_∞ - and G_∞ -structures. In the second section we state and prove the main theorem. In the last section we show that any two G_∞ -morphisms given by Tamarkin's method are homotopic.

Remark. In the sequel, unless otherwise stated, the manifold M is $\mathbb{R}^n$ for some $n \geq 1$. Most results could be generalized to other manifolds using techniques of Kontsevich [Ko1] (see also [TS], [CFT]).

1. C_∞ -, L_∞ - AND G_∞ -STRUCTURES

For any graded vector space $\mathfrak{g}$, we choose the following degree on $\wedge^\bullet \mathfrak{g}$: if $X_1, \dots, X_k$ are homogeneous elements of respective degree $|X_1|, \dots, |X_k|$, then

$$|X_1 \wedge \dots \wedge X_k| = |X_1| + \dots + |X_k| - k.$$

In particular the component $\mathfrak{g} = \wedge^1 \mathfrak{g} \subset \wedge^\bullet \mathfrak{g}$ is the same as the space $\mathfrak{g}$ with degree shifted by one. The space $\wedge^\bullet \mathfrak{g}$ with the deconcatenation cobracket is the cofree cocommutative coalgebra on $\mathfrak{g}$ with degree shifted by one (see [LS], Section 2). Any degree one map $d^k : \wedge^k \mathfrak{g} \rightarrow \mathfrak{g}$ ($k \geq 1$) extends into a derivation $d^k : \wedge^\bullet \mathfrak{g} \rightarrow \wedge^\bullet \mathfrak{g}$ of the coalgebra $\wedge^\bullet \mathfrak{g}$ by cofreeness property.

Definition 1.1. A vector space $\mathfrak{g}$ is endowed with a L_∞ -algebra (Lie algebras “up to homotopy”) structure if there are degree one linear maps $m^{1, \dots, 1}$, with k ones: $\wedge^k \mathfrak{g} \rightarrow \mathfrak{g}$ such that if we extend them to maps $\wedge^\bullet \mathfrak{g} \rightarrow \wedge^\bullet \mathfrak{g}$, then $d \circ d = 0$ where d is the derivation

$$d = m^1 + m^{1,1} + \dots + m^{1, \dots, 1} + \dots.$$

For more details on L_∞ -structures, see [LS]. It follows from the definition that a L_∞ -algebra structure induces a differential coalgebra structure on $\wedge^\bullet \mathfrak{g}$ and that the map $m^1 : \mathfrak{g} \rightarrow \mathfrak{g}$ is a differential. If $m^{1, \dots, 1} : \wedge^k \mathfrak{g} \rightarrow \mathfrak{g}$ are 0 for $k \geq 3$, we get the usual definition of (differential if $m^1 \neq 0$) graded Lie algebras.

For any graded vector space $\mathfrak{g}$, we denote $\underline{\mathfrak{g}}^{\otimes n}$ to be the quotient of $\mathfrak{g}^{\otimes n}$ by the image of all shuffles of length n (see [GK] or [GH] for details). The graded vector space $\bigoplus_{n \geq 0} \underline{\mathfrak{g}}^{\otimes n}$ is a quotient coalgebra of the tensor coalgebra $\bigoplus_{n \geq 0} \mathfrak{g}^{\otimes n}$. It is well known that this coalgebra $\bigoplus_{n \geq 0} \underline{\mathfrak{g}}^{\otimes n}$ is the cofree Lie coalgebra on the vector space $\mathfrak{g}$ (with degree shifted by minus one).

Definition 1.2. A C_∞ -algebra (commutative and associative “up to homotopy” algebra) structure on a vector space $\mathfrak{g}$ is given by a collection of degree one linear maps $m^k : \underline{\mathfrak{g}}^{\otimes k} \rightarrow \mathfrak{g}$ such that if we extend them to maps $\bigoplus \underline{\mathfrak{g}}^{\otimes \bullet} \rightarrow \bigoplus \underline{\mathfrak{g}}^{\otimes \bullet}$, then $d \circ d = 0$ where d is the derivation

$$d = m^1 + m^2 + m^3 + \dots.$$

In particular a C_∞ -algebra is an A_∞ -algebra.

For any space $\mathfrak{g}$, we denote by $\wedge^\bullet \underline{\mathfrak{g}}^{\otimes \bullet}$ the graded space

$$\wedge^\bullet \underline{\mathfrak{g}}^{\otimes \bullet} = \bigoplus_{m \geq 1, p_1 + \dots + p_n = m} \underline{\mathfrak{g}}^{\otimes p_1} \wedge \dots \wedge \underline{\mathfrak{g}}^{\otimes p_n}.$$

We use the following grading on $\wedge^\bullet \underline{\mathfrak{g}}^{\otimes \bullet}$: for $x_1^1, \dots, x_n^{p_n} \in \mathfrak{g}$, we define

$$|\underline{x_1^1} \otimes \dots \otimes \underline{x_1^{p_1}} \wedge \dots \wedge \underline{x_n^1} \otimes \dots \otimes \underline{x_n^{p_n}}| = \sum_{i_1}^{p_1} |x_1^{i_1}| + \dots + \sum_{i_n}^{p_n} |x_n^{i_n}| - n.$$

Note that the induced grading on $\wedge^\bullet \mathfrak{g} \subset \wedge^\bullet \underline{\mathfrak{g}}^{\otimes \bullet}$ is the same as the one introduced above. The cobracket on $\oplus \underline{\mathfrak{g}}^{\otimes \bullet}$ and the coproduct on $\wedge^\bullet \mathfrak{g}$ extend to a cobracket and a coproduct on $\wedge^\bullet \underline{\mathfrak{g}}^{\otimes \bullet}$ which yield a Gerstenhaber coalgebra structure on $\wedge^\bullet \underline{\mathfrak{g}}^{\otimes \bullet}$. It is well known that this coalgebra structure is cofree (see [Gi], Section 3, for example).

Definition 1.3. A G_∞ -algebra (Gerstenhaber algebra “up to homotopy”) structure on a graded vector space $\mathfrak{g}$ is given by a collection of degree one maps

$$m^{p_1, \dots, p_n} : \underline{\mathfrak{g}}^{\otimes p_1} \wedge \dots \wedge \underline{\mathfrak{g}}^{\otimes p_n} \rightarrow \mathfrak{g}$$

indexed by $p_1, \dots, p_n \geq 1$ such that their canonical extension $\wedge^\bullet \underline{\mathfrak{g}}^{\otimes \bullet} \rightarrow \wedge^\bullet \underline{\mathfrak{g}}^{\otimes \bullet}$ satisfies $d \circ d = 0$, where

$$d = \sum_{m \geq 1, p_1 + \dots + p_n = m} m^{p_1, \dots, p_n}.$$

Again, as the coalgebra structure of $\wedge^\bullet \underline{\mathfrak{g}}^{\otimes \bullet}$ is cofree, the map d makes $\wedge^\bullet \underline{\mathfrak{g}}^{\otimes \bullet}$ into a differential coalgebra. If the maps $m^{p_1, \dots, p_n}$ are 0 for $(p_1, p_2, \dots) \neq (1, 0, \dots)$, $(1, 1, 0, \dots)$ or $(2, 0, \dots)$, we get the usual definition of (differential if $m^1 \neq 0$) Gerstenhaber algebra.

The space of multivector fields $\mathfrak{g}_1$ is endowed with a graded Lie bracket $[-, -]_S$ called the Schouten bracket (see [Kos]). This Lie algebra can be extended into a Gerstenhaber algebra, with commutative structure given by the exterior product $(\alpha, \beta) \mapsto \alpha \wedge \beta$.

Setting $d_1 = m_1^{1,1} + m_1^2$, where $m_1^{1,1} : \wedge^2 \mathfrak{g}_1 \rightarrow \mathfrak{g}_1$, and $m_1^2 : \underline{\mathfrak{g}}_1^{\otimes 2} \rightarrow \mathfrak{g}_1$ are the extension of the Schouten bracket and the exterior product, we find that $(\mathfrak{g}_1, d_1)$ is a G_∞ -algebra.

In the same way, one can define a differential Lie algebra structure on the vector space $\mathfrak{g}_2 = C(A, A) = \bigoplus_{k \geq 0} C^k(A, A)$, the space of Hochschild cochains (generated by differential k -linear maps from A^k to A), where $A = C^\infty(M)$ is the algebra of smooth differential functions over M . Its bracket $[-, -]_G$, called the Gerstenhaber bracket, is defined, for $D, E \in \mathfrak{g}_2$, by

$$[D, E]_G = \{D|E\} - (-1)^{|E||D|} \{E|D\},$$

where

$$\{D|E\}(x_1, \dots, x_{d+e-1}) = \sum_{i \geq 0} (-1)^{|E| \cdot i} D(x_1, \dots, x_i, E(x_{i+1}, \dots, x_{i+e}), \dots).$$

The space $\mathfrak{g}_2$ has a grading defined by $|D| = k \Leftrightarrow D \in C^{k+1}(A, A)$ and its differential is $b = [m, -]_G$, where $m \in C^2(A, A)$ is the commutative multiplication on A .

Tamarkin (see [Ta] or also [GH]) stated the existence of a G_∞ -structure on $\mathfrak{g}_2$ (depending on a choice of a Drinfeld associator) given by a differential $d_2 = m_2^1 + m_2^{1,1} + m_2^2 + \dots + m_2^{p_1, \dots, p_n} + \dots$, on $\wedge^\bullet \underline{\mathfrak{g}}_2^{\otimes \bullet}$ satisfying $d_2 \circ d_2 = 0$. Although

this structure is non-explicit, it satisfies the following three properties:

- (a) m_2^1 is the extension of the differential b .
- (b) $m_2^{1,1}$ is the extension of the Gerstenhaber bracket $[-, -]_G$
- (1.1) and $m_2^{1,1,\dots,1} = 0$.
- (c) m_2^2 induces the exterior product in cohomology and the collection of the $(m^k)_{k \geq 1}$ defines a C_∞ -structure on $\mathfrak{g}_2$.

Definition 1.4. A L_∞ -morphism between two L_∞ -algebras $(\mathfrak{g}_1, d_1 = m_1^1 + \dots)$ and $(\mathfrak{g}_2, d_2 = m_2^1 + \dots)$ is a morphism of differential coalgebras

$$(1.2) \quad \varphi : (\wedge^\bullet \mathfrak{g}_1, d_1) \rightarrow (\wedge^\bullet \mathfrak{g}_2, d_2).$$

Such a map φ is uniquely determined by a collection of maps $\varphi^n : \wedge^n \mathfrak{g}_1 \rightarrow \mathfrak{g}_2$ (again by cofreeness properties). In the case $\mathfrak{g}_1$ and $\mathfrak{g}_2$ are respectively the graded Lie algebra $(\Gamma(M, \wedge TM), [-, -]_S)$ and the differential graded Lie algebra $(C(A, A), [-, -]_G)$, the formality theorems of Kontsevich and Tamarkin state the existence of a L_∞ -morphism between $\mathfrak{g}_1$ and $\mathfrak{g}_2$ such that φ^1 is the Hochschild-Kostant-Rosenberg quasi-isomorphism.

Definition 1.5. A morphism of C_∞ -algebras between two C_∞ -algebras $(\mathfrak{g}_1, d_1)$ and $(\mathfrak{g}_2, d_2)$ is a map $\phi : (\oplus \mathfrak{g}_1^{\otimes \bullet}, d_1) \rightarrow (\oplus \mathfrak{g}_2^{\otimes \bullet}, d_2)$ of codifferential coalgebras.

A C_∞ -morphism is in particular a morphism of A_∞ -algebras and is uniquely determined by maps $\partial^k : \mathfrak{g}^{\otimes k} \rightarrow \mathfrak{g}$.

Definition 1.6. A morphism of G_∞ -algebras between two G_∞ -algebras $(\mathfrak{g}_1, d_1)$ and $(\mathfrak{g}_2, d_2)$ is a map $\phi : (\wedge^\bullet \mathfrak{g}_1^{\otimes \bullet}, d_1) \rightarrow (\wedge^\bullet \mathfrak{g}_2^{\otimes \bullet}, d_2)$ of codifferential coalgebras.

There are coalgebra inclusions $\wedge^\bullet \mathfrak{g} \rightarrow \wedge^\bullet \mathfrak{g}^{\otimes \bullet}$, $\oplus \mathfrak{g}^{\otimes \bullet} \rightarrow \wedge^\bullet \mathfrak{g}^{\otimes \bullet}$, and it is easy to check that any G_∞ -morphism between two G_∞ -algebras $(\mathfrak{g}, \sum m^{p_1, \dots, p_n})$ and $(\mathfrak{g}', \sum m'^{p_1, \dots, p_n})$ restricts to a L_∞ -morphism $(\wedge^\bullet \mathfrak{g}, \sum m^{1, \dots, 1}) \rightarrow (\wedge^\bullet \mathfrak{g}', \sum m'^{1, \dots, 1})$ and a C_∞ -morphism $(\oplus \mathfrak{g}^{\otimes \bullet}, \sum m^k) \rightarrow (\oplus \mathfrak{g}'^{\otimes \bullet}, \sum m'^k)$. In the case where $\mathfrak{g}_1$ and $\mathfrak{g}_2$ are as above, Tamarkin's theorem states that there exists a G_∞ -morphism between the two G_∞ -algebras $\mathfrak{g}_1$ and $\mathfrak{g}_2$ (with the G_∞ structure he built) that restricts to a C_∞ - and a L_∞ -morphism.

2. MAIN THEOREM

We keep the notations of the previous section, in particular $\mathfrak{g}_2$ is the Hochschild complex of cochains on $C^\infty(M)$ and $\mathfrak{g}_1$ its cohomology. Here is our main theorem.

Theorem 2.1. *Given any G_∞ -structure d_2 on $\mathfrak{g}_2$ satisfying the three properties of (1.1), and any C_∞ -morphism φ between $\mathfrak{g}_1$ and $\mathfrak{g}_2$ such that φ^1 is the Hochschild-Kostant-Rosenberg map, there exists a G_∞ -morphism $\Phi : (\mathfrak{g}_1, d_1) \rightarrow (\mathfrak{g}_2, d_2)$ that restricts to φ .*

Also, given any L_∞ -morphism γ between $\mathfrak{g}_1$ and $\mathfrak{g}_2$ such that γ^1 is the Hochschild-Kostant-Rosenberg map, there exists a G_∞ -structure $(\mathfrak{g}_1, d'_1)$ on $\mathfrak{g}_1$ and G_∞ -morphism $\Gamma : (\mathfrak{g}_1, d'_1) \rightarrow (\mathfrak{g}_2, d_2)$ that restricts to γ . Moreover there exists a G_∞ -morphism $\Gamma' : (\mathfrak{g}_1, d_1) \rightarrow (\mathfrak{g}_1, d'_1)$.

In particular, Theorem 2.1 applies to the formality map of Kontsevich and also to any C_∞ -map derived (see [Ta], [GH]) from any B_∞ -structure on $\mathfrak{g}_2$ lifting the Gerstenhaber structure of $\mathfrak{g}_1$.

Let us first recall the proof of Tamarkin's formality theorem (see [GH] for more details):

1. First one proves there exists a G_∞ -structure on $\mathfrak{g}_2$, with differential d_2 , as in (1.1).
2. Then, one constructs a G_∞ -structure on $\mathfrak{g}_1$ given by a differential d'_1 together with a G_∞ -morphism Φ between $(\mathfrak{g}_1, d'_1)$ and $(\mathfrak{g}_2, d_2)$.
3. Finally, one constructs a G_∞ -morphism Φ' between $(\mathfrak{g}_1, d_1)$ and $(\mathfrak{g}_1, d'_1)$.

The composition $\Phi \circ \Phi'$ is then a G_∞ -morphism between $(\mathfrak{g}_1, d_1)$ and $(\mathfrak{g}_2, d_2)$, thus it restricts to a L_∞ -morphism between the differential graded Lie algebras $\mathfrak{g}_1$ and $\mathfrak{g}_2$.

We suppose now that, in the first step, we take any G_∞ -structure on $\mathfrak{g}_2$ given by a differential d_2 . We also suppose we are given a C_∞ -morphism φ and a L_∞ -morphism γ between $\mathfrak{g}_1$ and $\mathfrak{g}_2$ satisfying $\gamma^1 = \varphi^1 = \varphi_{\text{HKR}}$, the Hochschild-Kostant-Rosenberg quasi-isomorphism.

Proof of Theorem 2.1. The theorem will follow if we prove that steps 2 and 3 of Tamarkin's construction are still true with the extra conditions that the restriction of the G_∞ -morphism Φ (resp. Φ') on the C_∞ -structures is the C_∞ -morphism $\varphi : \mathfrak{g}_1 \rightarrow \mathfrak{g}_2$ (resp. id).

Let us recall (see [GH]) that the constructions of Φ and d'_1 can be made by induction. For $i = 1, 2$ and $n \geq 0$, let us set

$$V_i^{[n]} = \bigoplus_{p_1 + \dots + p_k = n} \underline{\mathfrak{g}_i^{\otimes p_1}} \wedge \dots \wedge \underline{\mathfrak{g}_i^{\otimes p_k}}$$

and $V_i^{[\leq n]} = \sum_{k \leq n} V_i^{[k]}$. Let $d_2^{[n]}$ and $d_2^{[\leq n]}$ be the sums

$$d_2^{[n]} = \sum_{p_1 + \dots + p_k = n} d_2^{p_1, \dots, p_k} \quad \text{and} \quad d_2^{[\leq n]} = \sum_{p \leq n} d_2^{[p]}.$$

Clearly, $d_2 = \sum_{n \geq 1} d_2^{[n]}$. In the same way, we denote $d'_1 = \sum_{n \geq 1} d'_1^{[n]}$ by

$$d'_1^{[n]} = \sum_{p_1 + \dots + p_k = n} d'_1^{p_1, \dots, p_k} \quad \text{and} \quad d'_1^{[\leq n]} = \sum_{1 \leq k \leq n} d'_1^{[k]}.$$

We know from Section 1 that a morphism $\Phi : (\wedge^\bullet \underline{\mathfrak{g}_1^{\otimes \bullet}}, d'_1) \rightarrow (\wedge^\bullet \underline{\mathfrak{g}_2^{\otimes \bullet}}, d_2)$ is uniquely determined by its components $\Phi^{p_1, \dots, p_k} : \underline{\mathfrak{g}_1^{\otimes p_1}} \wedge \dots \wedge \underline{\mathfrak{g}_1^{\otimes p_k}} \rightarrow \mathfrak{g}_2$. Again, we have $\Phi = \sum_{n \geq 1} \Phi^{[n]}$ with

$$\Phi^{[n]} = \sum_{p_1 + \dots + p_k = n} \Phi^{p_1, \dots, p_k} \quad \text{and} \quad \Phi^{[\leq n]} = \sum_{1 \leq k \leq n} \Phi^{[k]}.$$

We want to construct the maps $d'_1^{[n]}$ and $\Phi^{[n]}$ by induction with the initial condition

$$d'_1^{[1]} = 0 \quad \text{and} \quad \Phi^{[1]} = \varphi_{\text{HKR}},$$

where $\varphi_{\text{HKR}} : (\mathfrak{g}_1, 0) \rightarrow (\mathfrak{g}_2, b)$ is the Hochschild-Kostant-Rosenberg quasi-isomorphism (see [HKR]) defined, for $\alpha \in \mathfrak{g}_1$, $f_1, \dots, f_n \in A$, by

$$\varphi_{\text{HKR}} : \alpha \mapsto ((f_1, \dots, f_n) \mapsto \langle \alpha, df_1 \wedge \dots \wedge df_n \rangle).$$

Moreover, we want the following extra conditions to be true:

$$(2.3) \quad \Phi^{k \geq 2} = \varphi^k, \quad d_1'^2 = d_1^2, \quad d_1'^{k \geq 3} = 0.$$

Now suppose the construction is done for $n-1$ ($n \geq 2$), i.e., we have built maps $(d_1'^{[i]})_{i \leq n-1}$ and $(\Phi^{[i]})_{i \leq n-1}$ satisfying conditions (2.3) and

$$(2.4) \quad \begin{aligned} \Phi^{[\leq n-1]} \circ d_1'^{[\leq n-1]} &= d_2^{[\leq n-1]} \circ \Phi^{[\leq n-1]} \text{ on } V_1^{[\leq n-1]} \text{ and} \\ d_1'^{[\leq n-1]} \circ d_1'^{[\leq n-1]} &= 0 \text{ on } V_1^{[\leq n]}. \end{aligned}$$

In [GH], we prove that for any such $(d_1'^{[i]})_{i \leq n-1}$ and $(\Phi^{[i]})_{i \leq n-1}$, one can construct $d_1'^{[n]}$ and $\Phi^{[n]}$ such that condition (2.4) is true for n instead of $n-1$. To complete the proof of Theorem 2.1 (step 2), we have to show that $d_1'^{[n]}$ and $\Phi^{[n]}$ can be chosen to satisfy conditions (2.3). In equation (2.4), the terms $d_1'^k$ and Φ^k only act on V_1^k . So one can replace Φ^n with φ^n , $d_1'^2$ with d_1^2 (or $d_1'^i$, $i \geq 3$ with 0) provided conditions (2.4) are still satisfied on V_1^n . The other terms acting on V_1^n in equation (2.4) only involve terms $\Phi^m = \varphi^m$ and $d_1'^m$. Then conditions (2.4) on $V_1^{1, \dots, 1}$ are the equations that should be satisfied by a C_∞ -morphism between the C_∞ -algebras $(\mathfrak{g}_1, d_1'^{1,1} = d_1^{1,1})$ and $(\mathfrak{g}_2, \sum_{k \geq 1} d_2^k)$ restricted to V_1^n . Hence by hypothesis on φ the conditions hold.

Similarly the construction of Φ' can be made by induction. Let us recall the proof given in [GH]. Again a morphism $\Phi' : (\wedge^\bullet \underline{\mathfrak{g}}_1^{\otimes \bullet}, d_1) \rightarrow (\wedge^\bullet \underline{\mathfrak{g}}_2^{\otimes \bullet}, d_1')$ is uniquely determined by its components $\Phi'^{p_1, \dots, p_k} : \underline{\mathfrak{g}}_1^{\otimes p_1} \wedge \dots \wedge \underline{\mathfrak{g}}_1^{\otimes p_k} \rightarrow \underline{\mathfrak{g}}_1$. We write $\Phi' = \sum_{n \geq 1} \Phi'^{[n]}$ with

$$\Phi'^{[n]} = \sum_{p_1 + \dots + p_k = n} \Phi'^{p_1, \dots, p_k} \quad \text{and} \quad \Phi'^{[\leq n]} = \sum_{1 \leq k \leq n} \Phi'^{[k]}.$$

We construct the maps $\Phi'^{[n]}$ by induction with the initial condition $\Phi'^{[1]} = \text{id}$. Moreover, we want the following extra conditions to be true:

$$(2.5) \quad \Phi'^n = 0 \text{ for } n \geq 2.$$

Now suppose the construction is done for $n-1$ ($n \geq 2$), i.e., we have built maps $(\Phi'^{[i]})_{i \leq n-1}$ satisfying conditions (2.5) and

$$(2.6) \quad \Phi'^{[\leq n-1]} d_1'^{[\leq n]} = d_1'^{[\leq n]} \Phi'^{[\leq n-1]} \text{ on } V_1^{[\leq n]}.$$

In [GH], we prove that for any such $(\Phi'^{[i]})_{i \leq n-1}$, one can construct $\Phi'^{[n]}$ such that condition (2.6) is true for n instead of $n-1$ in the following way: making the equation $\Phi' d_1 = d_1' \Phi'$ on $V_1^{[n+1]}$ explicit, we get

$$(2.7) \quad \Phi'^{[\leq n]} d_1'^{[\leq n+1]} = d_1'^{[\leq n+1]} \Phi'^{[\leq n]}.$$

If we now take into account that $d_1'^{[i]} = 0$ for $i \neq 2$, $d_1'^{[1]} = 0$ and that on $V_1^{[n+1]}$ we have $\Phi'^{[k]} d_1'^{[l]} = d_1'^{[\leq k]} \Phi'^{[l]} = 0$ for $k+l > n+2$, the identity (2.7) becomes

$$\Phi'^{[\leq n]} d_1'^{[2]} = \sum_{k=2}^{n+1} d_1'^{[k]} \Phi'^{[\leq n-k+2]}.$$

Since $d_1'^{[2]} = d_1^{[2]}$, (2.7) is equivalent to

$$d_1^{[2]} \Psi'^{[\leq n]} - \Phi'^{[\leq n]} d_1^{[2]} = \left[d_1^{[2]}, \Phi'^{[\leq n]} \right] = - \sum_{k=3}^{n+1} d_1'^{[k]} \Phi'^{[\leq n-k+2]}.$$

Note that $d_1^{[2]} = m_1^{1,1} + m_1^2$. Then the construction will be possible when the term $\sum_{k=3}^{n+1} d_1'^{[k]} \Psi'^{[\leq n-k+2]}$ is a coboundary in the subcomplex of $(\text{End}(\wedge^\bullet \underline{\mathfrak{g}}_1^{\otimes \bullet}), [d_1^{[2]}, -])$ consisting of maps which restrict to zero on $\bigoplus_{n \geq 2} \underline{\mathfrak{g}}_1^{\otimes n}$. It is always a cocycle by straightforward computation (see [GH]), and the subcomplex is acyclic because both $(\text{End}(\wedge^\bullet \underline{\mathfrak{g}}_1^{\otimes \bullet}), [d_1^{[2]}, -])$ and the Harrison cohomology of $\mathfrak{g}_1$ are trivial according to Tamarkin [Ta] (see also [GH], Proposition 5.1, and [Hi], 5.4).

In the case of the L_∞ -morphism γ , the first step is similar: the fact that γ is a L_∞ -map enables us to build a G_∞ -structure $(\mathfrak{g}_1, d_1')$ on $\mathfrak{g}_1$ and a G_∞ -morphism $\Gamma : (\mathfrak{g}_1, d_1') \rightarrow (\mathfrak{g}_2, d_2)$ such that:

$$(2.8) \quad \Gamma^{1, \dots, 1} = \gamma^{1, \dots, 1}, \quad d_1'^{1,1} = d_1^{1,1}, \quad d_1'^{1,1, \dots, 1} = 0.$$

For the second step, we have to build a map Γ' satisfying the equation

$$d_1^{[2]} \Gamma'^{[\leq n]} - \Gamma'^{[\leq n]} d_1^{[2]} = \left[d_1^{[2]}, \Gamma'^{[\leq n]} \right] = - \sum_{k=3}^{n+1} d_1'^{[k]} \Gamma'^{[\leq n-k+2]}$$

on $V_1^{[n+1]}$ for any $n \geq 1$. Again, because Tamarkin has proved that the complex $(\text{End}(\wedge^\bullet \underline{\mathfrak{g}}_1^{\otimes \bullet}), [d_1^{[2]}, -])$ is acyclic (we are in the case $M = \mathbb{R}^n$), the result follows from the fact that $\sum_{k=3}^{n+1} d_1'^{[k]} \Gamma'^{[\leq n-k+2]}$ is a cocycle. The difference with the C_∞ -case is that the $\Gamma'^{1, \dots, 1}$ could be non-zero. $\square$

3. THE DIFFERENCE BETWEEN TWO G_∞ -MAPS

In this section we investigate the difference between two different G_∞ -formality maps.

We fix once and for all a G_∞ -structure on $\mathfrak{g}_2$ (given by a differential d_2) satisfying the conditions (1.1) and a morphism of G_∞ -algebras $T : (\mathfrak{g}_1, d_1) \rightarrow (\mathfrak{g}_2, d_2)$ such that $T^1 : \mathfrak{g}_1 \rightarrow \mathfrak{g}_2$ is φ_{HKR} . Let $K : (\mathfrak{g}_1, d_1) \rightarrow (\mathfrak{g}_2, d_2)$ be any other G_∞ -morphism with $K^1 = \varphi_{\text{HKR}}$ (for example any lift of a Kontsevich formality map or any G_∞ -maps lifting another C_∞ -morphism).

Theorem 3.1. *There exists a map $h : \wedge^\bullet \underline{\mathfrak{g}}_1^{\otimes \bullet} \rightarrow \wedge^\bullet \underline{\mathfrak{g}}_2^{\otimes \bullet}$ such that*

$$T - K = h \circ d_1 + d_2 \circ h.$$

In other words the formality G_∞ -morphisms K and T are homotopic.

The maps T and K are elements of the cochain complex $(\text{Hom}(\wedge^\bullet \underline{\mathfrak{g}}_1^{\otimes \bullet}, \wedge^\bullet \underline{\mathfrak{g}}_2^{\otimes \bullet}), \delta)$ with differential given, for all $f \in \text{Hom}(\wedge^\bullet \underline{\mathfrak{g}}_1^{\otimes \bullet}, \wedge^\bullet \underline{\mathfrak{g}}_2^{\otimes \bullet})$, $|f| = k$, by

$$\delta(f) = d_2 \circ f - (-1)^k f \circ d_1.$$

We first compare this cochain complex with the complexes $(\text{End}(\wedge^\bullet \underline{\mathfrak{g}}_1^{\otimes \bullet}), [d_1; -])$ and $(\text{End}(\wedge^\bullet \underline{\mathfrak{g}}_2^{\otimes \bullet}), [d_2; -])$, where $[-; -]$ is the graded commutator of morphisms. There are morphisms

$$T_* : \text{End}(\wedge^\bullet \underline{\mathfrak{g}}_1^{\otimes \bullet}) \rightarrow \text{Hom}(\wedge^\bullet \underline{\mathfrak{g}}_1^{\otimes \bullet}, \wedge^\bullet \underline{\mathfrak{g}}_2^{\otimes \bullet}), \quad T^* : \text{End}(\wedge^\bullet \underline{\mathfrak{g}}_2^{\otimes \bullet}) \rightarrow \text{Hom}(\wedge^\bullet \underline{\mathfrak{g}}_1^{\otimes \bullet}, \wedge^\bullet \underline{\mathfrak{g}}_2^{\otimes \bullet})$$

defined, for $f \in \text{End}(\wedge^{\bullet} \underline{\mathfrak{g}}_2^{\otimes \bullet})$ and $g \in \text{Hom}(\wedge^{\bullet} \underline{\mathfrak{g}}_1^{\otimes \bullet}, \wedge^{\bullet} \underline{\mathfrak{g}}_2^{\otimes \bullet})$, by

$$T_*(f) = T \circ f, \quad T^*(g) = g \circ T.$$

Lemma 3.2. *The morphisms*

$$\begin{aligned} T_* : \left(\text{End}(\wedge^{\bullet} \underline{\mathfrak{g}}_1^{\otimes \bullet}), [d_1; -] \right) &\rightarrow \left(\text{Hom}(\wedge^{\bullet} \underline{\mathfrak{g}}_1^{\otimes \bullet}, \wedge^{\bullet} \underline{\mathfrak{g}}_2^{\otimes \bullet}), \delta \right) \\ &\leftarrow \left(\text{End}(\wedge^{\bullet} \underline{\mathfrak{g}}_2^{\otimes \bullet}), [d_2; -] \right) : T^* \end{aligned}$$

of cochain complexes are quasi-isomorphisms.

Remark. This lemma holds for every manifold M and any G_{∞} -morphism $T : (\mathfrak{g}_1, d_1) \rightarrow (\mathfrak{g}_2, d_2)$.

Proof. First we show that T_* is a morphism of complexes. Let $f \in \text{End}(\wedge^{\bullet} \underline{\mathfrak{g}}_2^{\otimes \bullet})$ with $|f| = k$; then

$$\begin{aligned} T_*([d_1; f]) &= T \circ d_1 \circ f - (-1)^k T \circ f \circ d_1 \\ &= d_2 \circ (T \circ f) - (-1)^k (T \circ f) \circ d_1 \\ &= \delta(T_*(f)). \end{aligned}$$

Let us now prove that T_* is a quasi-isomorphism. For any graded vector space $\mathfrak{g}$, the space $\wedge^{\bullet} \underline{\mathfrak{g}}^{\otimes \bullet}$ has the structure of a filtered space where the m -level of the filtration is $F^m(\wedge^{\bullet} \underline{\mathfrak{g}}^{\otimes \bullet}) = \bigoplus_{p_1 + \dots + p_n - 1 \leq m} \underline{\mathfrak{g}}^{\otimes p_1} \wedge \dots \wedge \underline{\mathfrak{g}}^{\otimes p_n}$. Clearly the differential d_1 and d_2 are compatible with the filtrations on $\wedge^{\bullet} \underline{\mathfrak{g}}_1^{\otimes \bullet}$ and $\wedge^{\bullet} \underline{\mathfrak{g}}_2^{\otimes \bullet}$, hence $\text{End}(\wedge^{\bullet} \underline{\mathfrak{g}}_1^{\otimes \bullet})$ and $\text{Hom}(\wedge^{\bullet} \underline{\mathfrak{g}}_1^{\otimes \bullet}, \wedge^{\bullet} \underline{\mathfrak{g}}_2^{\otimes \bullet})$ are filtered cochain complex. This yields two spectral sequences (lying in the first quadrant) $E_{\bullet, \bullet}^*$ and $\tilde{E}_{\bullet, \bullet}^*$ which converge respectively toward the cohomology $H^*(\text{End}(\wedge^{\bullet} \underline{\mathfrak{g}}_1^{\otimes \bullet}))$ and $H^*(\text{Hom}(\wedge^{\bullet} \underline{\mathfrak{g}}_1^{\otimes \bullet}, \wedge^{\bullet} \underline{\mathfrak{g}}_2^{\otimes \bullet}))$. By standard spectral sequence techniques it is enough to prove that the map $T_*^0 : E_0^{\bullet, \bullet} \rightarrow \tilde{E}_0^{\bullet, \bullet}$ induced by T_* on the associated graded is a quasi-isomorphism.

The induced differentials on $E_0^{\bullet, \bullet}$ and $\tilde{E}_0^{\bullet, \bullet}$ are respectively $[d_1^1, -] = 0$ and $d_2^1 \circ (-) - (-) \circ d_1^1 = b \circ (-)$, where b is the Hochschild coboundary. By cofreeness property we have the following two isomorphisms:

$$E_0^{\bullet, \bullet} \cong \text{End}(\mathfrak{g}_1), \quad \tilde{E}_0^{\bullet, \bullet} \cong \text{Hom}(\mathfrak{g}_1, \mathfrak{g}_2).$$

The map $T_*^0 : E_0^{\bullet, \bullet} \rightarrow \tilde{E}_0^{\bullet, \bullet}$ induced by T_* is $\varphi_{\text{HKR}} \circ (-)$. Let $p : \mathfrak{g}_2 \rightarrow \mathfrak{g}_1$ be the projection onto the cohomology, *i.e.* $p \circ \varphi_{\text{HKR}} = \text{id}$. Let $u : \mathfrak{g}_1 \rightarrow \mathfrak{g}_2$ be any map satisfying $b(u) = 0$ and set $v = p \circ u \in \text{End}(\mathfrak{g}_1)$. One can choose a map $w : \mathfrak{g}_1 \rightarrow \mathfrak{g}_2$ which satisfies for any $x \in \mathfrak{g}_1$ the following identity:

$$\varphi_{\text{HKR}} \circ p \circ u(x) - u(x) = b \circ w(x).$$

It follows that $\varphi_{\text{HKR}}(v)$ has the same class of homology as u which proves the surjectivity of T_*^0 in cohomology. The identity $p \circ \varphi_{\text{HKR}} = \text{id}$ easily implies that T_*^0 is also injective in cohomology which finishes the proof of the lemma for T_* .

The proof that T^* is also a quasi-isomorphism is analogous. $\square$

Proof of Theorem 3.1. It is easy to check that $T - K$ is a cocycle in

$$\left(\text{Hom}(\wedge^{\bullet} \underline{\mathfrak{g}}_1^{\otimes \bullet}, \wedge^{\bullet} \underline{\mathfrak{g}}_2^{\otimes \bullet}), \delta \right).$$

The complex of cochain $\left(\text{End}(\wedge^\bullet \underline{\mathfrak{g}}_1^{\otimes \bullet}), [d_1, -]\right) \cong \left(\text{Hom}(\wedge^\bullet \underline{\mathfrak{g}}_1^{\otimes \bullet}, \underline{\mathfrak{g}}_1), [d_1, -]\right)$ is tri-graded with $||_1$ being the degree coming from the graduation of $\underline{\mathfrak{g}}_1$ and any element x lying in $\underline{\mathfrak{g}}_1^{\otimes p_1} \wedge \cdots \wedge \underline{\mathfrak{g}}_1^{\otimes p_q}$ satisfies $|x|_2 = q - 1$, $|x|_3 = p_1 + \cdots + p_q - q$. In the case $M = \mathbb{R}^n$, the cohomology $H^\bullet \left(\text{End}(\wedge^\bullet \underline{\mathfrak{g}}_1^{\otimes \bullet}), [d_1, -]\right)$ is concentrated in bidegree $(||_2, ||_3) = (0, 0)$ (see [Ta], [Hi]). By Lemma 3.2, this is also the case for the cochain complex $\left(\text{Hom}(\wedge^\bullet \underline{\mathfrak{g}}_1^{\otimes \bullet}, \wedge^\bullet \underline{\mathfrak{g}}_2^{\otimes \bullet}), \delta\right)$. Thus, its cohomology classes are determined by complex morphisms $(\underline{\mathfrak{g}}_1, 0) \rightarrow (\underline{\mathfrak{g}}_2, d_2^1)$, and it is enough to prove that T and K determine the same complex morphism $(\underline{\mathfrak{g}}_1, 0) \rightarrow (\underline{\mathfrak{g}}_2, d_2^1 = b)$ which is clear because T^1 and K^1 are both equal to the Hochschild-Kostant-Rosenberg map. $\square$

Remark 3.3. It is possible to have an explicit formula for the map h in Theorem 3.1. In fact the quasi-isomorphism coming from Lemma 3.2 can be made explicit using explicit homotopy formulae for the Hochschild-Kostant-Rosenberg map (see [Ha] for example) and deformation retract techniques (instead of spectral sequences) as in [Ka]. The same techniques also apply to give explicit formulae for the quasi-isomorphism giving the acyclicity of $\left(\text{End}(\wedge^\bullet \underline{\mathfrak{g}}_1^{\otimes \bullet}), [d_1, -]\right)$ in the proof of Theorem 3.1 (see [GH] for example).

REFERENCES

- [BFFLS1] F. Bayen, M. Flato, C. Fronsdal, A. Lichnerowicz, D. Sternheimer, *Quantum mechanics as a deformation of classical mechanics*, Lett. Math. Phys. **1** (1975), 521–530. MR0674337 (58:32617)
- [BFFLS2] F. Bayen, M. Flato, C. Fronsdal, A. Lichnerowicz, D. Sternheimer, *Deformation theory and quantization, I and II*, Ann. Phys. **111** (1977), 61–151. MR496157 (58:14737a); MR0496158 (58:14737b)
- [CFT] A. S. Cattaneo, G. Felder, L. Tomassini, *From local to global deformation quantization of Poisson manifolds*, Duke Math. J. **115** (2002), 329–352. MR1944574 (2004a:53114)
- [Gi] G. Ginot, *Homologie et modèle minimal des algèbres de Gerstenhaber*, Ann. Math. Blaise Pascal **11** (2004), 91–127. MR2077240
- [GH] G. Ginot, G. Halbout, *A formality theorem for Poisson manifold*, Lett. Math. Phys. **66** (2003), 37–64. MR2064591
- [GK] V. Ginzburg, M. Kapranov, *Koszul duality for operads*, Duke Math. J. **76** (1994), 203–272. MR1301191 (96a:18004)
- [Ha] G. Halbout, *Formule d’homotopie entre les complexes de Hochschild et de de Rham*, Compositio Math. **126** (2001), 123–145. MR1827641 (2002d:16010)
- [Hi] V. Hinich, *Tamarkin’s proof of Kontsevich’s formality theorem*, Forum Math. **15** (2003), 591–614. MR1978336 (2004j:17029)
- [HKR] G. Hochschild, B. Kostant and A. Rosenberg, *Differential forms on regular affine algebras*, Transactions AMS **102** (1962), 383–408. MR0142598 (26:167)
- [Ka] C. Kassel, *Homologie cyclique, caractère de Chern et lemme de perturbation*, J. Reine Angew. Math. **408** (1990), 159–180. MR1058987 (92d:19002)
- [Ko1] M. Kontsevich, *Formality conjecture. Deformation theory and symplectic geometry*, Math. Phys. Stud. **20** (1996), 139–156. MR1480721 (98m:58044)
- [Ko2] M. Kontsevich, *Deformation quantization of Poisson manifolds*, Lett. Math. Phys. **66** (2003), 157–216. MR2062626
- [KS] M. Kontsevich, Y. Soibelman, *Deformations of algebras over operads and the Deligne conjecture*, Math. Phys. Stud. **21** (2000), 255–307. MR1805894 (2002e:18012)
- [Kos] J. L. Koszul, *Crochet de Schouten-Nijenhuis et cohomologie in “Elie Cartan et les mathématiques d’aujourd’hui”*, Astérisque (1985), 257–271. MR0837203 (88m:17013)
- [LS] T. Lada, J. D. Stasheff, *Introduction to SH Lie algebras for physicists*, Internat. J. Theoret. Phys. **32** (1993), 1087–1103. MR1235010 (94g:17059)

- [Ta] D. Tamarkin, *Another proof of M. Kontsevich's formality theorem*, math. QA/9803025.
- [TS] D. Tamarkin, B. Tsygan, *Noncommutative differential calculus, homotopy BV algebras and formality conjectures*, Methods Funct. Anal. Topology **6** (2000), 85–97. MR1783778 (2001i:16017)

LABORATOIRE ANALYSE GÉOMÉTRIE ET APPLICATIONS, UNIVERSITÉ PARIS 13 ET ÉCOLE NORMALE SUPÉRIEURE DE CACHAN, FRANCE

E-mail address: `ginot@cmla.ens-cachan.fr`

INSTITUTE DE RECHERCHE MATHÉMATIQUE AVANCÉE, UNIVERSITÉ LOUIS PASTEUR ET CNRS, STRASBOURG, FRANCE

E-mail address: `halbout@math.u-strasbg.fr`