THE SET OF COMMON FIXED POINTS OF A ONE-PARAMETER CONTINUOUS SEMIGROUP OF MAPPINGS IS $F(T(1)) \cap F(T(\sqrt{2}))$

TOMONARI SUZUKI

(Communicated by Jonathan M. Borwein)

ABSTRACT. In this paper we prove the following theorem: Let $\{T(t) : t \geq 0\}$ be a one-parameter continuous semigroup of mappings on a subset C of a Banach space E . The set of all fixed points of $T(t)$ is denoted by $F(T(t))$ for each $t \geq 0$. Then

$$\bigcap_{t \geq 0} F(T(t)) = F(T(1)) \cap F(T(\sqrt{2}))$$

holds. Using this theorem, we discuss convergence theorems to a common fixed point of $\{T(t) : t \geq 0\}$.

1. INTRODUCTION

Let C be a subset of a Banach space E , and let T be a *nonexpansive mapping* on C , i.e., $\|Tx - Ty\| \leq \|x - y\|$ for all $x, y \in C$. We know that T has a fixed point in the case that E is uniformly convex and C is bounded, closed and convex; see Browder [5], Göhde [9], and Kirk [13]. We denote by $F(T)$ the set of all fixed points of T .

Let τ be a Hausdorff topology on E . A family of mappings $\{T(t) : t \geq 0\}$ is called a *one-parameter τ -continuous semigroup of mappings* on C if the following are satisfied:

- (sg 1) $T(s + t) = T(s) \circ T(t)$ for all $s, t \geq 0$;
- (sg 2) for each $x \in C$, the mapping $t \mapsto T(t)x$ from $[0, \infty)$ into C is continuous with respect to τ .

As the topology τ , we usually consider the strong topology of E . A family of mappings $\{T(t) : t \geq 0\}$ is called a *one-parameter τ -continuous semigroup of nonexpansive mappings* on C (*nonexpansive semigroup*, for short) if (sg 1), (sg 2) and the following (sg 3) are satisfied:

- (sg 3) for each $t \geq 0$, $T(t)$ is a nonexpansive mapping on C .

We know that $\{T(t) : t \geq 0\}$ has a common fixed point in the case that E is uniformly convex and C is bounded, closed and convex; see Browder [5]. Moreover,

Received by the editors December 17, 2003.

2000 *Mathematics Subject Classification*. Primary 47H20, 47H10.

Key words and phrases. Nonexpansive semigroup, common fixed point, irrational number.

The author was supported in part by Grants-in-Aid for Scientific Research from the Japanese Ministry of Education, Culture, Sports, Science and Technology.

©2005 American Mathematical Society
 Reverts to public domain 28 years from publication

in 1974, Bruck [8] proved that $\{T(t) : t \geq 0\}$ has a common fixed point in the case that C is weakly compact, convex, and has the fixed point property for nonexpansive mappings.

In this paper we prove the following theorem: Let E be a Banach space and let τ be a Hausdorff topology on E . Let $\{T(t) : t \geq 0\}$ be a one-parameter τ -continuous semigroup of mappings on a subset C of E . Then

$$\bigcap_{t \geq 0} F(T(t)) = F(T(1)) \cap F(T(\sqrt{2}))$$

holds. Using this theorem, we discuss convergence theorems to a common fixed point of $\{T(t) : t \geq 0\}$.

2. PRELIMINARIES

Throughout this paper we denote by $\mathbb{Q}$ the set of all rational numbers, and by $\mathbb{N}$ the set of all positive integers. For a real number t , we denote by $[t]$ the maximum integer not exceeding t . It is obvious that for each real number t , there exists $\varepsilon \in [0, 1)$ such that $t = [t] + \varepsilon$.

We recall that a Banach space E is called *strictly convex* if $\|x + y\|/2 < 1$ for all $x, y \in E$ with $\|x\| = \|y\| = 1$ and $x \neq y$. E is called *uniformly convex* if for each $\varepsilon > 0$, there exists $\delta > 0$ such that $\|x + y\|/2 < 1 - \delta$ for all $x, y \in E$ with $\|x\| = \|y\| = 1$ and $\|x - y\| \geq \varepsilon$. It is clear that a uniformly convex Banach space is strictly convex. The norm of E is called *Fréchet differentiable* if for each $x \in E$ with $\|x\| = 1$, $\lim_{t \rightarrow 0} (\|x + ty\| - \|x\|)/t$ exists and is attained uniformly in $y \in E$ with $\|y\| = 1$.

The following lemma is a corollary of Bruck's result in [7].

Lemma 1 (Bruck [7]). *Let C be a subset of a strictly convex Banach space E . Let S and T be nonexpansive mappings from C into E with a common fixed point. Then for each $\lambda \in (0, 1)$, a mapping U from C into E defined by $Ux = \lambda Sx + (1 - \lambda)Tx$ for $x \in C$ is nonexpansive and $F(U) = F(S) \cap F(T)$ holds.*

Proof. It is obvious that $F(U) \supset F(S) \cap F(T)$. Fix $x \in F(U)$ and $w \in F(S) \cap F(T)$. Then we have

$$\begin{aligned} \|x - w\| &= \|\lambda Sx + (1 - \lambda)Tx - w\| \\ &\leq \lambda \|Sx - w\| + (1 - \lambda) \|Tx - w\| \\ &\leq \lambda \|x - w\| + (1 - \lambda) \|x - w\| \\ &= \|x - w\| \end{aligned}$$

and hence

$$\|x - w\| = \|\lambda Sx + (1 - \lambda)Tx - w\| = \|Sx - w\| = \|Tx - w\|.$$

So, from the strict convexity of E , we obtain

$$x = \lambda Sx + (1 - \lambda)Tx = Sx = Tx.$$

That is, $x \in F(S) \cap F(T)$. This completes the proof. $\square$

The following four convergence theorems for nonexpansive mappings are well known.

Theorem 1 (Baillon [2]). *Let T be a nonexpansive mapping on a bounded closed convex subset C of a Hilbert space E . Let $x \in C$ and define a sequence $\{x_n\}$ in C by $x_n = \sum_{k=1}^n T^k x / n$ for $n \in \mathbb{N}$. Then $\{x_n\}$ converges weakly to a fixed point of T .*

Theorem 2 (Reich [17]). *Let E be a uniformly convex Banach space whose norm is Fréchet differentiable. Let T be a nonexpansive mapping on a bounded closed convex subset C of E . Define a sequence $\{x_n\}$ in C by $x_1 \in C$ and $x_{n+1} = \alpha_n T x_n + (1 - \alpha_n) x_n$ for $n \in \mathbb{N}$, where $\{\alpha_n\}$ is a sequence in $[0, 1]$ satisfying $\sum_{n=1}^{\infty} \alpha_n (1 - \alpha_n) = \infty$. Then $\{x_n\}$ converges weakly to a fixed point of T .*

Theorem 3 (Browder [6]). *Let T be a nonexpansive mapping on a bounded closed convex subset C of a Hilbert space E . Let $\{\lambda_n\}$ be a sequence in $(0, 1)$ converging to 0. Fix $u \in C$ and define a sequence $\{x_n\}$ in C by $x_n = (1 - \lambda_n) T x_n + \lambda_n u$ for $n \in \mathbb{N}$. Then $\{x_n\}$ converges strongly to a fixed point of T .*

Theorem 4 (Wittmann [24]). *Let T be a nonexpansive mapping on a bounded closed convex subset C of a Hilbert space E . Let $u \in C$ and define a sequence $\{x_n\}$ in C by $x_1 \in C$ and $x_{n+1} = (1 - \lambda_n) T x_n + \lambda_n u$ for $n \in \mathbb{N}$, where $\{\lambda_n\}$ is a sequence in $[0, 1]$ satisfying $\lim_n \lambda_n = 0$, $\sum_{n=1}^{\infty} \lambda_n = \infty$, and $\sum_{n=1}^{\infty} |\lambda_{n+1} - \lambda_n| < \infty$. Then $\{x_n\}$ converges strongly to a fixed point of T .*

3. LEMMAS

In this section we prove two lemmas, which are used in Section 4.

Lemma 2. *Let t be a nonnegative real number and let $\{\beta_n\}$ be a sequence in $(0, \infty)$ converging to 0. Define sequences $\{\delta_n\}$ in $[0, \infty)$ and $\{k_n\}$ in $\mathbb{N} \cup \{0\}$ as follows:*

- $\delta_1 = t$;
- $k_n = [\delta_n / \beta_n]$ for $n \in \mathbb{N}$;
- $\delta_{n+1} = \delta_n - k_n \beta_n$ for $n \in \mathbb{N}$.

Then the following hold:

- (i) $0 \leq \delta_{n+1} < \beta_n$ for all $n \in \mathbb{N}$;
- (ii) $k_n \in \mathbb{N} \cup \{0\}$ for all $n \in \mathbb{N}$;
- (iii) $\{\delta_n\}$ converges to 0;
- (iv) $\sum_{j=1}^n k_j \beta_j + \delta_{n+1} = t$ for all $n \in \mathbb{N}$;
- (v) $\sum_{j=1}^{\infty} k_j \beta_j = t$.

Proof. We put $\varepsilon_n \in [0, 1)$ with $\delta_n / \beta_n = k_n + \varepsilon_n$ for $n \in \mathbb{N}$. We have

$$\delta_{n+1} = \delta_n - k_n \beta_n = \varepsilon_n \beta_n < \beta_n$$

for all $n \in \mathbb{N}$. From this, we also have $\delta_{n+1} = \varepsilon_n \beta_n \geq 0$. This implies (i). It is obvious that (ii) and (iii) follow from (i). Let us prove (iv) by induction. We have

$$k_1 \beta_1 + \delta_2 = k_1 \beta_1 + (\delta_1 - k_1 \beta_1) = \delta_1 = t.$$

We assume (iv) holds for some $n \in \mathbb{N}$. Then we have

$$\begin{aligned} \sum_{j=1}^{n+1} k_j \beta_j + \delta_{n+2} &= \sum_{j=1}^{n+1} k_j \beta_j + (\delta_{n+1} - k_{n+1} \beta_{n+1}) \\ &= \sum_{j=1}^n k_j \beta_j + \delta_{n+1} = t. \end{aligned}$$

So, by induction, we obtain (iv). It is obvious that (v) follows from (iii) and (iv). This completes the proof. $\square$

Lemma 3. *Let α and β be positive real numbers satisfying $\alpha/\beta \notin \mathbb{Q}$. Define sequences $\{\alpha_n\}$ in $(0, \infty)$ and $\{k_n\}$ in $\mathbb{N}$ as follows:*

- $\alpha_1 = \max\{\alpha, \beta\};$
- $\alpha_2 = \min\{\alpha, \beta\};$
- $k_n = \lfloor \alpha_n / \alpha_{n+1} \rfloor$ for all $n \in \mathbb{N};$
- $\alpha_{n+2} = \alpha_n - k_n \alpha_{n+1}$ for all $n \in \mathbb{N}.$

Then the following hold:

- (i) $0 < \alpha_{n+1} < \alpha_n$ for all $n \in \mathbb{N};$
- (ii) $k_n \in \mathbb{N}$ for all $n \in \mathbb{N};$
- (iii) $\alpha_n / \alpha_{n+1} \notin \mathbb{Q}$ for all $n \in \mathbb{N};$
- (iv) $\{\alpha_n\}$ converges to 0.

Proof. We note that (i) implies (ii). We shall prove (i), (ii) and (iii) by induction. By the assumption of $\alpha/\beta \notin \mathbb{Q}$, we have $\alpha \neq \beta$ and hence

$$\alpha_1 = \max\{\alpha, \beta\} > \min\{\alpha, \beta\} = \alpha_2 > 0.$$

It is obvious that $\alpha_1 / \alpha_2 \notin \mathbb{Q}$. We assume that $0 < \alpha_{j+1} < \alpha_j$ and $\alpha_j / \alpha_{j+1} \notin \mathbb{Q}$ for some $j \in \mathbb{N}$. Since $\alpha_{j+2} = \alpha_j - k_j \alpha_{j+1}$, we have

$$\frac{\alpha_{j+2}}{\alpha_{j+1}} = \frac{\alpha_j}{\alpha_{j+1}} - k_j \notin \mathbb{Q}$$

and hence $\alpha_{j+1} / \alpha_{j+2} \notin \mathbb{Q}$. Put $\varepsilon_j \in [0, 1)$ satisfying $\alpha_j / \alpha_{j+1} = k_j + \varepsilon_j$. Since $\alpha_j / \alpha_{j+1} \notin \mathbb{Q}$, we note that $\varepsilon_j > 0$. We have

$$\alpha_{j+2} = \alpha_j - k_j \alpha_{j+1} = \varepsilon_j \alpha_{j+1} < \alpha_{j+1}.$$

From this, we also have $\alpha_{j+2} = \varepsilon_j \alpha_{j+1} > 0$. Therefore we have shown that $0 < \alpha_{j+2} < \alpha_{j+1}$ and $\alpha_{j+1} / \alpha_{j+2} \notin \mathbb{Q}$. By induction, we obtain (i), (ii) and (iii). Let us prove (iv). Since $\{\alpha_n\}$ is a sequence of positive real numbers and strictly decreasing, $\{\alpha_n\}$ converges to some $\alpha_\infty \in [0, \infty)$. Arguing by contradiction, we assume $\alpha_\infty > 0$. Then we can choose $j \in \mathbb{N}$ such that

$$\alpha_\infty < \alpha_{j+1} < \alpha_j < 2\alpha_\infty.$$

We have

$$k_j = \left\lfloor \frac{\alpha_j}{\alpha_{j+1}} \right\rfloor = 1 \quad \text{and} \quad \alpha_{j+2} = \alpha_j - k_j \alpha_{j+1} = \alpha_j - \alpha_{j+1} < \alpha_\infty.$$

This is a contradiction. Therefore $\alpha_\infty = 0$, that is, we obtain (iv). This completes the proof. $\square$

4. MAIN RESULTS

In this section we give our main results. We first prove the following.

Proposition 1. *Let E be a Banach space and let τ be a Hausdorff topology on E . Let $\{T(t) : t \geq 0\}$ be a one-parameter τ -continuous semigroup of mappings on a subset C of E . Let $\{\alpha_n\}$ be a sequence in $[0, \infty)$ converging to $\alpha_\infty \in [0, \infty)$, and satisfying $\alpha_n \neq \alpha_\infty$ for all $n \in \mathbb{N}$. Suppose that $z \in C$ satisfies $T(\alpha_n)z = z$ for all $n \in \mathbb{N}$. Then z is a common fixed point of $\{T(t) : t \geq 0\}$.*

Proof. We note that

$$T(\alpha_\infty)z = \tau\text{-}\lim_{n \rightarrow \infty} T(\alpha_n)z = z.$$

We put $\beta_n = |\alpha_n - \alpha_\infty| > 0$ for $n \in \mathbb{N}$. By the assumption, $\{\beta_n\}$ is a sequence in $(0, \infty)$ converging to 0. Since

$$\max\{\alpha_n, \alpha_\infty\} = \min\{\alpha_n, \alpha_\infty\} + \beta_n,$$

we have

$$\begin{aligned} T(\beta_n)z &= T(\beta_n) \circ T(\min\{\alpha_n, \alpha_\infty\})z \\ &= T(\beta_n + \min\{\alpha_n, \alpha_\infty\})z = T(\max\{\alpha_n, \alpha_\infty\})z \\ &= z \end{aligned}$$

for all $n \in \mathbb{N}$. We also have

$$T(0)z = T(0) \circ T(\alpha_1)z = T(0 + \alpha_1)z = T(\alpha_1)z = z.$$

Fix $t > 0$. Then by Lemma 2, there exists a sequence $\{k_n\}$ in $\mathbb{N} \cup \{0\}$ such that

$$\sum_{n=1}^{\infty} k_n \beta_n = t.$$

For each $n \in \mathbb{N}$ with $\sum_{j=1}^n k_j \beta_j > 0$, we obtain

$$\begin{aligned} T\left(\sum_{j=1}^n k_j \beta_j\right)z &= T(\beta_n)^{k_n} \circ T(\beta_{n-1})^{k_{n-1}} \circ \dots \circ T(\beta_2)^{k_2} \circ T(\beta_1)^{k_1}z \\ &= T(\beta_n)^{k_n} \circ T(\beta_{n-1})^{k_{n-1}} \circ \dots \circ T(\beta_2)^{k_2}z \\ &= \dots = T(\beta_n)^{k_n}z \\ &= z, \end{aligned}$$

where $T(\beta_j)^0$ is the identity mapping on C . This implies

$$T(t)z = \tau\text{-}\lim_{n \rightarrow \infty} T\left(\sum_{j=1}^n k_j \beta_j\right)z = z.$$

This completes the proof. $\square$

Now, we prove one of our main results.

Proposition 2. *Let E be a Banach space and let τ be a Hausdorff topology on E . Let $\{T(t) : t \geq 0\}$ be a one-parameter τ -continuous semigroup of mappings on a subset C of E . Let α and β be positive real numbers satisfying $\alpha/\beta \notin \mathbb{Q}$. Then*

$$\bigcap_{t \geq 0} F(T(t)) = F(T(\alpha)) \cap F(T(\beta))$$

holds.

Proof. It is obvious that

$$\bigcap_{t \geq 0} F(T(t)) \subset F(T(\alpha)) \cap F(T(\beta)).$$

So we shall prove the converse inclusion. We fix $z \in F(T(\alpha)) \cap F(T(\beta))$. Define sequences $\{\alpha_n\}$ in $(0, \infty)$ and $\{k_n\}$ in $\mathbb{N}$ as in Lemma 3. By the assumption, we have

$$T(\alpha_1)z = T(\max\{\alpha, \beta\})z = z \quad \text{and} \quad T(\alpha_2)z = T(\min\{\alpha, \beta\})z = z.$$

If $T(\alpha_j)z = T(\alpha_{j+1})z = z$, then we have

$$T(\alpha_{j+2})z = T(\alpha_{j+2}) \circ T(\alpha_{j+1})^{k_j} z = T(\alpha_{j+2} + k_j \alpha_{j+1})z = T(\alpha_j)z = z.$$

So, by induction, we obtain $T(\alpha_n)z = z$ for all $n \in \mathbb{N}$. Since $\{\alpha_n\}$ is a sequence of positive real numbers converging to 0, Proposition 1 yields that z is a common fixed point of $\{T(t) : t \geq 0\}$. This completes the proof. $\square$

As a direct consequence of Proposition 2, we obtain the following.

Corollary 1. *Let E , τ , C and $\{T(t) : t \geq 0\}$ be the same as in Proposition 2. Then*

$$\bigcap_{t \geq 0} F(T(t)) = F(T(1)) \cap F(T(\sqrt{2}))$$

holds.

Using Lemma 1, we obtain the following.

Corollary 2. *Let E , τ , C , α and β be the same as in Proposition 2. Let $\{T(t) : t \geq 0\}$ be a τ -continuous semigroup of nonexpansive mappings on C . Assume that E is strictly convex, and $F(T(\alpha)) \cap F(T(\beta)) \neq \emptyset$. Then*

$$\bigcap_{t \geq 0} F(T(t)) = \{z \in C : \lambda T(\alpha)z + (1 - \lambda)T(\beta)z = z\}$$

holds for every $\lambda \in (0, 1)$.

Corollary 3. *Let E , τ , C , α and β be the same as in Proposition 2. Let $\{T(t) : t \geq 0\}$ be a τ -continuous semigroup of nonexpansive mappings on C . Assume that E is uniformly convex, and C is bounded, closed and convex. Then*

$$\bigcap_{t \geq 0} F(T(t)) = \{z \in C : \lambda T(\alpha)z + (1 - \lambda)T(\beta)z = z\}$$

holds for every $\lambda \in (0, 1)$.

5. CONVERGENCE THEOREMS

Several authors have studied convergence theorems for nonexpansive semigroups; see [1, 3, 11, 16, 18, 20, 22] and others. For example, Suzuki and Takahashi in [22] proved the following: Let C be a compact convex subset of a Banach space E and let $\{T(t) : t \geq 0\}$ be a strongly continuous semigroup of nonexpansive mappings on C . Define a sequence $\{x_n\}$ in C by $x_1 \in C$ and

$$x_{n+1} = \frac{\lambda}{t_n} \int_0^{t_n} T(s)x_n ds + (1 - \lambda)x_n$$

for $n \in \mathbb{N}$, where λ is a constant in $(0, 1)$, and $\{t_n\}$ is a sequence in $(0, \infty)$ satisfying $\lim_n t_n = \infty$ and $\lim_n t_{n+1}/t_n = 1$. Then $\{x_n\}$ converges strongly to a common fixed point of $\{T(t) : t \geq 0\}$.

Using Proposition 2, we can prove many convergence theorems for nonexpansive semigroups. In this section, we state some of them. In the following theorems, we make the assumption:

- Let E be a Banach space and let τ be a Hausdorff topology on E . Let $\{T(t) : t \geq 0\}$ be a one-parameter τ -continuous semigroup of nonexpansive mappings on a bounded closed convex subset C of E . Let α and β be positive real numbers satisfying $\alpha/\beta \notin \mathbb{Q}$.

We first state the following, which are connected with Baillon's type iteration [2]; see pages 63 and 83 in [23].

Theorem 5. *Assume that E is a Hilbert space. Let $x \in C$ and define a sequence $\{x_n\}$ in C by*

$$x_n = \frac{\sum_{k=1}^n \sum_{\ell=1}^n T(k\alpha + \ell\beta)x}{n^2}$$

for $n \in \mathbb{N}$. Then $\{x_n\}$ converges weakly to a common fixed point of $\{T(t) : t \geq 0\}$.

Proof. We note that

$$\sum_{k=1}^n \sum_{\ell=1}^n T(k\alpha + \ell\beta)x = \sum_{k=1}^n \sum_{\ell=1}^n T(\alpha)^k \circ T(\beta)^\ell x.$$

Thus, $\{x_n\}$ converges weakly to a common fixed point z of $T(\alpha)$ and $T(\beta)$. Such z is a common fixed point of $\{T(t) : t \geq 0\}$ by Proposition 2. This completes the proof. $\square$

Theorem 6. *Assume that E is a Hilbert space. Let $x \in C$ and define a sequence $\{x_n\}$ in C by*

$$x_n = \frac{\sum_{k=1}^n \left(\frac{T(\alpha) + T(\beta)}{2} \right)^k x}{n}$$

for $n \in \mathbb{N}$. Then $\{x_n\}$ converges weakly to a common fixed point of $\{T(t) : t \geq 0\}$.

Proof. By Theorem 1, $\{x_n\}$ converges weakly to some $z \in C$ which is a fixed point of $(T(\alpha) + T(\beta))/2$. By Corollary 3, we obtain that z is a common fixed point of $\{T(t) : t \geq 0\}$. This completes the proof. $\square$

We next state the following, which are connected with Krasnosel'skiĭ-Mann's type iteration [14, 15]; see Reich [17] and Suzuki [19, 21].

Theorem 7. *Assume that E is a uniformly convex Banach space whose norm is Fréchet differentiable. Fix $\kappa, \lambda > 0$ with $\kappa + \lambda < 1$. Define a sequence $\{x_n\}$ in C by $x_1 \in C$ and*

$$x_{n+1} = \kappa T(\alpha)x_n + \lambda T(\beta)x_n + (1 - \kappa - \lambda)x_n,$$

for $n \in \mathbb{N}$. Then $\{x_n\}$ converges weakly to a common fixed point of $\{T(t) : t \geq 0\}$.

Proof. By Theorem 2, $\{x_n\}$ converges weakly to some $z \in C$ which is a fixed point of

$$\frac{\kappa}{\kappa + \lambda} T(\alpha) + \frac{\lambda}{\kappa + \lambda} T(\beta).$$

So, by Corollary 3, we obtain that z is a common fixed point of $\{T(t) : t \geq 0\}$. This completes the proof. $\square$

Theorem 8. Assume that C is compact. Fix $\lambda \in (0, 1)$. Define a sequence $\{x_n\}$ in C by $x_1 \in C$ and

$$x_{n+1} = \lambda \frac{\sum_{k=1}^n \sum_{\ell=1}^n T(k\alpha + \ell\beta)x_n}{n^2} + (1 - \lambda)x_n,$$

for $n \in \mathbb{N}$. Then $\{x_n\}$ converges strongly to a common fixed point of $\{T(t) : t \geq 0\}$.

Using Ishikawa's result in [12], we obtain the following.

Theorem 9. Assume that C is compact. Fix $\kappa, \lambda \in (0, 1)$. Define a sequence $\{x_n\}$ in C by $x_1 \in C$ and

$$x_{n+1} = (\lambda T(\alpha) + (1 - \lambda)I) \circ (\kappa T(\beta) + (1 - \kappa)I)^n x_n$$

for $n \in \mathbb{N}$, where I is the identity mapping on C . Then $\{x_n\}$ converges strongly to a common fixed point of $\{T(t) : t \geq 0\}$.

We next state the following, which is connected with Browder's type implicit iteration [6]. We note that $x \mapsto (1 - \lambda)Tx + \lambda u$ is a contractive mapping if T is a nonexpansive mapping and $\lambda \in (0, 1)$. Therefore, the Banach contraction principle [4] yields that such a mapping has a unique fixed point.

Theorem 10. Assume that E is a Hilbert space. Fix $u \in C$ and define a sequence $\{x_n\}$ in C by

$$x_n = \frac{1 - \lambda_n}{2} T(\alpha)x_n + \frac{1 - \lambda_n}{2} T(\beta)x_n + \lambda_n u$$

for $n \in \mathbb{N}$, where $\{\lambda_n\}$ is a sequence in $(0, 1)$ converging to 0. Then $\{x_n\}$ converges strongly to a common fixed point of $\{T(t) : t \geq 0\}$.

We finally state the following, which is connected with Halpern's type explicit iteration [10]; see Wittmann [24].

Theorem 11. Assume that E is a Hilbert space. Fix $u \in C$ and define a sequence $\{x_n\}$ in C by $x_1 \in C$ and

$$x_{n+1} = \frac{1 - \lambda_n}{2} T(\alpha)x_n + \frac{1 - \lambda_n}{2} T(\beta)x_n + \lambda_n u$$

for $n \in \mathbb{N}$, where $\{\lambda_n\}$ is a sequence in $[0, 1]$ satisfying the following:

$$\lim_{n \rightarrow \infty} \lambda_n = 0; \quad \sum_{n=1}^{\infty} \lambda_n = \infty; \quad \text{and} \quad \sum_{n=1}^{\infty} |\lambda_{n+1} - \lambda_n| < \infty.$$

Then $\{x_n\}$ converges strongly to a common fixed point of $\{T(t) : t \geq 0\}$.

REFERENCES

- [1] S. Atsushiba and W. Takahashi, "Strong convergence theorems for one-parameter nonexpansive semigroups with compact domains", in *Fixed Point Theory and Applications*, Volume 3 (Y. J. Cho, J. K. Kim and S. M. Kang Eds.), pp. 15–31, Nova Science Publishers, New York, 2002. MR2081406 (2005i:47081)
- [2] J. B. Baillon, "Un théorème de type ergodique pour les contractions non linéaires dans un espace de Hilbert", *C. R. Acad. Sci. Paris, Sér. A-B*, **280** (1975), 1511–1514. MR0375009 (51:11205)
- [3] J. B. Baillon, "Quelques propriétés de convergence asymptotique pour les semigroupes de contractions impaires", *C. R. Acad. Sci. Paris*, **283** (1976), A75–A78. MR0425702 (54:13655)

- [4] S. Banach, “*Sur les opérations dans les ensembles abstraits et leur application aux équations intégrales*”, Fund. Math., **3** (1922), 133–181.
- [5] F. E. Browder, “*Nonexpansive nonlinear operators in a Banach space*”, Proc. Nat. Acad. Sci. USA, **54** (1965), 1041–1044. MR0187120 (32:4574)
- [6] F. E. Browder, “*Convergence of approximants to fixed points of nonexpansive nonlinear mappings in Banach spaces*”, Arch. Ration. Mech. Anal., **24** (1967), 82–90. MR0206765 (34:6582)
- [7] R. E. Bruck, “*Properties of fixed-point sets of nonexpansive mappings in Banach spaces*”, Trans. Amer. Math. Soc., **179** (1973), 251–262. MR0324491 (48:2843)
- [8] R. E. Bruck, “*A common fixed point theorem for a commuting family of nonexpansive mappings*”, Pacific J. Math., **53** (1974), 59–71. MR0361945 (50:14387)
- [9] D. Göhde, “*Zum Prinzip der kontraktiven Abbildung*”, Math. Nachr., **30** (1965), 251–258. MR0190718 (32:8129)
- [10] B. Halpern, “*Fixed points of nonexpanding maps*”, Bull. Amer. Math. Soc., **73** (1967), 957–961. MR0218938 (36:2022)
- [11] N. Hirano, “*Nonlinear ergodic theorems and weak convergence theorems*”, J. Math. Soc. Japan, **34** (1982), 35–46. MR0639803 (83d:47059)
- [12] S. Ishikawa, “*Common fixed points and iteration of commuting nonexpansive mappings*”, Pacific J. Math., **80** (1979), 493–501. MR0539430 (80g:47065)
- [13] W. A. Kirk, “*A fixed point theorem for mappings which do not increase distances*”, Amer. Math. Monthly, **72** (1965), 1004–1006. MR0189009 (32:6436)
- [14] M. A. Krasnosel’skiĭ, “*Two remarks on the method of successive approximations*” (in Russian), Uspehi Mat. Nauk **10** (1955), 123–127. MR0068119 (16:833a)
- [15] W. R. Mann, “*Mean value methods in iteration*”, Proc. Amer. Math. Soc., **4** (1953), 506–510. MR0054846 (14:988f)
- [16] I. Miyadera and K. Kobayasi, “*On the asymptotic behaviour of almost-orbits of nonlinear contraction semigroups in Banach spaces*”, Nonlinear Anal., **6** (1982), 349–365. MR0654811 (84c:47064)
- [17] S. Reich, “*Weak convergence theorems for nonexpansive mappings in Banach spaces*”, J. Math. Anal. Appl., **67** (1979), 274–276. MR0528688 (80d:47090)
- [18] N. Shioji and W. Takahashi, “*Strong convergence theorems for asymptotically nonexpansive semigroups in Hilbert spaces*”, Nonlinear Anal., **34** (1998), 87–99. MR1631657 (99e:47071)
- [19] T. Suzuki, “*Strong convergence theorem to common fixed points of two nonexpansive mappings in general Banach spaces*”, J. Nonlinear Convex Anal., **3** (2002), 381–391. MR1947106 (2003i:47066)
- [20] T. Suzuki, “*On strong convergence to common fixed points of nonexpansive semigroups in Hilbert spaces*”, Proc. Amer. Math. Soc., **131** (2003), 2133–2136. MR1963759 (2004b:47108)
- [21] T. Suzuki, “*Common fixed points of two nonexpansive mappings in Banach spaces*”, Bull. Austral. Math. Soc., **69** (2004), 1–18. MR2040045 (2004m:47146)
- [22] T. Suzuki and W. Takahashi, “*Strong convergence of Mann’s type sequences for one-parameter nonexpansive semigroups in general Banach spaces*”, J. Nonlinear Convex Anal., **5** (2004), 209–216. MR2083912 (2005c:47093)
- [23] W. Takahashi, “*Nonlinear Functional Analysis*”, Yokohama Publishers, Yokohama, 2000. MR1864294 (2002k:47001)
- [24] R. Wittmann, “*Approximation of fixed points of nonexpansive mappings*”, Arch. Math. (Basel), **58** (1992), 486–491. MR1156581 (93c:47069)

DEPARTMENT OF MATHEMATICS, KYUSHU INSTITUTE OF TECHNOLOGY, SENSUICHO, TOBATA, KITAKYUSHU 804-8550, JAPAN

E-mail address: `suzuki-t@mns.kyutech.ac.jp`