A CHARACTERISATION OF $0^\#$ IN TERMS OF FORCING

SY D. FRIEDMAN

(Communicated by Carl G. Jockusch, Jr.)

ABSTRACT. We show that “saturation” of the universe with respect to forcing over L with partial orders on ω_1 is equivalent to the existence of $0^\#$.

If P is a constructible forcing notion, then we say that $G \subseteq P$ is P -generic iff G is P -generic over L . The statement that all countable constructible forcings have generics is rather weak, and holds for example in $L[R]$, where R is a Cohen real over L . But it is not possible that all constructible forcings have generics: consider the forcing that collapses ω_1 to ω with finite conditions.

Definition. V is L -saturated for ω_1 -forcings iff whenever P is a constructible forcing of L -cardinality ω_1 such that for any $p \in P$ there is a P -generic containing p in some ω_1 -preserving extension of V , then there is a P -generic in V .

Theorem 1. *The following are equivalent:*

- (a) V is L -saturated for ω_1 -forcings.
- (b) $0^\#$ exists.

Proof. (a) $\rightarrow$ (b) The existence of $0^\#$ is equivalent to the statement that every stationary constructible subset of ω_1 contains a CUB subset (see [2]). Now use the following:

Fact (Baumgartner; see [1]). If X is a stationary constructible subset of ω_1 , then there is a forcing $P \in L$ of L -cardinality ω_1 which preserves cardinals over V and adds a CUB subset to X . (P adds a CUB subset of X using “finite conditions”.)

(b) $\rightarrow$ (a) Assume that $0^\#$ exists, and suppose that P is a constructible forcing of L -cardinality ω_1 such that every condition in P belongs to a generic in an ω_1 -preserving extension of V . We will show that there is a P -generic in V . Assume that the universe of P is exactly ω_1 . Let P be of the form $t(\vec{i}, \omega_1, \vec{\omega})$, where $\vec{i} < \omega_1 < \vec{\omega}$ is a finite increasing sequence of indiscernibles and t is an L -term. We claim that if $\vec{i} < k_0 < k_1$ are countable indiscernibles and G_{k_0} is P_{k_0} -generic over L , then there is G_{k_1} containing G_{k_0} which is P_{k_1} -generic over L , where $P_k = t(\vec{i}, k, \vec{\omega})$. If not, then player I wins the open game $\mathcal{G}(k_0, k_1, G_{k_0})$, where I chooses constructible dense subsets of P_{k_1} and II responds with increasingly strong conditions meeting

Received by the editors January 4, 2005.

2000 *Mathematics Subject Classification.* Primary 03E40, 03E45, 03E55.

Key words and phrases. Forcing, Silver indiscernibles, constructibility.

This research was supported by Research Project Number P16334-NO5 of the Austrian Foundation for the Support of Scientific Research (FWF). The author thanks David Asperò for posing the question answered here.

©2005 American Mathematical Society
 Reverts to public domain 28 years from publication

these dense sets which are compatible with all conditions in G_{k_0} . The latter is a property of the model $L[G_{k_0}]$. Let $p \in P_{k_0}$ be a condition forcing that I wins $\mathcal{G}(k_0, k_1, G_{k_0})$. Then p forces that I wins $\mathcal{G}(k_2, k_3, G_{k_2})$, where $k_2 < k_3$ are any indiscernibles $\geq k_0$ and G_{k_2} denotes the P_{k_2} -generic. But now let G be a P -generic containing p in an ω_1 -preserving extension of V . As G preserves ω_1 over V , there are indiscernibles $k_2 < k_3$ with $k_0 \leq k_2$ such that $G \cap k_2$ is P_{k_2} -generic and $G \cap P_{k_3}$ is P_{k_3} -generic, so clearly player II has a winning strategy in the game $\mathcal{G}(k_2, k_3, G \cap P_{k_2})$, in contradiction to the choice of p .

Now it is easy to build a P -generic: List the countable indiscernibles greater than $\vec{i}$ as $j_0 < j_1 < j_2 < \dots$ and inductively choose P_{j_α} -generic G_α such that $\alpha < \beta$ implies $G_\alpha \subseteq G_\beta$. At the first step, G_{j_0} is an arbitrary P_{j_0} -generic. By the previous paragraph there is no difficulty at the successor steps, where one extends G_{j_α} to $G_{j_{\alpha+1}}$. At limit stages λ , the P_{j_λ} -genericity of the union G_{j_λ} of the G_{j_α} , $\alpha < \lambda$, follows by indiscernibility. The desired P -generic is the union of the G_{j_α} , $\alpha < \omega_1$. $\square$

Remark. The proof of (a) implies (b) shows that the theorem still holds if “ ω_1 -preserving extension” is taken to be “ ω_1 -preserving set-generic extension” in the definition of L -saturation for ω_1 -forcings.

Question. Suppose that $0^\#$ exists. Then does part (a) of the theorem hold (in the obvious sense) for constructible ω_1^{+L} -forcings, i.e. constructible P of L -cardinality ω_1^+ of L ?

REFERENCES

- [1] Abraham, U. and Shelah, S. Forcing closed unbounded sets, J. Symbolic Logic 48 (1983) 643-657. MR0716625 (85i:03112)
- [2] Jech, T. *Set Theory: Millenium Edition*, Springer-Verlag, 2003. MR1940513 (2004g:03071)

KURT GÖDEL RESEARCH CENTER FOR MATHEMATICAL LOGIC, WÄHRINGERSTRASSE 25, A-1090 WIEN, AUSTRIA

E-mail address: sdf@logic.univie.ac.at