ON THE INVARIANT TRANSLATION APPROXIMATION PROPERTY FOR DISCRETE GROUPS

JOACHIM ZACHARIAS

(Communicated by David R. Larson)

ABSTRACT. Recently J. Roe considered the question of whether for a discrete group the reduced group C^* -algebra $C_r^*(\Gamma)$ is the fixed point algebra of $\{\text{Ad}(\rho_t) \mid t \in \Gamma\}$ acting on the uniform Roe algebra $UC_r^*(\Gamma)$. Γ is said to have the invariant translation approximation property in this case. We consider a slight generalization of this property which, for exact Γ , is equivalent to the operator space approximation property of $C_r^*(\Gamma)$. We also give a new characterization of exactness and a short proof of the equivalence of exactness of Γ and exactness of $C_r^*(\Gamma)$ for discrete groups.

1. INTRODUCTION

A discrete group Γ has a natural coarse structure which allows us to define the uniform Roe algebra $UC_r^*(\Gamma)$. This algebra may be thought of as the closure of scalar $\Gamma \times \Gamma$ -matrices $[\alpha_{s,t}]$ of finite width (i.e. $\{st^{-1} \mid \alpha_{s,t} \neq 0\}$ is finite) with uniformly bounded entries acting on $\ell^2(\Gamma)$. The reduced group C^* -algebra $C_r^*(\Gamma)$ is naturally contained in $UC_r^*(\Gamma)$. Indeed the left translation λ_r on $\ell^2(\Gamma)$ is given by the matrix where $\alpha_{s,t} = 1$ if $st^{-1} = r$ and $\alpha_{s,t} = 0$ otherwise; so the matrices $[\alpha_{s,t}]$ of finite width such that $\alpha_{sr,tr} = \alpha_{s,t}$ for all $s, t, r \in \Gamma$ form precisely the group ring $\mathbb{C}[\Gamma]$. They may also be characterized as those finite width matrices fixed by all automorphisms of the form $\text{Ad}(\rho_t)$, where $t \in \Gamma$ and ρ is the right regular representation of Γ .

Now consider the set of fixed points $UC_r^*(\Gamma)^\Gamma$ of this action in the whole of $UC_r^*(\Gamma)$, not just in the dense subalgebra of finite width matrices. According to [Ro] Γ is said to have the invariant translation approximation property if $C_r^*(\Gamma) = UC_r^*(\Gamma)^\Gamma$. One might wonder whether all discrete groups have this property.

In this note we do not answer this question but consider a slightly stronger invariance property defined as follows. One may define $UC_r^*(\Gamma, S)$ with coefficients in a (concrete) operator space S . $UC_r^*(\Gamma, S)$ is an operator space in general and is a C^* -algebra if S is a C^* -algebra. This very natural idea to allow nontrivial coefficients in the definition of the uniform Roe algebra might be useful elsewhere. $UC_r^*(\Gamma, -)$ may be regarded as a functor (on C^* -algebras or on operator spaces). In Theorem 2.3 we show that Γ is exact iff this functor is exact on C^* -algebras and

Received by the editors March 22, 2004 and, in revised form, February 1, 2005.

2000 *Mathematics Subject Classification*. Primary 46L06, 46L85, 20F69.

Key words and phrases. Exact groups, uniform Roe algebra, invariant translation approximation property, operator approximation property.

©2006 American Mathematical Society
 Reverts to public domain 28 years from publication

give a quick proof of the result of Kirchberg and Wassermann on the equivalence of exactness of Γ and exactness of $C_r^*(\Gamma)$ for discrete groups ([KW]) together with another characterization of exactness.

Instead of looking at the fixed points in $UC_r^*(\Gamma)$ we consider fixed points in $UC_r^*(\Gamma, S)$. Using results and techniques from [HK] we show that for exact Γ the following conditions are equivalent:

- (1) $UC_r^*(\Gamma, S)^\Gamma = C_r^*(\Gamma) \otimes S$ for all closed subspaces S of the compact operators $\mathcal{K}$ (on $\ell^2(\mathbb{N})$).
- (2) $(UC_r^*(\Gamma) \otimes S)^\Gamma = C_r^*(\Gamma) \otimes S$ for all closed subspaces S of $\mathcal{K}$.
- (3) For any operator space S (not necessarily contained in $\mathcal{K}$) $UC_r^*(\Gamma, S)^\Gamma = (UC_r^*(\Gamma) \otimes S)^\Gamma = C_r^*(\Gamma) \otimes S$.
- (4) $C_r^*(\Gamma)$ has the operator approximation property (OAP) or equivalently Γ has the AP of [HK].

Here and throughout $\otimes$ denotes the minimal tensor product. For group C^* -algebras of discrete groups the OAP implies exactness, and so condition (4) implies the other conditions for all discrete groups Γ . Moreover, the invariant translation approximation property with coefficients implies the one without coefficients, so if Γ has the AP, then $C_r^*(\Gamma) = UC_r^*(\Gamma)^\Gamma$. We also remark that it is not known whether there are exact groups without the AP, but it has been conjectured that $SL_3(\mathbb{Z})$ is such a group (cf. [HK] or [ER]).

2. PRELIMINARIES

A C^* -algebra A is *exact* if given any exact sequence $0 \rightarrow J \rightarrow B \rightarrow C \rightarrow 0$ of C^* -algebras the sequence $0 \rightarrow A \otimes J \rightarrow A \otimes B \rightarrow A \otimes C \rightarrow 0$ is again exact, i.e. $A \otimes -$ is an exact functor. Inspection shows that for arbitrary A the kernel of the map $A \otimes B \rightarrow A \otimes C$ is the *Fubini product* $F(A, J) = \{x \in A \otimes B \mid (\varphi \otimes \text{id})(x) \in J \forall \varphi \in A^*\}$ (cf. [Wa]). Thus exactness of A is equivalent to requiring $F(A, J) = A \otimes J$ for all pairs $J \subseteq B$, where B is a C^* -algebra and J a closed ideal of B .

Kraus [Kr] considered the same condition for subspaces, i.e. $F(A, S) = A \otimes S$ for all pairs $S \subseteq B$, where B is a C^* -algebra and S a closed subspace of B . (Wassermann had considered this earlier for subalgebras.) He showed (cf. [Kr], [ER]) that this condition, the *slice map property for subspaces*, is equivalent to the *strong operator approximation property (strong OAP)* which means the following: for every C^* -algebra B there is a net (ϕ_α) of finite rank maps such that $\phi_\alpha \otimes \text{id}(x) \rightarrow x$ for all $x \in A \otimes B$. In particular if A has the strong OAP, then it is exact. A weaker variant is the *operator approximation property (OAP)* which requires only a net (ϕ_α) of finite rank maps such that $\phi_\alpha \otimes \text{id}(x) \rightarrow x$ for all $x \in A \otimes \mathcal{K}$ ($\mathcal{K} = \mathcal{K}(\ell^2(\mathbb{N}))$). It is the operator space version of Grothendieck's AP (cf. [ER]). The OAP is equivalent to the slice map property for closed subspaces of $\mathcal{K}$.

Most regularity properties of C^* -algebras define, when required for $C_r^*(\Gamma)$, reasonable regularity properties for the group Γ , e.g. nuclearity of $C_r^*(\Gamma)$ is equivalent to amenability of Γ . Similarly $C_r^*(\Gamma)$ is an exact C^* -algebra iff Γ is amenable at infinity. Another formally stronger property is exactness of Γ defined as follows ([KW]). Let $\alpha : \Gamma \rightarrow \text{Aut}(A)$ be a Γ -action on a C^* -algebra A such that $\alpha_s(J) = J$ for all $s \in \Gamma$; then $\dot{\alpha}_s(a + J) = \alpha_s(a) + J$ defines an action $\dot{\alpha}$ on $B = A/J$. Denoting the restriction of α to J by α again, we obtain the exact sequence of Γ -dynamical

systems $0 \rightarrow (J, \alpha) \rightarrow (A, \alpha) \rightarrow (B, \dot{\alpha}) \rightarrow 0$. Γ is said to be *exact* if the corresponding sequence of reduced crossed products $0 \rightarrow J \rtimes_{\alpha, r} \Gamma \rightarrow A \rtimes_{\alpha, r} \Gamma \rightarrow B \rtimes_{\dot{\alpha}, r} \Gamma \rightarrow 0$ is exact. A discrete group Γ is exact iff $C_r^*(\Gamma)$ is an exact C^* -algebra ([KW]), and we will give a short proof of this fact in Theorem 2.3.

As for the strong OAP and the OAP, Haagerup and Kraus [HK] show that they are equivalent for $C_r^*(\Gamma)$ if Γ is a discrete group. They also establish an equivalent condition in terms of the Fourier algebra $A(\Gamma)$ which they call the approximation property (AP). The Fourier algebra $A(\Gamma)$ is the set of all coefficients $s \mapsto \langle \xi, \lambda_s \eta \rangle$ in the regular representation of Γ with the pointwise product and the norm $\|u\| = \inf\{\|\xi\| \|\eta\| \mid u(s) = \langle \xi, \lambda_s \eta \rangle \text{ for all } s \in \Gamma\}$. Amenability of Γ is equivalent to the existence of a bounded approximate unit in $A(\Gamma)$ (even one consisting of positive definite functions). In order to explain the AP recall that a complex function u on Γ is a *multiplier* of $A(\Gamma)$ if $uv \in A(\Gamma)$ for all $v \in A(\Gamma)$. In $A(\Gamma)$ convergence in norm implies pointwise convergence, and therefore, by the closed graph theorem, such a multiplier u defines a bounded linear map on $A(\Gamma)$. Since $A(\Gamma) = VN(\Gamma)_*$, by duality, it also defines a bounded linear $\overline{M}_u \in B(VN(\Gamma))$, and since $\overline{M}_u$ leaves the group ring $\mathbb{C}[\Gamma]$ invariant we finally obtain a bounded $M_u \in B(C_r^*(\Gamma))$. u is said to be a completely bounded multiplier if M_u is a completely bounded map, i.e. $M_u \in CB(C_r^*(\Gamma))$. The space $M_0A(\Gamma)$ of completely bounded multipliers can then be equipped with the completely bounded norm $\|u\|_{M_0}$. As cited in [HK] (cf. [Pi]) $u \in M_0A(\Gamma)$ iff there is a Hilbert space K and bounded maps $p, q : \Gamma \rightarrow K$ such that $u(st^{-1}) = \langle p(s), q(t) \rangle$ for all $s, t \in \Gamma$. Moreover, the completely bounded norm is given by

$$\|u\|_{M_0} = \inf\{\|p\|_\infty \|q\|_\infty \mid \exists K \text{ s.t. } p, q : \Gamma \rightarrow K, u(st^{-1}) = \langle p(s), q(t) \rangle \forall s, t \in \Gamma\}.$$

Since all $u \in M_0A(\Gamma)$ are bounded functions, every summable sequence of scalars $(\alpha_s) \in \ell^1(\Gamma)$ defines a functional on $M_0A(\Gamma)$ by $\omega_{(\alpha_s)}(u) = \sum_s \alpha_s u(s)$. Let $Q(\Gamma)$ be the closure of $\{\omega_{(\alpha_s)} \mid (\alpha_s) \in \ell^1(\Gamma)\}$ in $M_0A(\Gamma)^*$. Then Γ is said to have the *approximation property* (AP) if there is a net $(u_\alpha) \subseteq A(\Gamma)$ converging to 1 in the $\sigma(M_0A(\Gamma), Q(\Gamma))$ -topology. By ([HK], 2.1) $C_r^*(\Gamma)$ has the OAP iff Γ has the AP. It is pointed out in [HK] that one may assume $u_\alpha \in A_c(\Gamma)$, i.e. all the u_α to have finite support.

A matrix $[\alpha_{s,t}]_{s,t \in \Gamma}$ is said to have *width* F , where $F \subseteq \Gamma$ is a subset if $\alpha_{s,t} \neq 0$ implies $st^{-1} \in F$. If $[\alpha_{s,t}]_{s,t \in \Gamma}$ has finite width (i.e. width F for some finite $F \subseteq \Gamma$) and there is $M \geq 0$ such that $|\alpha_{s,t}| \leq M$ for all $s, t \in \Gamma$, then this matrix defines a bounded operator acting on $\ell^2(\Gamma)$. Such matrices form a $*$ -algebra and the uniform Roe algebra $UC_r^*(\Gamma)$ is the closure of it in $B(\ell^2(\Gamma))$. We have $UC_r^*(\Gamma) = \ell^\infty(\Gamma) \rtimes_{\tau, r} \Gamma$, where τ is the translation action on $\ell^\infty(\Gamma)$. $UC_r^*(\Gamma)$ is closely related to exactness of Γ . More precisely Guentner, Kaminker ([GK]) and Ozawa ([Oz]) showed the following important result (cf. [Pi] for an excellent exposition).

Theorem 2.1. *For a discrete group Γ the following conditions are equivalent:*

- (1) $C_r^*(\Gamma)$ is exact.
- (2) $UC_r^*(\Gamma)$ is nuclear.
- (3) *There exists a net of finite width positive definite kernels $k_\alpha : \Gamma \times \Gamma \rightarrow \mathbb{C}$ such that for all $\varepsilon > 0$ and every finite subset $F \subseteq \Gamma$ there is α_0 such that $|k_\alpha(s, t) - 1| < \varepsilon$ whenever $st^{-1} \in F$ and $\alpha \geq \alpha_0$.*

In (3) k_α positive definite means $\sum_{s,t} k_\alpha(s, t) \bar{\alpha}_s \alpha_t \geq 0$ for all $(\alpha_s)_{s \in \Gamma}$ of finite support. Condition (3) is quite similar to classical amenability which is equivalent

to the existence of a net of positive definite functions on Γ with analogous properties. Another equivalent condition says that Γ is amenable at infinity, i.e. acts amenably on a compact space.

By a GNS-type construction attributed to Kolmogoroff any positive definite kernel $k : \Gamma \times \Gamma \rightarrow \mathbb{C}$ is of the form $k(s, t) = \langle v(s), v(t) \rangle$, where $v : \Gamma \rightarrow K$ is a map into a suitable Hilbert space K . Thus $V(e_s) = e_s \otimes v(s)$ defines a linear map $V : \ell^2(\Gamma) \rightarrow \ell^2(\Gamma) \otimes K$ such that $\|V^*V\| = \sup\{\|v(s)\|^2 \mid s \in \Gamma\}$, and if $X \in B(\ell^2(\Gamma))$ has the matrix representation $[\alpha_{s,t}]$, then $V^*(X \otimes 1)V$ has the matrix representation $[k(s, t)\alpha_{s,t}]$. Thus $\Theta_k([\alpha_{s,t}]) = [k(s, t)\alpha_{s,t}]$ defines a completely positive map (Schur multiplier) $\Theta_k \in B(B(\ell^2(\Gamma)))$ which is used in [Oz]. If k has finite width, then $\Theta_k(B(\ell^2(\Gamma)))$ consists of finite width matrices. Moreover, if (k_α) is as in Theorem 2.1(3), then $\Theta_{k_\alpha}(x) \rightarrow x$ for $x \in UC_r^*(\Gamma)$ since this is true for finite width matrices and $\|\Theta_{k_\alpha}\| \rightarrow 1$ as $\alpha \rightarrow \infty$.

If $S \subseteq B(H)$ is a closed subspace (i.e. a concrete operator space) we may define the operator space $UC_r^*(\Gamma, S)$ as the closure of finite width matrices $[a_{s,t}]_{s,t \in \Gamma}$, where $a_{s,t} \in S$ and $\|a_{s,t}\|$ is uniformly bounded for all $s, t \in \Gamma$ acting on $\ell^2(\Gamma) \otimes H$. Clearly $UC_r^*(\Gamma) \otimes S \subseteq UC_r^*(\Gamma, S)$. Similar to tensor product functors like $C_r^*(\Gamma) \otimes -$, we may regard $UC_r^*(\Gamma, -)$ as a functor on the category of C^* -algebras and $*$ -homomorphisms, and will say that $UC_r^*(\Gamma, -)$ is an exact functor if for every exact sequence $0 \rightarrow J \rightarrow A \rightarrow B \rightarrow 0$ of C^* -algebras the induced sequence $0 \rightarrow UC_r^*(\Gamma, J) \rightarrow UC_r^*(\Gamma, A) \rightarrow UC_r^*(\Gamma, B) \rightarrow 0$ is exact. In Theorem 2.3 we will see that exactness of Γ and of $UC_r^*(\Gamma, -)$ are equivalent. The proof uses Theorem 2.1 and ideas from [KW] and [Ki]. We start with some preliminaries.

Let $0 \rightarrow (J, \alpha) \rightarrow (A, \alpha) \xrightarrow{p} (B, \dot{\alpha}) \rightarrow 0$ be an exact sequence of Γ -dynamical systems. Denote by p_α the surjective $*$ -homomorphism $p_\alpha : A \rtimes_{\alpha, r} \Gamma \rightarrow B \rtimes_{\dot{\alpha}, r} \Gamma$ induced by p . $A \rtimes_{\alpha, r} \Gamma$ may be regarded naturally as a subalgebra of $UC_r^*(\Gamma, A)$ since for $a \in A$ and $s \in \Gamma$ the element $a\lambda_s \in A \rtimes_{\alpha, r} \Gamma$ corresponds to the matrix $\sum_{t \in \Gamma} e_{t, s^{-1}t} \otimes \alpha_{t^{-1}}(a) \in UC_r^*(\Gamma, A)$. Let $(e_\lambda)_{\lambda \in \Lambda}$ be an approximate unit in J consisting of positive contractions. Then $\hat{e}_\lambda = \sum_{t \in \Gamma} e_{t, t} \otimes \alpha_{t^{-1}}(e_\lambda)$ corresponds to the element e_λ in the regular representation and is an approximate unit for $J \rtimes_{\alpha, r} \Gamma$. Since $\hat{e}_\lambda(A \rtimes_{\alpha, r} \Gamma)\hat{e}_\lambda \subseteq J \rtimes_{\alpha, r} \Gamma$ and $\hat{e}_\lambda(a\lambda_s)\hat{e}_\lambda \rightarrow a\lambda_s$ for all $a \in J$ and $s \in \Gamma$, it follows that $x \in A \rtimes_{\alpha, r} \Gamma$ lies in $J \rtimes_{\alpha, r} \Gamma$ iff $\hat{e}_\lambda x \hat{e}_\lambda \rightarrow x$ as $\lambda \rightarrow \infty$.

Let $\ell^\infty(\Lambda, A)$ be the bounded functions on Λ with values in A and $c_0(\Lambda, A) = \{(a_\lambda) \in \ell^\infty(\Lambda, A) \mid a_\lambda \rightarrow 0 \text{ as } \lambda \rightarrow \infty\}$. Denoting the quotient C^* -algebra $\ell^\infty(\Lambda, A)/c_0(\Lambda, A)$ by $q^\infty(\Lambda, A)$ we obtain the exact sequence

$$0 \longrightarrow c_0(\Lambda, A) \longrightarrow \ell^\infty(\Lambda, A) \xrightarrow{q} q^\infty(\Lambda, A) \longrightarrow 0$$

and the induced surjective $*$ -homomorphism

$$\hat{q} : UC_r^*(\Gamma, \ell^\infty(\Lambda, A)) \longrightarrow UC_r^*(\Gamma, q^\infty(\Lambda, A)).$$

Following ideas in [Ki] we define the completely bounded map

$$\delta : A \rtimes_{\alpha, r} \Gamma \longrightarrow UC_r^*(\Gamma, \ell^\infty(\Lambda, A))$$

by $\delta(x)_\lambda = x - \hat{e}_\lambda x \hat{e}_\lambda$, where we think of $UC_r^*(\Gamma, \ell^\infty(\Lambda, A)) \subseteq \ell^\infty(\Lambda, UC_r^*(\Gamma, A))$ as the closure of families (x_λ) , where all $x_\lambda \in UC_r^*(\Gamma, A)$ have the same finite width. Then it is clear that $\delta(a\lambda_s) \in UC_r^*(\Gamma, \ell^\infty(\Lambda, A))$ for all $a \in A$ and $s \in \Gamma$, and so $\delta(A \rtimes_{\alpha, r} \Gamma) \subseteq UC_r^*(\Gamma, \ell^\infty(\Lambda, A))$.

Lemma 2.2. *For $x \in A \rtimes_{\alpha,r} \Gamma$ consider the following conditions:*

- (1) $x \in J \rtimes_{\alpha,r} \Gamma$.
- (2) $\delta(x) \in UC_r^*(\Gamma, c_0(\Lambda, A))$.
- (3) $x \in \ker(p_\alpha : A \rtimes_{\alpha,r} \Gamma \rightarrow B \rtimes_{\dot{\alpha},r} \Gamma)$.
- (4) $x_{s,t} \in J$ for all $s, t \in \Gamma$, where $x_{s,t}$ denote the matrix elements of x .
- (5) $\delta(x) \in \ker(\hat{q} : UC_r^*(\Gamma, \ell^\infty(\Lambda, A)) \rightarrow UC_r^*(\Gamma, q^\infty(\Lambda, A)))$.

Then (1) $\Leftrightarrow$ (2) and (3) $\Leftrightarrow$ (4) $\Leftrightarrow$ (5).

Proof. (1) $\Leftrightarrow$ (2): Follows from the fact that $\hat{e}_\lambda x \hat{e}_\lambda \xrightarrow{\lambda \rightarrow \infty} x$ iff $x \in J \rtimes_{\alpha,r} \Gamma$.

(3) $\Leftrightarrow$ (4): This is evident if we note that $p_\alpha : A \rtimes_{\alpha,r} \Gamma \rightarrow B \rtimes_{\dot{\alpha},r} \Gamma$ is simply the restriction of $\hat{p} : UC_r^*(\Gamma, A) \rightarrow UC_r^*(\Gamma, B)$ and the kernel of $\hat{p}$ consists precisely of those matrices with entries in J .

(4) $\Leftrightarrow$ (5): We have

$$\ker \hat{q} = \{X \in UC_r^*(\Gamma, \ell^\infty(\Lambda, A)) \mid X_{s,t} \in c_0(\Lambda, A) \text{ for all } s, t \in \Gamma\}.$$

Regarding $\delta(x)$ as in its definition as a family $(\delta(x)_\lambda)_{\lambda \in \Lambda}$, we have

$$\delta(x)_{\lambda;s,t} = x_{s,t} - \alpha_{s^{-1}}(e_\lambda)x_{s,t}\alpha_{t^{-1}}(e_\lambda).$$

$(\alpha_{s^{-1}}(e_\lambda))_{\lambda \in \Lambda}$ and $(\alpha_{t^{-1}}(e_\lambda))_{\lambda \in \Lambda}$ are also approximate units in J , whenever $s, t \in \Gamma$ are arbitrary and fixed so that

$$x_{s,t} \in J \iff \delta(x)_{\lambda;s,t} \rightarrow 0 \text{ as } \lambda \rightarrow \infty$$

or $x_{s,t} \in J$ iff $(\lambda \mapsto \delta(x)_{\lambda;s,t}) \in c_0(\Lambda, A)$. This proves (4) $\Leftrightarrow$ (5). $\square$

We are now ready to prove our characterizations of exactness. Condition (3) of the following theorem will be used later.

Theorem 2.3. *For a discrete group the following conditions are equivalent:*

- (1) Γ is exact.
- (2) $C_r^*(\Gamma)$ is exact.
- (3) For all Hilbert spaces H and closed subspaces $S \subseteq B(H)$,

$$UC_r^*(\Gamma, S) = \{x \in UC_r^*(\Gamma, B(H)) \mid x_{s,t} \in S \text{ for all } s, t \in \Gamma\}.$$

- (4) $UC_r^*(\Gamma, -)$ is an exact functor.

Proof. (1) $\Rightarrow$ (2): This is clear since $0 \rightarrow J \rtimes_{\alpha,r} \Gamma \rightarrow A \rtimes_{\alpha,r} \Gamma \rightarrow B \rtimes_{\dot{\alpha},r} \Gamma \rightarrow 0$ is just $0 \rightarrow J \otimes C_r^*(\Gamma) \rightarrow A \otimes C_r^*(\Gamma) \rightarrow B \otimes C_r^*(\Gamma) \rightarrow 0$ when α is the trivial action $\alpha_s = \text{id}$ for all s .

(2) $\Rightarrow$ (3): Clearly $\subseteq$ in (3) always holds. For the reverse inclusion note that if $C_r^*(\Gamma)$ is exact, then there exists a net (k_α) as in Theorem 2.1(3), and the Schur multiplier Θ_{k_α} defines a completely positive map on $B(\ell^2(\Gamma) \otimes H)$ such that $\Theta_{k_\alpha}(x) \rightarrow x$ is norm for all $x \in UC_r^*(\Gamma, B(H))$. Moreover, all the images $\Theta_{k_\alpha}(\{x \in UC_r^*(\Gamma, B(H)) \mid x_{s,t} \in S \text{ for all } s, t \in \Gamma\})$ consist of finite width matrices only, which must have entries in S whose norms are uniformly bounded. Therefore they form subsets of $UC_r^*(\Gamma, S)$. Since $UC_r^*(\Gamma, S)$ is norm closed and $\Theta_{k_\alpha}(x) \rightarrow x$ for $x \in UC_r^*(\Gamma, B(H))$, the claim follows.

(3) $\Rightarrow$ (4): Let $0 \rightarrow J \rightarrow A \xrightarrow{p} B \rightarrow 0$ be any exact sequence of C^* -algebras (without a Γ -action). p induces a surjective $*$ -homomorphism $\hat{p} : UC_r^*(\Gamma, A) \rightarrow UC_r^*(\Gamma, B)$, where $\ker \hat{p} = \{x \in UC_r^*(\Gamma, A) \mid x_{s,t} \in J \text{ for all } s, t \in \Gamma\}$. Thinking of

A as being faithfully represented on a Hilbert space H , i.e. $A \subseteq B(H)$, we have $UC_r^*(\Gamma, J) \subseteq \ker \hat{p} \subseteq \{x \in UC_r^*(\Gamma, B(H)) \mid x_{s,t} \in J \text{ for all } s, t \in \Gamma\} = UC_r^*(\Gamma, J)$ by assumption, hence $\ker \hat{p} = UC_r^*(\Gamma, J)$.

(4) $\Rightarrow$ (1): Let $0 \rightarrow (J, \alpha) \rightarrow (A, \alpha) \rightarrow (B, \dot{\alpha}) \rightarrow 0$ be an exact sequence of Γ -dynamical systems and $p_\alpha : A \rtimes_{\alpha, r} \Gamma \rightarrow B \rtimes_{\dot{\alpha}, r} \Gamma$ the induced surjective $*$ -homomorphism. Let $0 \rightarrow c_0(\Lambda, A) \rightarrow \ell^\infty(\Lambda, A) \xrightarrow{q} q^\infty(\Lambda, A) \rightarrow 0$ be the exact sequence before Lemma 2.2. Then by assumption the sequence

$$0 \longrightarrow UC_r^*(\Gamma, c_0(\Lambda, A)) \longrightarrow UC_r^*(\Gamma, \ell^\infty(\Lambda, A)) \xrightarrow{\hat{q}} UC_r^*(\Gamma, q^\infty(\Lambda, A)) \longrightarrow 0$$

is exact. Suppose that $x \in \ker p_\alpha$. Then by Lemma 2.2(3) $\Rightarrow$ (5) we know that $\delta(x) \in \ker \hat{q} = UC_r^*(\Gamma, c_0(\Lambda, A))$ and by Lemma 2.2(2) $\Rightarrow$ (1) it follows that $x \in J \rtimes_{\alpha, r} \Gamma$ and hence Γ is an exact group. $\square$

Now we get back to the OAP, respectively AP for Γ . Let us first note that a completely bounded multiplier $u \in M_0 A(\Gamma)$ defines a completely bounded map $\hat{M}_u \in CB(UC_r^*(\Gamma, S))$ for any operator space $S \subseteq B(H)$ by $\hat{M}_u([x_{s,t}]) = [u(st^{-1})x_{s,t}]$. (Here CB stands for completely bounded linear map.) Indeed, as in the definition of Schur multipliers, let K be a Hilbert space and $p, q : \Gamma \rightarrow K$ be bounded with $u(st^{-1}) = \langle p(s), q(t) \rangle$ for all $s, t \in \Gamma$. Define bounded linear maps $V, W : \ell^2(\Gamma) \otimes H \rightarrow \ell^2(\Gamma) \otimes H \otimes K$ by

$$V(e_s \otimes \xi) = e_s \otimes \xi \otimes p(s) \text{ and } W(e_s \otimes \xi) = e_s \otimes \xi \otimes q(s),$$

where $(e_s) \subseteq \ell^2(\Gamma)$ is the canonical orthonormal basis in $\ell^2(\Gamma)$ and $\xi \in H$. Then $V^*(x \otimes 1_K)W = \hat{M}_u(x)$ for all $x \in UC_r^*(\Gamma, S)$. This map is clearly completely bounded with $\text{cb-norm} \leq \|V\|\|W\|$. It follows that $\|\hat{M}_u\| \leq \|u\|_{M_0}$ and $\sup_S \|\hat{M}_u\| = \|u\|_{M_0}$, where the supremum is taken over all operator spaces.

For convergence properties of $\hat{M}_{u_\alpha}$ (as $\alpha \rightarrow \infty$), where (u_α) is a net in $A_c(\Gamma)$, we can now use the same kind of arguments as in [HK] for tensor products which we sketch for convenience. By a Hahn-Banach type argument the point-norm closure and the point-weak closure of the subspace $\{\hat{M}_u \mid u \in A_c(\Gamma)\} \subseteq B(UC_r^*(\Gamma, S))$ are identical. Thus the identity map is in the point-norm closure of this subspace iff there is a net $(u_\alpha) \subseteq A_c(\Gamma)$ such that $\varphi(\hat{M}_{u_\alpha}(x)) \rightarrow \varphi(x)$ for all $\varphi \in UC_r^*(\Gamma, S)^*$ and $x \in UC_r^*(\Gamma, S)$. Now let us view $u \mapsto \varphi(\hat{M}_u(x))$ as a bounded linear functional $\omega_{\varphi, x} \in M_0 A(\Gamma)^*$. Then $u_\alpha \rightarrow 1$ in the $\sigma(M_0 A(\Gamma), Q(\Gamma))$ -topology implies $\varphi(\hat{M}_{u_\alpha}(x)) \rightarrow \varphi(x)$ for all $\varphi \in UC_r^*(\Gamma, S)^*$ and $x \in UC_r^*(\Gamma, S)$, provided we can show the following.

Lemma 2.4. $\omega_{\varphi, x}$ is in $Q(\Gamma)$ for all $\varphi \in UC_r^*(\Gamma, S)^*$ and $x \in UC_r^*(\Gamma, S)$.

Proof. Note first that $\|\hat{M}_u\| \leq \|u\|_{M_0}$ implies $\|\omega_{\varphi, x}\| \leq \|\varphi\|\|x\|$. Since $Q(\Gamma)$ is complete we may thus assume that x has finite width. But then $\omega_{\varphi, x}(u)$ actually only depends on the value of the function $u : \Gamma \rightarrow \mathbb{C}$ on a finite subset of Γ which clearly means that $u \mapsto \omega_{\varphi, x}(u)$ is in $Q(\Gamma)$ in this case. $\square$

Thus the existence of (u_α) in $A_c(\Gamma)$ with $u_\alpha \rightarrow 1$ in the $\sigma(M_0 A(\Gamma), Q(\Gamma))$ -topology guarantees that we may choose (u_α) so that $\hat{M}_{u_\alpha}(x) \rightarrow x$ for all $x \in UC_r^*(\Gamma, S)$.

3. THE INVARIANT TRANSLATION APPROXIMATION PROPERTY AND THE AP

Note that if $S \subseteq B(H)$ is a concrete operator space, then $\text{Ad}(\lambda_t \otimes \text{id})$ and $\text{Ad}(\rho_t \otimes \text{id})$ both leave $UC_r^*(\Gamma, S) \subseteq B(\ell^2(\Gamma) \otimes H)$ invariant. We say that a discrete group Γ has the *operator invariant translation approximation property* if for any closed subspace S of the compact operators $\mathcal{K}$ (on $\ell^2(\mathbb{N})$) the equality

$$UC_r^*(\Gamma, S)^\Gamma = C_r^*(\Gamma) \otimes S$$

is obtained, where $UC_r^*(\Gamma, S)^\Gamma$ denotes the set of elements in $UC_r^*(\Gamma, S)$ which are fixed under $\text{Ad}(\rho_t \otimes \text{id})$ for all $t \in \Gamma$. If Γ is exact we may replace this condition by $(UC_r^*(\Gamma) \otimes S)^\Gamma = C_r^*(\Gamma) \otimes S$:

Lemma 3.1. *Suppose that Γ is exact and $S \subseteq B(H)$ is an arbitrary closed subspace. Then $UC_r^*(\Gamma, S)^\Gamma = (UC_r^*(\Gamma) \otimes S)^\Gamma$.*

Proof. Let (k_α) be as in Theorem 2.1(3). We show first that $\Theta_{k_\alpha}(UC_r^*(\Gamma, S)^\Gamma) \subseteq UC_r^*(\Gamma) \odot S$, where $\odot$ denotes the algebraic tensor product. Given $x \in UC_r^*(\Gamma, S)^\Gamma$ we have $x_{s,t} = x_{sr,tr} \in S$ for all $s, t, r \in \Gamma$. So $x_{s,t}$ only depends on the value of st^{-1} . In particular $x_{s,t} = x_{st^{-1},e}$ for all $s, t \in \Gamma$. Let $F \subseteq \Gamma$ be finite and x^F the element obtained from x by replacing $x_{s,t}$ by 0 whenever $st^{-1} \notin F$. Then $x^F = \sum_{r \in F} \lambda_r \otimes x_{r,e}$ is in $\mathbb{C}[\Gamma] \odot S$, where $\mathbb{C}[\Gamma]$ is the group ring. Moreover, $\Theta_{k_\alpha}(x) = \Theta_{k_\alpha}(x^F)$ provided k_α has width F . But $\Theta_{k_\alpha}(x^F) = \sum_{r \in F} \Theta_{k_\alpha}(\lambda_r) \otimes x_{r,e} \in UC_r^*(\Gamma) \odot S$, thus $\Theta_{k_\alpha}(UC_r^*(\Gamma, S)^\Gamma) \subseteq UC_r^*(\Gamma) \odot S$ for all α . Since $\Theta_{k_\alpha}(x) \rightarrow x$ for $x \in UC_r^*(\Gamma, S)$, it follows that $UC_r^*(\Gamma, S)^\Gamma = UC_r^*(\Gamma) \otimes S \cap UC_r^*(\Gamma, S)^\Gamma = (UC_r^*(\Gamma) \otimes S)^\Gamma$. $\square$

Finally we can prove our result.

Theorem 3.2. *For a discrete exact group Γ the following conditions are equivalent:*

- (1) Γ has the operator invariant translation approximation property.
- (2) $UC_r^*(\Gamma, S)^\Gamma = (UC_r^*(\Gamma) \otimes S)^\Gamma = C_r^*(\Gamma) \otimes S$ for any closed subspace $S \subseteq \mathcal{K}$.
- (3) $UC_r^*(\Gamma, S)^\Gamma = (UC_r^*(\Gamma) \otimes S)^\Gamma = C_r^*(\Gamma) \otimes S$ for any Hilbert space H and any closed subspace $S \subseteq B(H)$.
- (4) Γ has the AP.

Proof. (1) $\Leftrightarrow$ (2): follows from Lemma 3.1.

(1) $\Rightarrow$ (4): given Γ exact with the operator invariant translation approximation property, it suffices to show that $C_r^*(\Gamma)$ has the slice map property for closed subspaces $S \subseteq \mathcal{K}$. Thus given such an S we have to show that $F(C_r^*(\Gamma), S) = \{x \in C_r^*(\Gamma) \otimes \mathcal{K} \mid (\varphi \otimes \text{id})(x) \in S \text{ for all } \varphi \in C_r^*(\Gamma)^*\}$ equals $C_r^*(\Gamma) \otimes S$. Now $C_r^*(\Gamma) \otimes \mathcal{K} \subseteq UC_r^*(\Gamma, \mathcal{K})$, so we may view $x \in F(C_r^*(\Gamma), S)$ as a $\Gamma \times \Gamma$ -matrix with entries in $\mathcal{K}$. Since $(\langle e_s, \cdot \rangle \otimes \text{id})(x) = x_{s,t}$, all its entries must be in S and if all entries $x_{s,t}$ are in S , then one checks that $x \in F(C_r^*(\Gamma), S)$. Since Γ is exact Theorem 2.3(3) now implies that $F(C_r^*(\Gamma), S) = C_r^*(\Gamma) \otimes \mathcal{K} \cap UC_r^*(\Gamma, S)$, and this equals $UC_r^*(\Gamma, \mathcal{K})^\Gamma \cap UC_r^*(\Gamma, S) = UC_r^*(\Gamma, S)^\Gamma = C_r^*(\Gamma) \otimes S$ if Γ has the operator invariant translation approximation property.

(4) $\Rightarrow$ (3): let $(u_\alpha) \subseteq A_c(\Gamma)$ (all of finite support) such that $\hat{M}_{u_\alpha}(x) \rightarrow x$ for all $x \in UC_r^*(\Gamma, B(H))$. $\hat{M}_{u_\alpha}$ commutes with $\text{Ad}\rho_t$ for all $t \in \Gamma$, so for all closed subspaces $S \subseteq B(H)$ the completely bounded operator $\hat{M}_{u_\alpha} \in CB(UC_r^*(\Gamma, S))$ maps invariant elements into invariant elements. Moreover, $\hat{M}_{u_\alpha}(UC_r^*(\Gamma, S)^\Gamma) \subseteq \mathbb{C}[\Gamma] \odot S$ by a similar argument as in the proof of Lemma 3.1. Since $\hat{M}_{u_\alpha}(x) \rightarrow x$

whenever $x \in UC_r^*(\Gamma, S)$, it follows that $UC_r^*(\Gamma, S)^\Gamma \subseteq C_r^*(\Gamma) \otimes S$, i.e. $UC_r^*(\Gamma, S)^\Gamma = C_r^*(\Gamma) \otimes S$.

(3) $\Rightarrow$ (1): obvious. $\square$

Remark. Is it true that if Γ is a discrete group, then $(UC_r^*(\Gamma) \otimes S)^\Gamma = UC_r^*(\Gamma)^\Gamma \otimes S$ for all closed subspaces $S \subseteq \mathcal{K}$? If so, then Roe's invariant translation approximation property is equivalent to the OAP at least if Γ is exact. We leave this question open.

ACKNOWLEDGEMENT

I thank J. Kaminker for interesting discussions and for drawing my attention to this problem during a short visit to Nottingham University supported by the London Mathematical Society. I also thank the referee for useful suggestions which improved the paper.

REFERENCES

- [ER] E. Effros, Z.-J. Ruan *Operator spaces*, London Mathematical Society Monographs **23** Oxford University Press (2000). MR1793753 (2002a:46082)
- [GK] E. Guentner, J. Kaminker *Exactness and the Novikov conjecture*, Topology **41** (2002) 411–418. MR1876896 (2003e:46097a)
- [HK] U. Haagerup, J. Kraus *Approximation properties for group C^* -algebras and group von Neumann algebras*, Trans. AMS **344** (1994) 667–699. MR1220905 (94k:22008)
- [Ki] E. Kirchberg *The Fubini theorem for exact C^* -algebras*, JOT **10** (1983) 3–8. MR0715549 (85d:46081)
- [KW] E. Kirchberg, S. Wassermann *Exact groups and continuous bundles of C^* -algebras*, Math. Ann. **315** (1999) 169–203. MR1721796 (2000i:46050)
- [Kr] J. Kraus *The slice map problem and approximation properties*, JFA **102** (1991) 116–155. MR1138840 (92m:47083)
- [Oz] N. Ozawa *Amenable actions and exactness for discrete groups*, C.R. Acad. Sci. Paris Sér. I Math. **330** (2000) 691–695. MR1763912 (2001g:22007)
- [Pi] G. Pisier *Introduction to operator space theory*, London Math. Soc. Lecture Notes Series **294** CUP (2003). MR2006539 (2004k:46097)
- [Ro] J. Roe *Lectures on coarse geometry*, University Lecture Series **31** AMS (2003). MR2007488 (2004g:53050)
- [Wa] S. Wassermann *Exact C^* -algebras and related topics*, Lecture Notes Series **19**, Seoul National University Research Institute of Mathematics (1994). MR1271145 (95b:46081)

SCHOOL OF MATHEMATICAL SCIENCES, UNIVERSITY OF NOTTINGHAM, NOTTINGHAM NG7 2RD, UNITED KINGDOM

E-mail address: jz@maths.nott.ac.uk