ON KORENBLUM'S MAXIMUM PRINCIPLE

CHUNJIE WANG

(Communicated by Joseph A. Ball)

ABSTRACT. Let $A^2(\mathbb{D})$ be the Bergman space over the open unit disk $\mathbb{D}$ in the complex plane. Korenblum's maximum principle states that there is an absolute constant $c \in (0, 1)$, such that whenever $|f(z)| \leq |g(z)|$ ($f, g \in A^2(\mathbb{D})$) in the annulus $c < |z| < 1$, then $\|f\|_{A^2} \leq \|g\|_{A^2}$. In this paper we prove that Korenblum's maximum principle holds with $c = 0.25018$.

1. INTRODUCTION

Let $\mathbb{D}$ be the open unit disk in the complex plane $\mathbb{C}$. The Bergman space $A^2(\mathbb{D})$ consists of analytic functions f in $\mathbb{D}$ such that

$$\|f\|_{A^2} = \left[\frac{1}{\pi} \int_{\mathbb{D}} |f(z)|^2 dA(z) \right]^{\frac{1}{2}} < +\infty,$$

where

$$dA(z) = dx dy = r dr d\theta, \quad z = x + iy = re^{i\theta},$$

is the Lebesgue area measure. Korenblum [4] conjectured that there is an absolute constant c , $0 < c < 1$, such that whenever $|f(z)| \leq |g(z)|$ in the annulus $c < |z| < 1$ ($f, g \in A^2(\mathbb{D})$), then $\|f\|_{A^2} \leq \|g\|_{A^2}$.

In 1999 Hayman [1] proved Korenblum's conjecture, and Hinkkanen [2] proved that Korenblum's conjecture is true for the Bergman space $A^p(\mathbb{D})$ ($p \geq 1$). But the sharp value of c even when $p = 2$ (we use γ to denote this sharp value and call it Korenblum's constant) is still unknown. However, Hayman [1] gave a lower bound on γ : $\gamma \geq 0.04$. Hinkkanen [2] improved Hayman's result that $\gamma \geq 0.15724 \dots$. Recently Schuster [5] has shown that $\gamma \geq 0.21$ in terms of Möbius pseudodistance for the annulus. On the other hand, an upper bound on γ can be found from Martin's example (see [4] or [6]): $\gamma < 0.70450 \dots$. Wang [6] gave an upper bound on γ : $\gamma < 0.69472$, and has recently [7] shown that $\gamma < 0.67795$. In this paper we prove that Korenblum's maximum principle holds with $c = 0.25018$, which is the following theorem.

Theorem 1. *Let $f(z)$ and $g(z)$ be analytic functions in the unit disk $\mathbb{D}$. Suppose that $|f(z)| \leq |g(z)|$ whenever $c < |z| < 1$, where $c = 0.25018$. Then $\|f\|_{A^2} \leq \|g\|_{A^2}$.*

Received by the editors December 10, 2004 and, in revised form, February 14, 2005.

2000 *Mathematics Subject Classification.* Primary 30C80, 30H05.

Key words and phrases. Bergman space, Korenblum's maximum principle, Fock space.

This work was supported by NNSF of China No. 10401002 and the Doctoral Foundation of Hebei University of Technology.

©2006 American Mathematical Society
 Reverts to public domain 28 years from publication

Let $c \in (0, 1)$ be a constant to be determined later. We write $D(c) = \{z \in \mathbb{C} : |z| < c\}$ and $A(c, 1) = \{z \in \mathbb{C} : c < |z| < 1\}$. Suppose that $f(z)$ and $g(z)$ are analytic functions in the unit disk $\mathbb{D}$ and that $|f(z)| \leq |g(z)|$ whenever $c < |z| < 1$. To prove Theorem 1, we need only to show that

$$(1) \quad \int_{D(c)} (|f(z)|^2 - |g(z)|^2) dA(z) \leq \int_{A(c, 1)} (|g(z)|^2 - |f(z)|^2) dA(z).$$

We may assume that $\|g\|_{A^2}$ is finite, for otherwise the result holds trivially. Hence $\|f\|_{A^2}$ is also finite, since $|f(z)| \leq |g(z)|$ whenever $c < |z| < 1$. We may also assume that g is not identically equal to 0. Define $\omega(z) = f(z)/g(z)$ for $c < |z| < 1$, so that $|\omega| \leq 1$. We may assume that $|\omega(z)| < 1$ in $c < |z| < 1$, since otherwise the result is trivial. Define $\omega_0 = \omega_0(\rho) = \max\{|\omega(z)| : |z| = \rho\}$.

To obtain a better constant c , we follow Hinkkanen [2] and Schuster [5] but use a different estimate. Note that for any $a \in (-1, 1)$,

$$(2) \quad |f|^2 - |g|^2 \leq \frac{|f - ag|^2}{1 - a^2},$$

which is a direct consequence of the following identity:

$$|f|^2 - |g|^2 = \frac{|f - ag|^2 - |af - g|^2}{1 - a^2}.$$

Using (2) and the fact that, for a subharmonic function h in $\mathbb{D}$ and $0 < r_1 < r_2 < 1$, $\int_0^{2\pi} h(r_1 e^{i\theta}) d\theta \leq \int_0^{2\pi} h(r_2 e^{i\theta}) d\theta$, we obtain

$$\begin{aligned} & \int_0^{2\pi} (|f(re^{i\theta})|^2 - |g(re^{i\theta})|^2) d\theta \\ & \leq \int_0^{2\pi} \frac{|f(re^{i\theta}) - ag(re^{i\theta})|^2}{1 - a^2} d\theta \\ & \leq \int_0^{2\pi} \frac{|f(\rho e^{i\theta}) - ag(\rho e^{i\theta})|^2}{1 - a^2} d\theta \\ & = \int_0^{2\pi} \frac{|\omega(\rho e^{i\theta}) - a|^2}{(1 - a^2)(1 - |\omega(\rho e^{i\theta})|^2)} (|g(\rho e^{i\theta})|^2 - |f(\rho e^{i\theta})|^2) d\theta \\ & \leq \gamma(\rho) \int_0^{2\pi} (|g(\rho e^{i\theta})|^2 - |f(\rho e^{i\theta})|^2) d\theta, \end{aligned}$$

where $a \in (0, 1)$ is a quantity depending on $\rho \in (c, 1)$ but not on r and θ , and

$$\gamma(\rho) = \max \left\{ \frac{|\omega(z) - a|^2}{(1 - a^2)(1 - |\omega(z)|^2)} : |z| = \rho \right\}.$$

Let us keep ρ fixed and multiply both sides of this inequality by r , and then integrate with respect to r from 0 to c . We obtain

$$(3) \quad \frac{1}{\gamma(\rho)} \int_{D(c)} (|f(z)|^2 - |g(z)|^2) dA(z) \leq \frac{c^2}{2} \int_0^{2\pi} (|g(\rho e^{i\theta})|^2 - |f(\rho e^{i\theta})|^2) d\theta.$$

Multiplying both sides of (3) by ρ and integrating with respect to ρ from c to 1 yields

$$(4) \quad \int_c^1 \frac{\rho}{\gamma(\rho)} d\rho \int_{D(c)} (|f(z)|^2 - |g(z)|^2) dA(z) \leq \frac{c^2}{2} \int_{A(c, 1)} (|g(z)|^2 - |f(z)|^2) dA(z).$$

Thus if we can choose c so that

$$(5) \quad \int_c^1 \frac{\rho}{\gamma(\rho)} d\rho \geq \frac{c^2}{2},$$

then Theorem 1 is proved.

So we only need to find a suitable bound for $\gamma(\rho)$.

2. THE MÖBIUS PSEUDODISTANCE

We recall that the pseudohyperbolic distance d between two points $\alpha, \beta \in \mathbb{D}$ is defined by

$$d(\alpha, \beta) = \left| \frac{\alpha - \beta}{1 - \bar{\alpha}\beta} \right|.$$

The Möbius pseudodistance for a domain $D \subset \mathbb{D}$ is defined by

$$c_D^*(a, z) = \sup\{d(\omega(a), \omega(z)) : \omega \in \mathcal{H}(D, \mathbb{D})\},$$

where $a, z \in D$ and $\mathcal{H}(D, \mathbb{D})$ denotes the set of analytic functions from D to $\mathbb{D}$ (see [3] for an excellent survey). Note that both of the two pseudodistances are Möbius invariant.

In [5] Schuster showed that

$$c_{A(c,1)}^*(\rho, \rho e^{i\theta}) = \rho \prod_{n=1}^{\infty} f_n(\rho, c) g_n(\rho, c, \theta),$$

where

$$f_n(\rho, c) = \frac{(1 + \rho^2 c^{2n-1})(1 + \rho^{-2} c^{2n-1})}{(1 + c^{2n-1})^2}$$

and

$$g_n(\rho, c, \theta) = \frac{\sqrt{1 - 2c^{2n-2} \cos \theta + c^{4n-4}} \sqrt{1 - 2c^{2n} \cos \theta + c^{4n}}}{\sqrt{1 - 2\rho^2 c^{2n-2} \cos \theta + \rho^4 c^{4n-4}} \sqrt{1 - 2\rho^{-2} c^{2n} \cos \theta + \rho^{-4} c^{4n}}}.$$

In the following we use $F \sim G$ to indicate that the quotient F/G of two functions F and G is positive. We want to give an estimate for $c_{A(c,1)}^*(\rho, \rho e^{i\theta})$ as in [5]. First, we proceed to show that when $n \geq 3$ and $c \leq 1/\sqrt{2}$,

$$(6) \quad g_n(\rho, c, \theta) \leq g_n(\rho, c, \pi).$$

Since $g_n(\rho, c, \theta)$ is an even function of θ , we may assume that $0 \leq \theta \leq \pi$. So it suffices for us to prove that $\frac{\partial g_n^2(\rho, c, \theta)}{\partial \theta} > 0$ for $0 < \theta < \pi$. A direct calculation shows that for $0 < \theta < \pi$,

$$\begin{aligned} \frac{\partial g_n^2(\rho, c, \theta)}{\partial \theta} &\sim -4c^{4n-2}(1 + c^{4n-2}) \cos^2 \theta - 4c^{6n-4}(1 + \rho^2)(1 + \rho^{-2}c^2) \cos \theta \\ &+ (1 + c^{4n-2})(1 - 2c^{4n-2} + c^{8n-4} - \rho^2 c^{4n-2} - \rho^2 c^{4n-4} - \rho^{-2} c^{4n-2} - \rho^{-2} c^{4n}). \end{aligned}$$

Write

$$\begin{aligned} x_0 &= \frac{-c^{4n-3}(1 + \rho^2)(1 + \rho^{-2}c^2)}{2c^{2n-1}(1 + c^{4n-2})} \\ &+ \frac{\sqrt{(1 - \rho^2 c^{4n-4})(1 - \rho^2 c^{4n-2})(1 - \rho^{-2} c^{4n-2})(1 - \rho^{-2} c^{4n})}}{2c^{2n-1}(1 + c^{4n-2})}. \end{aligned}$$

Then we only need to prove $x_0 > 1$, which is equivalent to

$$\begin{aligned} \beta(\rho, c) &= (1 - \rho^2 c^{4n-4})(1 - \rho^2 c^{4n-2})(1 - \rho^{-2} c^{4n-2})(1 - \rho^{-2} c^{4n}) \\ &\quad - c^{4n-2}(2(1 + c^{4n-2}) + c^{2n-2}(1 + \rho^2)(1 + \rho^{-2} c^2))^2 > 0. \end{aligned}$$

It is easy to check that

$$\frac{\partial \beta(\rho, c)}{\partial \rho} \sim c - \rho^2,$$

and for any $n \geq 3, c \leq 1/\sqrt{2}$,

$$\begin{aligned} \beta(1, c) &= \beta(c, c) = (1 - c^{4n-4})(1 - c^{4n-2})^2(1 - c^{4n}) \\ &\quad - 4c^{4n-2}(1 + c^{4n-2} + c^{2n-2}(1 + c^2))^2 \\ &\sim 1 + c^{8n-4} - 6c^{4n-2} - c^{2n-2} - c^{2n} - c^{6n-4} - c^{6n-2} > 0. \end{aligned}$$

Then we get

$$\beta(\rho, c) \geq \beta(1, c) = \beta(c, c) > 0.$$

The desired inequality (6) now follows.

Next, a long and tedious calculation shows that

$$(7) \quad \prod_{n=1}^4 g_n(\rho, c, \theta) \leq \prod_{n=1}^4 g_n(\rho, c, \pi).$$

Thus we have

$$\begin{aligned} c_{A(c,1)}^*(\rho, \rho e^{i\theta}) &\leq \rho \prod_{n=1}^{\infty} f_n(\rho, c) g_n(\rho, c, \pi) \\ &= \rho \prod_{n=1}^{\infty} \left[\frac{(1 + \rho^2 c^{2n-1})(1 + \rho^{-2} c^{2n-1})}{(1 + c^{2n-1})^2} \cdot \frac{(1 + c^{2n-2})(1 + c^{2n})}{(1 + \rho^2 c^{2n-2})(1 + \rho^{-2} c^{2n})} \right] \\ &= 2\rho(1 + \rho^{-2}c) \prod_{n=1}^{\infty} \frac{(1 + \rho^2 c^{2n-1})(1 + \rho^{-2} c^{2n+1})(1 + c^{2n})^2}{(1 + \rho^2 c^{2n-2})(1 + \rho^{-2} c^{2n})(1 + c^{2n-1})^2}. \end{aligned}$$

Note that every term in the last infinite product is less than 1. To use *Mathematica*, we choose

$$K = 2\rho(1 + \rho^{-2}c) \prod_{n=1}^{10} \frac{(1 + \rho^2 c^{2n-1})(1 + \rho^{-2} c^{2n+1})(1 + c^{2n})^2}{(1 + \rho^2 c^{2n-2})(1 + \rho^{-2} c^{2n})(1 + c^{2n-1})^2}.$$

3. ESTIMATES FOR $\gamma(\rho)$

In the previous section it has been proved that

$$d(\omega(z), \omega_0) = \left| \frac{\omega(z) - \omega_0}{1 - \omega_0 \omega(z)} \right| \leq K < 1$$

for $|z| = \rho$. Write

$$\frac{\omega(z) - \omega_0}{1 - \omega_0 \omega(z)} = t e^{i\theta},$$

where $0 \leq t \leq K, \theta \in [0, 2\pi]$. It is easy to check that $|\omega(z)| \leq \omega_0$ is equivalent to

$$\cos \theta \leq -\frac{t(1 + \omega_0^2)}{2\omega_0} < 0.$$

To get a bound for $\gamma(\rho)$, we choose $a = \omega_0^\alpha$, where

$$\alpha = \sqrt{\frac{1+3K^2}{1-K^2}}.$$

Then for $|z| = \rho$,

$$\begin{aligned} & \frac{|\omega(z) - a|^2}{(1-a^2)(1-|\omega(z)|^2)} \\ &= \frac{(\omega_0 - \omega_0^\alpha)^2 + t^2(1 - \omega_0^{\alpha+1})^2 + 2\omega_0 t(1 - \omega_0^{\alpha-1})(1 - \omega_0^{\alpha+1}) \cos \theta}{(1 - \omega_0^{2\alpha})(1 - \omega_0^2)(1 - t^2)} \\ &\leq \frac{(\omega_0^2 - \omega_0^\alpha)^2(1 - t^2) + t^2\omega_0^{\alpha-1}(1 - \omega_0^2)^2}{(1 - \omega_0^{2\alpha})(1 - \omega_0^2)(1 - t^2)} \\ &\triangleq H(\rho, t, \omega_0). \end{aligned}$$

To estimate $H(\rho, t, \omega_0)$, we need the following two lemmas.

Lemma 1. For any $0 \leq u \leq v < 1, x \in (0, 1)$, we have

$$(8) \quad \frac{x^u - x^v}{1 - x^{u+v}} \leq \frac{v - u}{v + u}.$$

It can be easily checked. So we omit the details.

Lemma 2. For any $u \geq 1, x \in (0, 1)$, we have

$$(9) \quad \frac{(x - x^u)^2}{(1 - x^2)(1 - x^{2u})} \leq \frac{(u - 1)^2}{4u}.$$

Proof. For any $u \geq 1, x \in (0, 1)$, define

$$\Phi(x) = \frac{(x - x^u)^2}{(1 - x^2)(1 - x^{2u})}.$$

It is easy to check that

$$\begin{aligned} \Phi'(x) &\sim 2(x - x^u)(1 - ux^{u-1})(1 - x^2)(1 - x^{2u}) \\ &\quad - (x - x^u)^2(-2x - 2ux^{2u-1} + (2u + 2)x^{2u+1}) \\ &\sim (1 - x^{u+1})[1 - x^{2u} - ux^{u-1}(1 - x^2)] \\ &\sim 1 - u \frac{x^{u-1} - x^{u+1}}{1 - x^{2u}}. \end{aligned}$$

Applying Lemma 1, we know $\Phi'(x) > 0$. Hence $\Phi(x)$ is an increasing function of x . The desired inequality (9) then follows. $\square$

Note that $\alpha \geq 1$. We use Lemma 2 and Lemma 1 to obtain

$$\begin{aligned} H(\rho, t, \omega_0) &= \frac{(\omega_0 - \omega_0^\alpha)^2}{(1 - \omega_0^{2\alpha})(1 - \omega_0^2)} + \frac{t^2(\omega_0^{\alpha-1} - \omega_0^{\alpha+1})}{(1 - t^2)(1 - \omega_0^{2\alpha})} \\ &\leq \frac{(\alpha - 1)^2}{4\alpha} + \frac{t^2}{\alpha(1 - t^2)} \\ &\leq \frac{(\alpha - 1)^2}{4\alpha} + \frac{K^2}{\alpha(1 - K^2)} \\ &= \frac{1}{2} \left(\sqrt{\frac{1+3K^2}{1-K^2}} - 1 \right). \end{aligned}$$

Putting things together, we obtain

$$(10) \quad \gamma(\rho) \leq \frac{1}{2} \left(\sqrt{\frac{1+3K^2}{1-K^2}} - 1 \right).$$

Using *Mathematica*, we obtain that (5) holds with $c = 0.25018$.

4. REMARKS

First, write

$$u_n(\rho, c, \theta) = \frac{\left| \exp\left\{\frac{\pi}{\log(1/c)}(\theta + 2n\pi)\right\} - 1 \right|}{\left\{ 1 + \exp\left\{\frac{2\pi}{\log(1/c)}(\theta + 2n\pi)\right\} - 2 \exp\left\{\frac{\pi}{\log(1/c)}(\theta + 2n\pi)\right\} \cos\left(2\pi \frac{\log \rho}{\log c}\right) \right\}^{\frac{1}{2}}}.$$

In fact, Hinkkanen gave an estimate for K (take a closer look at the proof in [2]): $K = \prod_{n=-\infty}^{\infty} u_n(\rho, c, \pi)$, which can be proved by a calculus argument. If we use the above estimate for K and Hinkkanen's estimate for $\gamma(\rho)$ (also use (4) and (5) here), a better constant than the original one can be obtained: $c = 0.1921 \dots$. Thus Korenblum's maximum principle holds for general Bergman space $A^p(\mathbb{D})$ ($p \geq 1$) with an absolute constant $c = 0.1921 \dots$. Also, this is true with $c = 0.21$. However, we cannot prove that is true with $c = 0.25018$.

Next, following Schuster [5] and using the estimate (10) in the Fock space setting, we can prove that Theorem 2 in [5] holds with $c = 0.7248$. Namely, we have

Theorem 2. *Suppose that $f(z)$ and $g(z)$ are two entire functions satisfying $|f(z)| \leq |g(z)|$ for $|z| > c$, where $c = 0.7248$. Then $\|f\|_F \leq \|g\|_F$.*

REFERENCES

- [1] W. K. Hayman, On a conjecture of Korenblum, *Analysis (Munich)* **19**(1999), 195-205. MR1705360 (2000e:30041)
- [2] A. Hinkkanen, On a maximum principle in Bergman space, *J. Anal. Math.* **79**(1999), 335-344. MR1749317 (2000m:30033)
- [3] M. Jarnicki and P. Pflug, *Invariant distances and metrics in complex analysis*, de Gruyter Expositions in Mathematics, 9. Walter de Gruyter and Co., Berlin, 1993. MR1242120 (94k:32039)
- [4] B. Korenblum, A maximum principle for the Bergman space, *Publ. Mat.* **35**(1991), 479-486. MR1201570 (93j:30018)
- [5] A. Schuster, The maximum principle for the Bergman space and the Möbius pseudodistance for the annulus, preprint.
- [6] C. Wang, Refining the constant in a maximum principle for the Bergman space, *Proc. Amer. Math. Soc.* **132**(2004), 853-855. MR2019965 (2004i:30017)
- [7] C. Wang, On Korenblum's constant, *J. Math. Anal. Appl.* **296**(2004), 262-264. MR2070507 (2005b:30038)

DEPARTMENT OF MATHEMATICS, HEBEI UNIVERSITY OF TECHNOLOGY, TIANJIN 300130, PEOPLE'S REPUBLIC OF CHINA

E-mail address: wcj498@eyou.com