A FAMILY OF SCHOTTKY GROUPS ARISING FROM THE HYPERGEOMETRIC EQUATION

TAKASHI ICHIKAWA AND MASAACKI YOSHIDA

(Communicated by Carmen C. Chicone)

ABSTRACT. We study a complex 3-dimensional family of classical Schottky groups of genus 2 as monodromy groups of the hypergeometric equation. We find non-trivial loops in the deformation space; these correspond to continuous integer-shifts of the parameters of the equation.

1. INTRODUCTION

When the three exponents of the hypergeometric differential equation are purely imaginary, its monodromy group is a classical Schottky group of genus 2; such groups form a real 3-dimensional family ([IY]). Since, under a small deformation, a Schottky group remains to be a Schottky group, by deforming the parameters of the hypergeometric equation, we have a complex 3-dimensional family of Schottky groups. We introduce (in §4) a complex 3-dimensional family S of classical Schottky groups, containing the above ones coming from pure-imaginary-exponents cases, equipped with some additional structure. We study a structure of S , and construct loops (real 1-dimensional families of classical Schottky groups) in S generating the fundamental group of S . These loops correspond to *continuous* interger-shifts of the parameters of the hypergeometric equation.

Here we give a few examples of general background references. The hypergeometric function: [IKSY], Schottky groups: Chapter 4 of [MSW] and Chapter 5 of [BBEIM], history: [Gray].

2. THE HYPERGEOMETRIC EQUATION

We consider the hypergeometric differential equation

$$E(a, b, c) : x(1-x) \frac{d^2 u}{dx^2} + \{c - (a+b+1)x\} \frac{du}{dx} - abu = 0.$$

For (any) two linearly independent solutions u_1 and u_2 , the (multi-valued) map

$$s : X := \mathbb{C} - \{0, 1\} \ni x \longmapsto z = u_1(x) : u_2(x) \in \mathbb{P}^1 := \mathbb{C} \cup \{\infty\}$$

is called a Schwarz map (or Schwarz's s -map). If we choose as solutions u_1 and u_2 , near $x = 0$, a holomorphic one and x^{1-c} times a holomorphic one, respectively,

Received by the editors October 1, 2004 and, in revised form, February 24, 2005.

2000 *Mathematics Subject Classification*. Primary 33C05, 30F10, 30F40.

Key words and phrases. Hypergeometric equation, monodromy group, Schottky group.

©2006 American Mathematical Society
 Reverts to public domain 28 years from publication

then the circuit matrices around $x = 0$ and 1 are given by

$$\gamma_1 = \begin{pmatrix} 1 & 0 \\ 0 & e^{2\pi i(1-c)} \end{pmatrix} \quad \text{and} \quad \gamma_2 = P^{-1} \begin{pmatrix} 1 & 0 \\ 0 & e^{2\pi i(c-a-b)} \end{pmatrix} P,$$

respectively, where P is a connection matrix given by

$$P = \begin{pmatrix} \frac{\Gamma(c)\Gamma(c-a-b)}{\Gamma(c-a)\Gamma(c-b)} & \frac{\Gamma(2-c)\Gamma(c-a-b)}{\Gamma(1-a)\Gamma(1-b)} \\ \frac{\Gamma(c)\Gamma(a+b-c)}{\Gamma(a)\Gamma(b)} & \frac{\Gamma(2-c)\Gamma(a+b-c)}{\Gamma(a-c+1)\Gamma(b-c+1)} \end{pmatrix};$$

here Γ denotes the Gamma function. Note that the matrices act on the row vector (u_1, u_2) from the right. These generate the monodromy group $M(a, b, c)$ of the equation $E(a, b, c)$.

3. HYPERGEOMETRIC EQUATIONS WITH PURELY IMAGINARY EXPONENT-DIFFERENCES ([SY], [IY])

If the exponent-differences

$$\lambda = 1 - c, \quad \mu = c - a - b, \quad \nu = b - a$$

(at the singular points $x = 0, 1$ and ∞ , respectively) are purely imaginary, the image of the upper-half part

$$X^+ := \{x \in X \mid \Im x \geq 0\}$$

is bounded by the three circles (in the s -plane) which are images of the three intervals $(-\infty, 0)$, $(0, 1)$ and $(1, +\infty)$; put

$$C_1 = s((-\infty, 0)), \quad C_3 = s((0, 1)), \quad C_2 = s((1, +\infty)).$$

Note that if we continue s analytically through the interval, say $(0, 1)$, to the lower-half part $X^- := \{x \in X \mid \Im x \leq 0\}$, then the image of X^- is the mirror image of that of X^+ under the reflection with center C_3 .

The reflection with center (mirror) C_j will be denoted by ρ_j ($j = 1, 2, 3$). The monodromy group $M(a, b, c)$ is the group consisting of even words of ρ_j ($j = 1, 2, 3$). This group is a Schottky group of genus 2. The domain of discontinuity modulo $M(a, b, c)$ is a curve of genus 2 defined over the reals.

Choosing the solutions u_1 and u_2 suitably, we can assume that the centers of the three circles are on the real axis, and that the circles C_2 and C_3 are inside the circle C_1 (see Figure 1). Let C'_1 and C'_2 be the mirror images with respect to C_3 of the circles C_1 and C_2 , respectively. Then $\gamma_1 := \rho_3\rho_1$ maps the inside of C_1 onto the inside of C'_1 ; and $\gamma_2 := \rho_3\rho_2$ maps the inside of C_2 onto the outside of C'_2 . Note that one of the two fixed points of γ_1 is inside C'_1 , and the other one is outside C_1 ; one of the two fixed points of γ_2 is inside C'_2 , and the other one is inside C_2 . The transformations γ_1 and γ_2 are conjugate to those given in §2.

We thus have four disjoint disks D_1, D'_1, D_2 , and D'_2 , whose centers are on the real axis, and two loxodromic transformations γ_1 and γ_2 taking D_1 and D_2 to the complementary disks of D'_1 and D'_2 , respectively. Note that one of the two fixed points of γ_1 is in D'_1 and the other one in D_1 ; one of the two fixed points of γ_2 is in D'_2 and the other one in D_2 .

FIGURE 1. The circles and the disks

4. THE MODULI SPACE S

Two loxodromic transformations γ_1 and γ_2 generate a (classical) Schottky group if and only if there are four disjoint closed disks D_1, D'_1, D_2 , and D'_2 , such that γ_1 and γ_2 take D_1 and D_2 to the complementary disks of D'_1 and D'_2 , respectively. Throughout the paper, disks are always assumed to be *closed*; we simply call them *disks*. The closure of the complement of a disk in $\mathbb{P}^1$ will simply be called *the complementary disk of the disk*.

Definition. Let S be the space of two loxodromic transformations γ_1 and γ_2 equipped with four disjoint disks D_1, D'_1, D_2 , and D'_2 , such that γ_1 and γ_2 take D_1 and D_2 to the complementary disks of D'_1 and D'_2 , respectively.

The space has a natural structure of a complex 3-dimensional manifold. We are interested in its homotopic property.

Note that if a loxodromic transformation γ takes a disk D onto the complementary disk of a disk D' ($D \cap D' = \emptyset$), then γ has a fixed point in D and the other fixed point in D' .

A loxodromic transformation is determined by the two fixed points and the multiplier.

Lemma 1. *For two given disjoint disks D and D' , there is a unique point $F \in D$ (resp. $F' \in D'$) such that by a(ny) linear fractional transformation taking F (resp. F') to ∞ , the two circles ∂D and $\partial D'$ are transformed into concentric circles.*

Proof. Let the two disks be given as

$$D : |z - a| \leq r, \quad D' : |z - a'| \leq r'.$$

By the transformation $z \rightarrow w$ defined by

$$z = \frac{1}{w} + \zeta$$

taking ζ to ∞ , the circles $C = \partial D$ and $C' = \partial D'$ are transformed into circles with centers

$$c := \frac{\bar{\zeta} - \bar{a}}{r^2 - |\zeta - a|^2}, \quad c' := \frac{\bar{\zeta} - \bar{a}'}{r'^2 - |\zeta - a'|^2},$$

respectively. Equating c and c' , we have the quadric equation

$$\zeta^2 + \left(-a - a' + \frac{r^2 - r'^2}{\bar{a} - \bar{a}'}\right) \zeta + aa' + \frac{ar'^2 - a'r^2}{\bar{a} - \bar{a}'} = 0.$$

Put $\zeta = (a - a')\eta + a'$. Note that $\zeta = a'$ and a correspond to $\eta = 0$ and 1 , respectively. Then the equation above for ζ reduces to

$$\eta^2 + \left(-1 + \frac{r^2 - r'^2}{|a - a'|^2}\right) \eta + \frac{r'^2}{|a - a'|^2} = 0.$$

It is a high school mathematics problem to see that each of the two roots of this equation is in each of the two intervals

$$\left(0, \frac{r'}{|a - a'|}\right) \quad \text{and} \quad \left(1 - \frac{r}{|a - a'|}, 1\right).$$

Since concentric circles are mapped to concentric circles under linear transformations, this completes the proof. $\square$

Remark 1. The circles ∂D and $\partial D'$ are Apollonius circles with respect to the two centers F and F' .

Lemma 2. *For two given disjoint disks D and D' and a point f' in the interior of D' , there is a 1-parameter family (parametrized by a circle) of loxodromic transformations γ which take D onto the complementary disk of D' , and fix f' . The absolute value $|m|$ of the multiplier m of γ is determined by the given data.*

(1) *If $f' \neq F'$, then the other fixed point $f \in D$ of γ must be on the Apollonius circle $A = A(f')$ in D determined by the two centers of the circles ∂D and $\partial D'$, and proportion $|m|$. $\arg m \in \mathbb{R}/2\pi\mathbb{Z}$ determines f , and vice versa.*

(2) *If $f' = F'$, then the other fixed point is $F \in D$. $\arg m \in \mathbb{R}/2\pi\mathbb{Z}$ remains free.*

Proof. We can assume that $f = 0$ and $f' = \infty$, so that the transformation in question can be presented by $z \mapsto mz$. Let c and r be the center and the radius of the disk D , and c' and r' those of the complementary disk of D' . Then we have $c' = mc$ and $r' = |m|r$.

(1) If $c \neq c'$, $f (= 0)$ is on the Apollonius circle

$$A : |f - c'| = |m||f - c|$$

with centers c and c' , and proportion $|m|$. It is easy to see that this circle is in D . (See Figure 2.)

(2) If $c = c' = 0$, then the Apollonius circle reduces to a point. $\square$

In case (2), we blow up the point F to be the circle $\mathbb{R}/2\pi\mathbb{Z}$, and call this circle also the Apollonius circle A . Under this convention, two disjoint disks D and D' , an interior point $f' \in D'$, and a point f on the Apollonius circle A uniquely determine a loxodromic transformation.

Thus an element of S can be determined by four disjoint disks D_1, D'_1, D_2, D'_2 , two interior points $f'_1 \in D'_1, f'_2 \in D'_2$, and two points f_1 on the Apollonius circle A_1 (determined by D_1, D'_1, f'_1) in D_1 , and f_2 on the Apollonius circle A_2 (determined by D_2, D'_2, f'_2) in D_2 . Since disks are contractible, we have

Proposition 1. *The fundamental group of S can be generated by the moves of the four disjoint disks D_1, D'_1, D_2, D'_2 , and the moves of f_1 in A_1 , and f_2 in A_2 .*

 FIGURE 2. Apollonius circle in D

In the next section we explicitly construct loops (real 1-dimensional families of classical Schottky groups) in S .

5. LOOPS IN S

Let γ_1 and γ_2 be as in §2. The fixed points of γ_1 are $\{0, \infty\}$, and those of γ_2 are

$$f_2 = \frac{\Gamma(c)\Gamma(a-c+1)\Gamma(b-c+1)}{\Gamma(2-c)\Gamma(a)\Gamma(b)} \quad \text{and} \quad f'_2 = \frac{\Gamma(c)\Gamma(1-a)\Gamma(1-b)}{\Gamma(2-c)\Gamma(c-a)\Gamma(c-b)}.$$

We change the coordinate z by multiplying by $1/f_2$. Then the fixed points of γ_1 remain to be $\{0, \infty\}$, and those of γ_2 become 1 and

$$\alpha = g(a)g(b), \quad \text{where} \quad g(x) = \frac{\sin(\pi c - \pi x)}{\sin(\pi x)}.$$

Indeed we have

$$\frac{f'_2}{f_2} = \frac{\Gamma(1-a)\Gamma(1-b)}{\Gamma(c-a)\Gamma(c-b)} \cdot \frac{\Gamma(a)\Gamma(b)}{\Gamma(a-c+1)\Gamma(b-c+1)} \quad \text{and} \quad \Gamma(x)\Gamma(1-x) = \frac{\pi}{\sin(\pi x)}.$$

Definition. When the three exponent-differences are purely imaginary:

$$\lambda = i\theta_0, \quad \mu = i\theta_1, \quad \nu = i\theta_2, \quad \theta_0, \theta_1, \theta_2 > 0,$$

the generators $\{\gamma_1, \gamma_2\}$ of the monodromy group of $E(a, b, c)$ given in §2 (take the four disks given in §3) form a simply connected real 3-dimensional submanifold of S ; this submanifold is called S_0 .

In this section, we construct loops in S , with base in S_0 , which generate the fundamental group of S . When the exponent-differences are as above, note that the parameters can be expressed as

$$a = \frac{1}{2} - \frac{i}{2}(\theta_0 + \theta_1 + \theta_2), \quad b = \frac{1}{2} - \frac{i}{2}(\theta_0 + \theta_1 - \theta_2), \quad c = 1 - i\theta_0.$$

For clarity, we denote the four disks D_1, D'_1, D_2, D'_2 by

$$D_0 (\ni 0), \quad D_\infty (\ni \infty), \quad D_1 (\ni 1), \quad D_\alpha (\ni \alpha).$$

FIGURE 3. The disk D_α travels around the disk D_0

5.1. The disk D_α travels around the disk D_0 . We construct a loop in S (with base in S_0), which is represented by a travel of the disk D_α around the disk D_0 with a change of argument (by 2π) of the multiplier of γ_2 . We fix c and the real part of a (as $1/2$), and let the real part of b move from $1/2$ to $3/2$; the imaginary parts of a and b are so chosen that the monodromy group $M(a, b, c)$ remains to be a Schottky group along the move. Putting $c = 1 - i\theta_0$, we have

$$g(x) = \frac{\varepsilon^{-1} - \varepsilon e^{2\pi i x}}{1 - e^{2\pi i x}} = \varepsilon + \frac{\varepsilon^{-1} - \varepsilon}{1 - e^{2\pi i x}}, \quad \text{where } \varepsilon = e^{-\pi\theta_0} < 1.$$

Set

$$r = g\left(\frac{1}{2} - \frac{i}{2}(\theta_0 + 2)\right), \quad R = g\left(\frac{1}{2} - \frac{i}{2}(\theta_0 + 1)\right).$$

Then we have

$$0 < \varepsilon < r = \varepsilon \frac{e^{2\pi} + \varepsilon^{-1}}{e^{2\pi} + \varepsilon} < R = \varepsilon \frac{e^\pi + \varepsilon^{-1}}{e^\pi + \varepsilon} < 1.$$

Let D_0 be the disk with center at 0 and with radius εr , and D_∞ the complementary disk of the disk with center at 0 and with radius $\varepsilon^{-1}r$. Note that we have

$$0 < \varepsilon r < \varepsilon R < R < 1 < \varepsilon^{-1}r,$$

and that γ_1 maps D_0 onto the outside disk of D_∞ .

Define real continuous functions $\phi(t)$ and $\psi(t)$ for $0 \leq t \leq 1$ by

$$Re^{2\pi i t} = g\left(\frac{1}{2} + \phi(t) - \frac{i}{2}(\theta_0 + \psi(t))\right).$$

Since

$$2\pi i \left(\frac{1}{2} + \phi(t) - \frac{i}{2}(\theta_0 + \psi(t))\right) = \log(Re^{2\pi i t} - \varepsilon^{-1}) - \log(Re^{2\pi i t} - \varepsilon),$$

the function $\phi(t)$ is monotone increasing with $\phi(0) = 0$, $\phi(1) = 1$, and $\psi(t)$ satisfies $\psi(0) = \psi(1) = 1$. Choose θ_1 so large that

- $\theta_1 > \max\{\psi(t) \mid 0 \leq t \leq 1\}$, and that
- for any z in the ring $\{\varepsilon R \leq |z| \leq R\}$, there are two disjoint disks D_1 ($\ni 1$) and D_z ($\ni z$), and a fractional linear transformation with multiplier $e^{-2\pi\theta_1}$ mapping D_1 onto the complementary disk of D_z .

Now set $\theta_2(t) = \theta_1 - \psi(t) > 0$ and deform the parameters a and b as

$$\begin{aligned} a(t) &= \frac{1}{2} - \frac{i}{2}(\theta_0 + \theta_1 + \theta_2(t)), \\ b(t) &= \frac{1}{2} + \phi(t) - \frac{i}{2}(\theta_0 + \theta_1 - \theta_2(t)). \end{aligned}$$

Then we have $\varepsilon < g(a(t)) < 1, |g(b(t))| = R$, and so $\alpha(t) = g(a(t))g(b(t))$ satisfies

$$\varepsilon R < |\alpha(t)| = g(a(t))|g(b(t))| < R.$$

Thus $\alpha(t)$ travels around the disk D_0 in the ring $\{\varepsilon R \leq |z| \leq R\}$, and there are two disjoint disks D_1 ($\ni 1$) and $D_{\alpha(t)}$ ($\ni \alpha$) in the ring $\{\varepsilon r < |z| < \varepsilon^{-1}r\}$, and a transformation $\gamma_2(t)$ with multiplier $e^{2\pi i\mu} = e^{-2\pi\theta_1}$ which maps D_1 onto the complementary disk of $D_{\alpha(t)}$.

5.2. The disk D_α travels around the disk D_1 . We construct a loop in S (with base in S_0), which is represented by a travel of the disk D_α around the disk D_1 with a change of argument (by 2π) of the multiplier of γ_2 . We fix c, a and the imaginary part of b , and let the real part of b move from $1/2$ to $3/2$. Take $\theta_0 > 0$ and $\theta' < 0$ satisfying

$$(1) \quad \frac{e^{\pi\theta'}(1 + e^{\pi\theta'})}{1 - e^{\pi\theta'}} < \varepsilon^{-2}, \quad \varepsilon = e^{-\pi\theta_0}.$$

Set

$$r = \varepsilon + \frac{\varepsilon^{-1} - \varepsilon}{1 + e^{\pi\theta'}}, \quad R = \varepsilon + \frac{\varepsilon^{-1} - \varepsilon}{1 - e^{\pi\theta'}}.$$

Since we have

$$\frac{1 - \varepsilon r}{r} = \frac{1 - \varepsilon^2}{r} \left(1 - \frac{1}{1 + e^{\pi\theta'}}\right) > 0$$

and

$$\varepsilon^{-1}r - \varepsilon R = (1 - \varepsilon^2) \left(1 + \frac{\varepsilon^{-2}}{1 + e^{\pi\theta'}} - \frac{1}{1 - e^{\pi\theta'}}\right) > 0,$$

where the last inequality holds thanks to (1), there is a positive number s satisfying

$$0 < s < \min \left\{ r, \frac{1 - \varepsilon r}{r}, \frac{\varepsilon^{-1}r - \varepsilon R}{\varepsilon^{-1} + R} \right\},$$

which implies

$$(2) \quad (\varepsilon + s)r < 1, \quad (\varepsilon + s)R < \varepsilon^{-1}(r - s).$$

Take θ_1 so large that

$$(3) \quad \varepsilon + \frac{\varepsilon^{-1} - \varepsilon}{1 + e^{\pi(\theta_0 + \theta_1)}} < \varepsilon + s.$$

Now set $\theta_2 := \theta_0 + \theta_1 - \theta'$ and

$$a = \frac{1}{2} - \frac{i}{2}(\theta_0 + \theta_1 + \theta_2),$$

and deform (the real part of) b as

$$b(t) = \frac{1}{2} + t - \frac{i}{2}(\theta_0 + \theta_1 - \theta_2) = \frac{1}{2} + t - \frac{i}{2}\theta'.$$

Note that

$$\varepsilon < g(a) = \varepsilon + \frac{\varepsilon^{-1} - \varepsilon}{1 + e^{\pi(\theta_0 + \theta_1 + \theta_2)}} < \varepsilon + s \quad (\text{here we used (3)}),$$

FIGURE 4. The disk D_α travels around the disk D_1

and that

$$g(b(t)) = \varepsilon + \frac{\varepsilon^{-1} - \varepsilon}{1 + e^{\pi\theta'} e^{2\pi it}} \quad \text{satisfies} \quad r \leq |g(b(t))| \leq R.$$

These inequalities together with (2) imply that $\alpha(t) = g(a)g(b(t))$ is in the ring $\{\varepsilon r \leq |z| \leq (\varepsilon + s)R\}$, and that

$$\alpha(0) = \alpha(1) \leq (\varepsilon + s)r < 1 < \varepsilon R \leq \alpha(1/2).$$

This shows that $\alpha(t)$ travels around 1. Let D_0 be the disk with center at 0 and with radius $\varepsilon(r - s)$, D_∞ the complementary disk of the disk with center at 0 and with radius $\varepsilon^{-1}(r - s)$. Then γ_1 maps D_0 onto the complementary disk of D_∞ , and by the right side of (2), $D_0 \cup D_\infty$ is disjoint from the ring

$$\{\varepsilon r \leq |z| \leq (\varepsilon + s)R\} \ni 1, \alpha(t).$$

Thus by taking $\theta_1 > 0$ sufficiently large, for any $0 \leq t \leq 1$, there are two disjoint disks $D_1 (\ni 1)$, $D_{\alpha(t)} (\ni \alpha(t))$ in the complement of $D_0 \cup D_\infty$ and a transformation $\gamma_2(t)$ with multiplier $e^{2\pi i\mu} = e^{-2\pi\theta_1} e^{-2\pi it}$ which maps D_1 onto the complementary disk of the $D_{\alpha(t)}$ (see Figure 4).

5.3. The multiplier of γ_2 travels around 0. We construct a loop in S (with base in S_0), which is represented by the change of argument (by 2π) of the multiplier of γ_2 . We fix b and c , and the imaginary part of a , and let the real part of a move from $1/2$ to $3/2$. Set

$$\theta := \theta_0 + \theta_1 + \theta_2, \quad \varepsilon := e^{-\pi\theta_0}, \quad r := \frac{\varepsilon^{-1} - \varepsilon}{e^{\pi\theta} - 1}.$$

Choose and fix θ_0, θ_1 and θ_2 so that $\theta_1 = \theta_2$ and

$$r < \min \left\{ 1 - \varepsilon, \frac{1}{1 + \varepsilon^{-1}} \right\}.$$

Then since

$$b = \frac{1}{2} - \frac{i}{2}(\theta_0 + \theta_1 - \theta_2) = \frac{1}{2} - \frac{i}{2}\theta_0,$$

we have $e^{2\pi ib} = -\varepsilon^{-1}$, and so $g(b) = 1$.

Now we deform the parameter a as

$$a(t) = \frac{1}{2} + t - \frac{i}{2}\theta, \quad 0 \leq t \leq 1.$$

FIGURE 5. The fixed point α of γ_2 travels along the Apollonius circle

Since we have

$$\alpha(t) = g(a(t))g(b) = g(a(t)) = \varepsilon + \frac{\varepsilon^{-1} - \varepsilon}{1 + e^{\pi\theta} e^{2\pi it}},$$

the point α is in the disk with center at ε and with radius r . We thus have

$$\varepsilon^2(1+r) < \varepsilon - r \leq |\alpha(t)| \leq \varepsilon + r < 1 < 1+r.$$

Let D_0 be the disk with center at 0 and with radius $\varepsilon^2(1+r)$, D_∞ the complementary disk of the disk with center at 0 and with radius $1+r$, $D_{\alpha(t)}$ ($\ni \alpha(t)$) the disk with center at ε and with radius r . Then γ_1 maps D_0 onto the complementary disk of D_∞ , and $\alpha(t)$ belongs to the disk $\{|z - \varepsilon| \leq r\}$ which is disjoint from $D_0 \cup D_\infty \cup \{1\}$. Thus by taking $\theta_1 = \theta_2 > 0$ sufficiently large, for any $0 \leq t \leq 1$, there are two disjoint disks D_1 ($\ni 1$), $D_{\alpha(t)}$ ($\ni \alpha(t)$) in the complement of $D_0 \cup D_\infty$ and a transformation $\gamma_2(t)$ with multiplier $e^{2\pi i\mu} = e^{-2\pi\theta_1} e^{-2\pi it}$ which maps D_1 onto the complementary disk of $D_{\alpha(t)}$ (see Figure 5).

5.4. The multiplier of γ_1 travels around 0. We construct a loop in S (with base in S_0), which is represented by the change of argument (by 2π) of the multiplier of γ_1 . Note that by the change of variable $x \rightarrow 1 - x$, the equation $E(a, b, c)$ changes into $E(a, b, a + b + 1 - c)$. So we have only to literally follow §5.3 exchanging c and $a + b + 1 - c$, and θ_0 and θ_1 . We thus fix b and the imaginary parts of a and c , and let the real part of c move from 1 to 2, and let the real part of a move from 1/2 to 3/2 keeping $a + b + 1 - c$ constant.

6. MISCELLANEA

For a Schottky group Γ of genus 2, the quotient of the domain of discontinuity modulo Γ is a curve of genus 2. A curve of genus 2 is a double cover of $\mathbb{P}^1$ branching at six points, which are uniquely determined modulo automorphisms of $\mathbb{P}^1$ by the curve. Thus a Schottky group determines a point of the configuration space $X\{6\}$ of six-point sets on $\mathbb{P}^1$. (The fundamental group of $X\{6\}$ is the Braid group with five strings.) When all the exponent-differences of the hypergeometric equation are purely imaginary, its monodromy group is a Schottky group, which determines six points on a line. Thus they determine a point of the configuration space $X(6)$ of six colored points on $\mathbb{P}^1$. (The fundamental group of $X(6)$ is the colored Braid group with five strings.) The space $X(6)$ is well studied.

Remark 2. When all the exponent-differences are purely imaginary, there is a Schottky automorphic function defined by an absolutely convergent infinite product, which induces a holomorphic map of the genus 2 curve onto $\mathbb{P}^1$ ([IY]). This infinite product remains to be convergent for groups represented by loops above if we take the multipliers of γ_1 and γ_2 sufficiently small. This is because there is a circle separating two disks among the four ([BBEIM, Chapter 5]).

Problems. The four loops constructed in §5 induce those in $X\{6\}$. Do they generate the fundamental group of $X\{6\}$? Same problem for $X(6)$.

ACKNOWLEDGEMENTS

The authors are grateful to Professors Susumu Hirose, Ken'ichi Ohshika, Hiroki Sato and Hiro-o Yamamoto for their encouragement.

REFERENCES

- [BBEIM] E. D. BELOKOLOS, A. I. BOBENKO, V. Z. ENOL'SKII, A. R. ITS AND V. B. MATVEEV, *Algebro-geometric Approach to Nonlinear Integrable Equations*, Springer Series in Nonlinear Dynamics (Springer-Verlag, 1994).
- [GP] L. GERRITZEN AND M. VAN DER PUT, *Schottky groups and Mumford curves*, Lect. Notes in Math. **817**(1980) Springer. MR0590243 (82j:10053)
- [IY] T. ICHIKAWA AND M. YOSHIDA, On Schottky groups arising from the hypergeometric equation with imaginary exponents, Proc AMS 132(2003), 447–454. MR2022368
- [IKSY] K. IWASAKI, H. KIMURA, S. SHIMOMURA AND M. YOSHIDA, *From Gauss to Painlevé – A modern theory of special functions*, Vieweg Verlag, Wiesbaden, 1991. MR1118604 (92j:33001)
- [Gray] J. GRAY, *Linear differential equations and group theory from Riemann to Poincaré*, Birkhäuser, 2000. MR1751835 (2000m:34002)
- [MSW] D. MUMFORD, C. SERIES AND D. WRIGHT, *Indra's Pearls*, Cambridge Univ. Press, 2002. MR1913879 (2003f:00005)
- [SY] T. SASAKI AND M. YOSHIDA, A geometric study of the hypergeometric function with imaginary exponents, Experimental Math. **10**(2000), 321–330. MR1917420 (2003f:33002)
- [Sch] F. SCHOTTKY, Über eine specielle Function, welche bei einer bestimmten linearen Transformation ihres Arguments unverändert bleibt, J. Reine Angew. Math. 101 (1887) 227–272.

DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE AND ENGINEERING, SAGA UNIVERSITY,
SAGA 840-8502, JAPAN

E-mail address: ichikawa@ms.saga-u.ac.jp

DEPARTMENT OF MATHEMATICS, KYUSHU UNIVERSITY, FUKUOKA 810-8560, JAPAN

E-mail address: myoshida@math.kyushu-u.ac.jp