A SHORT PROOF OF A CONJECTURE ON THE CONNECTIVITY OF GRAPH COLORING COMPLEXES

ALEXANDER ENGSTRÖM

(Communicated by Paul Goerss)

ABSTRACT. The Hom -complexes were introduced by Lovász to study topological obstructions to graph colorings. It was conjectured by Babson and Kozlov, and proved by Čukić and Kozlov, that $\text{Hom}(G, K_n)$ is $(n - d - 2)$ -connected, where d is the maximal degree of a vertex of G , and n the number of colors. We give a short proof of the conjecture.

INTRODUCTION

It was conjectured by Babson and Kozlov [1], and proved by Čukić and Kozlov [4], that $\text{Hom}(G, K_n)$ is $(n - d - 2)$ -connected, where d is the maximal degree of a vertex of G , and n the number of colors. We give a shorter proof of this, by generalizing the proof of that $\text{Hom}(K_m, K_n)$ is $(n - m - 1)$ -connected in Babson and Kozlov [1].

For definitions and basic theorems on Hom -complexes used in this text, see the papers mentioned above, or the survey by Kozlov [6].

1. AN ANALOGUE OF THE CHROMATIC NUMBER

An independent subset of vertices of a graph is a set, such that no vertices of it are adjacent. The minimal number of sets needed to partition the vertex set of a graph G into independent sets is the chromatic number $\chi(G)$.

Definition 1.1. A *covering* $I_1, I_2, \dots, I_k$ of G is a sequence of independent subsets of $V(G)$ such that they partition $V(G)$, and I_i is a maximal independent set in the induced subgraph of G with vertex set $I_i \cup I_{i+1} \cup \dots \cup I_k$, for all i , where $1 \leq i \leq k$.

A partition of G into $\chi(G)$ independent sets can always be transformed to a covering by ordering the independent sets and if needed enlarging them. But a covering can use more than $\chi(G)$ sets. Define $\dot{\chi}(G)$ to be the maximal number of sets in a covering of G . Clearly, $\dot{\chi}(G) \geq \chi(G)$.

Lemma 1.2. *If d is the maximal degree of a vertex of G , then $\dot{\chi}(G) \leq d + 1$.*

Received by the editors May 31, 2005 and, in revised form, July 6, 2005.

2000 *Mathematics Subject Classification.* Primary 57M15, 05C15.

Key words and phrases. Graph homomorphisms, k -connectivity, Hom -complexes, graph coloring complexes.

This research was supported by ETH and Swiss National Science Foundation Grant PP002-102738/1.

©2006 American Mathematical Society
 Reverts to public domain 28 years from publication

Proof. Let $I_1, I_2, \dots, I_{\dot{\chi}(G)}$ be a covering of G , and $v \in I_{\dot{\chi}(G)}$. For each i , where $1 \leq i < \dot{\chi}(G)$, there is a $w \in I_i$ adjacent to v , because otherwise I_i would not be a maximal independent set. Hence the degree of v is at least $\dot{\chi}(G) - 1$. The degree of v is at most d , thus $\dot{\chi}(G) \leq d + 1$. $\square$

Lemma 1.3. *If H is an induced subgraph of G , then $\dot{\chi}(H) \leq \dot{\chi}(G)$.*

Proof. It suffices to prove this when H and G only differ by a vertex v of G . Let $I_1, I_2, \dots, I_{\dot{\chi}(H)}$ be a covering of H . If v is adjacent to a vertex in each of the sets I_i , then $\{v\}, I_1, I_2, \dots, I_{\dot{\chi}(H)}$ is a covering of G and $\dot{\chi}(H) + 1 \leq \dot{\chi}(G)$. Otherwise, let I_j be the first set in the covering such that v is not adjacent to any vertex of I_j . Then $I_1, I_2, \dots, I_j \cup \{v\}, \dots, I_{\dot{\chi}(H)}$ is a covering of G , and $\dot{\chi}(H) \leq \dot{\chi}(G)$. $\square$

Lemma 1.4. *If I is a maximal independent set of G , then $\dot{\chi}(G) > \dot{\chi}(G \setminus I)$.*

Proof. Let $I_1, I_2, \dots, I_{\dot{\chi}(G \setminus I)}$ be a covering of $G \setminus I$. Then $I, I_1, I_2, \dots, I_{\dot{\chi}(G \setminus I)}$ is a covering of G with $1 + \dot{\chi}(G \setminus I)$ sets. $\square$

2. HIGHER CONNECTIVITY OF $\text{Hom}(G, K_n)$

Lemma 2.1. *If I is an independent set of G , and $I' \subset I$, then $\Delta = \{\eta \in \text{Hom}(G, K_n) \mid n \in \eta(i) \Rightarrow i \in I\}$ collapses onto $\Delta' = \{\eta \in \text{Hom}(G \setminus (I \setminus I'), K_n) \mid n \in \eta(i) \Rightarrow i \in I'\}$.*

Proof. It suffices to prove this when $I \setminus I' = \{v\}$. Let $\eta_1, \eta_2, \dots, \eta_k$ be an ordering of $\{\eta \in \Delta \mid n \notin \eta(v)\}$ such that if $\eta(w) \supseteq \eta'(w)$, for all $w \in V(G)$, then η is not after η' . Define η_i^* as $\eta_i^*(w) = \eta_i(w)$ for $w \neq v$, and $\eta_i^*(v) = \eta_i(v) \cup \{n\}$. Each successive removal of η_i^* together with η_i from Δ for $i = 1, 2, \dots, k$ is a collapse step. The cells left are $\Delta'' = \{\eta \in \Delta \mid \eta(v) = \{n\}\}$. Finally, there is a bijection between the face posets of Δ' and Δ'' by extending each $\eta \in \Delta'$ with $\eta(v) = \{n\}$. $\square$

The main use of Lemma 2.1 is when $I' = \emptyset$. Then $n \notin \eta(w)$ for all $\eta \in \Delta'$ and $w \in V(G) \setminus I$, so $\Delta' = \text{Hom}(G \setminus I, K_{n-1})$. Another way to prove the lemma is to use discrete Morse theory [5].

Lemma 2.2 (Nerve Lemma, [2, Theorem 10.6(ii)], [3]). *Let Δ be a regular cell complex, and $(\Delta_j)_{j \in J}$ a family of subcomplexes such that $\Delta = \bigcup_{j \in J} \Delta_j$ and every nonempty finite intersection $\Delta_{j_1} \cap \Delta_{j_2} \cap \dots \cap \Delta_{j_t}$ is $(m - t + 1)$ -connected. Then Δ is m -connected if and only if the nerve $\mathcal{N}(\Delta_j)$ is m -connected.*

Assume that $m \geq 0$. We will construct a family of subcomplexes such that $\Delta = \bigcup_{j \in J} \Delta_j$, all Δ_j are m -connected, and every intersection $\Delta_{j_1} \cap \Delta_{j_2} \cap \dots \cap \Delta_{j_t}$ is $(m - 1)$ -connected for $t \geq 2$. If a complex is $(m - 1)$ -connected, then it is $(m - t + 1)$ -connected for $t \geq 2$. Since all intersections are nonempty, the nerve is a simplex, and Δ is m -connected.

Theorem 2.3. *$\text{Hom}(G, K_n)$ is $(n - \dot{\chi}(G) - 1)$ -connected.*

Proof. We use induction on $\dot{\chi}(G)$ and on $n - \dot{\chi}(G)$. When $\dot{\chi}(G) = 1$, G has no edges, so $\text{Hom}(G, K_n)$ is contractible, and in particular, $(n - \dot{\chi}(G) - 1)$ -connected. If $n - \dot{\chi}(G) = 0$, then $n \geq \chi(G)$ so $\text{Hom}(G, K_n)$ is nonempty, and $(n - \dot{\chi}(G) - 1)$ -connected.

For all $I \in \mathcal{I}$, let $\Delta_I = \{\eta \in \text{Hom}(G, K_n) \mid n \in \eta(i) \Rightarrow i \in I\}$, where $\mathcal{I}$ is the family of maximal independent subsets of G . Clearly, $\text{Hom}(G, K_n) = \bigcup_{I \in \mathcal{I}} \Delta_I$. By

Lemma 2.1, the complex Δ_I is homotopy equivalent to $\text{Hom}(G \setminus I, K_{n-1})$, which is $((n-1) - (\dot{\chi}(G) - 1) - 1)$ -connected by Lemma 1.4 and induction. If $\mathcal{I} \supseteq \mathcal{I}' \neq \emptyset$, then $\bigcap_{I \in \mathcal{I}'} \Delta_I = \{\eta \in \text{Hom}(G, K_n) \mid n \in \eta(i) \Rightarrow i \in \bigcap_{I \in \mathcal{I}'} I\}$ is homotopy equivalent to $\text{Hom}(G \setminus (\bigcap_{I \in \mathcal{I}'} I), K_{n-1})$ by Lemma 2.1, and $((n-1) - \dot{\chi}(G) - 1)$ -connected by Lemma 1.3 and induction. By the Nerve Lemma we are done. $\square$

Corollary 2.4. $\text{Hom}(G, K_n)$ is $(n - d - 2)$ -connected.

Proof. Lemma 1.2 states that $\dot{\chi}(G) \leq d + 1$. $\square$

ACKNOWLEDGEMENTS

We thank Sonja Čukić, Dmitry Kozlov, and the referee for their comments.

REFERENCES

1. E. Babson, D.N. Kozlov, *Complexes of graph homomorphisms*. Israel J. Math. **152** (2006), 285–312.
2. A. Björner, *Topological Methods*, in: “Handbook of Combinatorics” (eds. R. Graham, M. Grötschel, and L. Lovász), North-Holland, 1995, 1819–1872. MR1373690 (96m:52012)
3. A. Björner, L. Lovász, S.T. Vrećica, R.T. Živaljević, *Chessboard complexes and matching complexes*, J. London Math. Soc. (2) **49** (1994), 25–49. MR1253009 (95c:52021)
4. S. Lj. Čukić, D.N. Kozlov, *Higher connectivity of graph coloring complexes*, Int. Math. Res. Notices. **2005:25** (2005), 1543–1562. MR2152894
5. R. Forman, *Morse theory for cell complexes*, Adv. Math. **134**, no. 1, (1998), 90–145. MR1612391 (99b:57050)
6. D.N. Kozlov, *Chromatic numbers, morphism complexes, and Stiefel-Whitney characteristic classes*, 67 pages, to appear in *Geometric Combinatorics*, IAS/Park City Mathematics Series **14**, American Mathematical Society, Providence, RI; Institute for Advanced Study (IAS), Princeton, NJ.

DEPARTMENT OF COMPUTER SCIENCE, Eidgenössische Technische Hochschule, ZÜRICH, SWITZERLAND

E-mail address: engstroa@inf.ethz.ch