CONTINUOUSLY EXTENDING PARTIAL FUNCTIONS

PHILLIP ZENOR

(Communicated by Alan Dow)

ABSTRACT. We characterize those Hausdorff spaces in which continuous functions defined on compact subsets can be continuously extended to continuous functions defined on the space.

1. INTRODUCTION

$C(X)$ will denote the set of continuous real valued functions defined on the space X . $\mathcal{T}_{norm}$ will denote the norm topology on $C(X)$, while $\mathcal{T}_{co}$ will denote the compact-open topology on $C(X)$. Let $K(X)$ denote the space of compact subsets of X endowed with the Vietoris topology.¹ Each function in $\mathcal{C}_K(X) = \{f \in C(H) : H \in K(X)\}$ is identified with its graph so that $\mathcal{C}_K(X)$ is a subspace of $K(X \times \mathcal{R})$. The space $\mathcal{C}_K(X)$ was first studied by Kuratowski in [5], [6]. In [8], the author shows that if X is a compact metric space, then there is a continuous function $e : \mathcal{C}_K(X) \rightarrow (C(X), \mathcal{T}_{norm})$ such that $ef(x) = f(x)$ for all $x \in \text{Domain}(f)$. (We will call such a function e an extender.) In [7], Stepanova defines a separating function to be a function $\varphi : X^2 \setminus \Delta \rightarrow C(X)$ such that $\varphi(x, y)(x) \neq \varphi(x, y)(y)$ and proves that the following conditions are equivalent for a paracompact p-space X :

- (1) X admits a continuous separating function $\varphi : X^2 \setminus \Delta \rightarrow (C(X), \mathcal{T}_{norm})$,
- (2) there is a continuous extender $e : \mathcal{C}_K(X) \rightarrow (C(X), \mathcal{T}_{norm})$, and
- (3) X is metrizable.

(Paracompact p-spaces can be characterized as those spaces that admit perfect maps onto metric spaces [1].)

In [4], it is shown that if X is metrizable, then there is a continuous extender $e : \mathcal{C}_K(X) \rightarrow (C(X), \mathcal{T}_{norm})$ that is linear on functions with common domains.

Definition 1. A function $f : X \rightarrow C(Y)$ can be thought of as a function $\varphi_f : X \times Y \rightarrow \mathcal{R}$, defined by $\varphi_f(x, y) = [f(x)](y)$. We will say that $f : X \rightarrow C(Y)$ is naturally continuous if $\varphi_f : X \times Y \rightarrow \mathcal{R}$ is continuous. Unless otherwise stated, we will think of a separating function for X as a function $\varphi : (X^2 \setminus \Delta) \times X \rightarrow \mathcal{R}$ such that $\varphi(x, y, x) \neq \varphi(x, y, y)$.

Received by the editors January 1, 2004 and, in revised form, July 15, 2005.

2000 *Mathematics Subject Classification.* Primary 54D15, 54C20, 54C30.

¹If $\{U_1, \dots, U_n\}$ is a finite collection of open subsets of X , then the set $\langle U_1, \dots, U_n \rangle = \{H \in K(X) : H \subset \bigcup_{i=1}^n U_i, \text{ and if } 1 \leq i \leq n, \text{ then } H \cap U_i \neq \emptyset\}$ is a basic open set for the Vietoris topology on $K(X)$.

©2006 American Mathematical Society
 Reverts to public domain 28 years from publication

Clearly, if $f : X \rightarrow (C(Y), \mathcal{T}_{norm})$ is continuous, then $\varphi_f : X \times Y \rightarrow \mathcal{R}$ is continuous. We will see in Proposition 1 that if $\varphi_f : X \times Y \rightarrow \mathcal{R}$ is continuous, then $f : X \rightarrow (C(Y), \mathcal{T}_{co})$ is continuous.

In [3], X is defined to be continuously Urysohn if X admits a continuous separating function $f : X^2 \setminus \Delta \rightarrow (C(X), \mathcal{T}_{norm})$.

Definition 2. We will say that X is weakly continuously Uryshon if X admits a continuous separating function $\varphi : (X^2 \setminus \Delta) \times X \rightarrow \mathcal{R}$.

The main result of this paper is the following.

Theorem 1. *The following conditions are equivalent on the Hausdorff space X :*

- (1) X is weakly continuously Urysohn,
- (2) there is a naturally continuous extender $e : \mathcal{C}_K(X) \rightarrow C(X)$, and
- (3) there is a naturally continuous extender $e : \mathcal{C}_K(X) \rightarrow C(X)$ such that if $\text{Domain}(f) = \text{Domain}(g)$ and $f(x) \leq g(x)$ for all $x \in \text{Domain}(f)$, then $ef(x) \leq eg(x)$ for all $x \in X$.

2. BACKGROUND, CONSTRUCTIONS AND PROOFS

If $U \subset Y$ and $V \subset \mathcal{R}$, then we will let $[U, V] = \{f \in C(Y) : f(U) \subset V\}$.

Proposition 1. *A function $f : X \rightarrow (C(Y), \mathcal{T}_{co})$ is continuous provided that the function $\varphi_f : X \times Y \rightarrow \mathcal{R}$ is continuous.*

Proof. Suppose that $\varphi_f : X \times Y \rightarrow \mathcal{R}$ is continuous, $x \in X$ and $[K, W]$ is a subbasic member of $\mathcal{T}_{co}$ containing $f(x)$. For each $k \in K$, there are open sets U_k in X and V_k in Y such that $x \in U_k$, $k \in V_k$, and $\varphi_f(U_k \times V_k) \subset W$. There is a finite subset $\{k_1, \dots, k_n\}$ of K such that $\{V_{k_1}, \dots, V_{k_n}\}$ covers K . It follows that if $x' \in \cap\{U_{k_1}, \dots, U_{k_n}\}$, then $f(x') \in [K, W]$.

Proposition 2. *If Y is locally compact, then $f : X \rightarrow (C(Y), \mathcal{T}_{co})$ is continuous if and only if $\varphi_f : X \times Y \rightarrow \mathcal{R}$ is continuous.*

Proof. Suppose that $f : X \rightarrow (C(Y), \mathcal{T}_{co})$ is continuous. Let W be an open set in $\mathcal{R}$ containing $\varphi_f(x, y) = f(x)(y)$. Since $f(x) \in C(Y)$, we may let V be an open set containing y with compact closure such that $f(x) \in [\overline{V}, W]$. Since f is continuous, there is an open set U in X containing x such that if $x' \in U$, then $f(x') \in [\overline{V}, W]$. Thus, if $(x', y') \in U \times V$, then $\varphi_f(x', y') = f(x')(y') \in W$. It follows that φ_f is continuous.

Recall that if X is compact, then the compact-open topology and the norm topology on $C(X)$ are the same topologies. Thus we have

Theorem 2 (Stepanova [7]). *The compact space X is metrizable if and only if X is weakly continuously Urysohn.*

In [7], Stepanova points out that a submetrizable space admits a continuous separating function. Indeed, we have the following observation.

Proposition 3. *If X has a zero-set diagonal, then X is weakly continuously Urysohn.*

Proof. Let $\rho : X^2 \rightarrow [0, 1]$ be a continuous function such that $\Delta = \rho^{-1}(0)$. Since ρ is continuous,

$$\varphi(x, y, t) = \frac{1}{2} \left[\frac{-\rho(y, t) + \rho(x, t)(1 - \rho(y, t))}{\rho(x, t) + \rho(y, t)} + 1 \right]$$

is a continuous function from $(X^2 \setminus \Delta) \times X$ into $\mathcal{R}$ such that $\varphi(x, y, x) = 0$ and $\varphi(x, y, y) = 1$.

In [2], Bennett and Lutzer give us an example of a linearly ordered space that is continuously Uryshon but that does not have a G_δ -diagonal.

We let $\mathcal{M}(X) = \{(H, K) \in K(X) \times K(X) : H \cap K = \emptyset\}$ and $\mathcal{A}(X) = \{(x, K) \in X \times K(X) : x \notin K\}$.

Lemma 1 (Stepanova [7]). *If X admits a continuous separating function $\varphi : (X^2 \setminus \Delta) \times X \rightarrow \mathcal{R}$, then there is a continuous function $\varphi' : (X^2 \setminus \Delta) \times X \rightarrow [0, 1]$ such that $\varphi'(x, y, x) = 0$ and $\varphi'(x, y, y) = 1$.*

Lemma 2. *If X admits a continuous separating function $\varphi : (X^2 \setminus \Delta) \times X \rightarrow \mathcal{R}$, then there is a continuous function $\tilde{\varphi} : \mathcal{M}(X) \times X \rightarrow [0, 1]$ such that*

- (1) *if $x \in H$, then $\tilde{\varphi}(H, K, x) = 0$, and if $x \in K$, then $\tilde{\varphi}(H, K, x) = 1$, and*
- (2) *if (H, K) and $(H', K') \in \mathcal{M}(X)$ with $H \subset H'$ and $K' \subset K$, then $\tilde{\varphi}(H', K', x) \leq \tilde{\varphi}(H, K, x)$.*

Proof. By Lemma 1, we may assume that $0 \leq \varphi(x, y, t) \leq 1$, $\varphi(x, y, x) = 0$, and $\varphi(x, y, y) = 1$.

Define $\varphi_1 : \mathcal{A}(X) \times X \rightarrow \mathcal{R}$ by $\varphi_1(x, H, t) = \max\{\varphi(x, y, t) : y \in H\}$.

Clearly, $\varphi_1(x, H, x) = 0$. If $t \in H$, then $\varphi_1(x, H, t) = 1$, and if $H \subset K$, then $\varphi_1(x, H, t) \leq \varphi_1(x, K, t)$. We will show that φ_1 is continuous.

Let $\varepsilon > 0$ and let $W = (\varphi_1(x_0, K_0, t_0) - \varepsilon, \varphi_1(x_0, K_0, t_0) + \varepsilon)$. For each $y \in K_0$ there are open sets O_y , Λ_y , and Ω_y containing x_0 , y , and t_0 , respectively, such that if $x' \in O_y$, $y' \in \Lambda_y$, and $t' \in \Omega_y$, then $\varphi(x', y', t') \in (\varphi(x_0, y, t_0) - \varepsilon, \varphi(x_0, y, t_0) + \varepsilon)$. Since K_0 is compact, there is a finite subset $\{y_1, \dots, y_n\}$ of K_0 such that $\{\Lambda_{y_1}, \dots, \Lambda_{y_n}\}$ covers K . We may assume that $\varphi_1(x_0, K_0, t_0) = \varphi(x_0, y_1, t_0)$. Let $O = \bigcap_{i=1}^n O_{y_i}$, $\Lambda = \langle \Lambda_{y_1}, \dots, \Lambda_{y_n} \rangle$, and $\Omega = \bigcap_{i=1}^n \Omega_{y_i}$. Let $(x', K', t') \in O \times \Lambda \times \Omega$. Let $k \in K' \cap \Lambda_{y_1}$. Then $\varphi_1(x', K', t') \geq \varphi(x', k, t') \geq \varphi(x_0, k_1, t_0) - \varepsilon = \varphi_1(x_0, y_0, t_0) - \varepsilon$. Now, let $k \in K'$ be such that $\varphi(x', k, t') = \varphi_1(x', K', t')$. Then $k \in \Lambda_{y_i}$ for some $1 \leq i \leq n$. We have $\varphi(x', k, t') \leq \varphi(x_0, y_i, t_0) + \varepsilon \leq \varphi(x_0, y_1, t_0) + \varepsilon = \varphi_1(x_0, K_0, t_0) + \varepsilon$.

We now define $\tilde{\varphi}(H, K, t) = \min\{\varphi_1(x, K, t) : x \in H\}$. We need only show that $\tilde{\varphi}$ is continuous. To this end, let $\varepsilon > 0$. For each $x \in H$, there are open sets U_x in X containing x , V_x in $K(X)$ containing K , and Λ_x containing t such that if $(x', K', t') \in U_x \times V_x \times \Lambda_x$, $\varphi_1(x, K, t) - \varepsilon < \varphi_1(x', K', t') < \varphi_1(x, K, t) + \varepsilon$. There is a finite set $\{x_1, \dots, x_n\} \subset H$ such that $\{U_{x_1}, \dots, U_{x_n}\}$ covers H . We may assume that $\tilde{\varphi}(H, K, t) = \varphi_1(x_1, K, t)$. Let $U = \langle U_{x_1}, \dots, U_{x_n} \rangle$, $V = \bigcap_{i=1}^n V_{x_i}$, and $\Lambda = \bigcap_{i=1}^n \Lambda_{x_i}$. Let $(H', K', t') \in U \times V \times \Lambda$. There is an $x' \in H' \cap U_{x_1}$, and so, $\tilde{\varphi}(H', K', t') \leq \varphi_1(x', K', t') < \varphi_1(x_1, K, t) + \varepsilon = \varphi_1(H, K, t) + \varepsilon$. Now, let $x' \in H'$ be such that $\tilde{\varphi}(H', K', t') = \varphi_1(x', H', t')$. Then $x' \in U_{x_j}$ for some j . Thus, $\tilde{\varphi}(H', K', t') = \varphi_1(x', K', t') > \varphi_1(x_j, K, t) - \varepsilon \geq \tilde{\varphi}(H, K, t) - \varepsilon$.

Observation 1. It follows from Part 2 of Lemma 2 that if $(H, K, t) \in \mathcal{M}(X) \times X$, and $\varepsilon > 0$, then

- (1) there are open sets $U \times V$ in $\mathcal{M}(X)$ containing (H, K) and W containing t such that if $\{(A, B), (A', B')\} \subset \mathcal{M}(X)$ with $A' \subset A$, $A' \in U$, $B \subset B' \in V$, and $x \in W$, then $\varphi(A, B, x) < \varphi(H, K, t) + \varepsilon$ and
- (2) there are open sets $U' \times V'$ in $\mathcal{M}(X)$ containing (H, K) and W' containing t such that if $\{(A, B), (A', B')\} \subset \mathcal{M}(X)$ with $A \subset A' \in U'$, $B' \subset B$, $B' \in V'$ and $x \in W'$, then $\varphi(A, B, x) > \varphi(H, K, t) - \varepsilon$.

Let $\mathcal{C}_K^*(X)$ denote the set $\{f : H \rightarrow (0, 1) \mid H \in K(X), f \text{ is continuous}\}$ endowed with the Vietoris topology.

Lemma 3. Let $\varphi : \mathcal{M}(X) \times X \rightarrow [0, 1]$ be the function $\tilde{\varphi}$ given in Lemma 2. We extend φ so that if $H \neq \emptyset$, then $\varphi(\emptyset, H, x) = 1$ and $\varphi(H, \emptyset, x) = 0$ for all $x \in X$. Suppose that $f \in \mathcal{C}_K^*(X)$, and $0 < a < b < 1$.

- (1) If $f^{-1}[0, a] \neq \emptyset$, $\varphi(f^{-1}[0, a], f^{-1}[b, 1], x) < \zeta$, and $\varepsilon > 0$, then there are open sets Γ in $\mathcal{C}_K^*(X)$ containing f and O in X containing x such that if $a' < b'$, $a' > a + \varepsilon$, $b' > b + \varepsilon$, $g \in \Gamma$, and $x' \in O$, then

$$\varphi(g^{-1}[0, a'], g^{-1}[b', 1], x') < \zeta.$$

- (2) If $f^{-1}[b, 1] \neq \emptyset$, $\varphi(f^{-1}[0, a], f^{-1}[b, 1], x) > \zeta$, and $\varepsilon > 0$, then there are open sets Γ in $\mathcal{C}_K^*(X)$ containing f and O in X containing x such that if $a' < b'$, $a' < a - \varepsilon$, $b' < b - \varepsilon$, $g \in \Gamma$, and $x' \in O$, then

$$\varphi(g^{-1}[0, a'], g^{-1}[b', 1], x') > \zeta.$$

Proof of Lemma 3(1). Case I: $f^{-1}[b, 1] = \emptyset$. Let $c = \sup(f)$ and $\delta < \frac{1}{2}\min\{\varepsilon, b - c\}$. Then $\varphi(f^{-1}[0, c], f^{-1}[b, 1], x) = 0$ for all $x \in X$. Let $\langle \gamma_1, \dots, \gamma_k \rangle$ be an open set in $K(X)$ containing the domain of f such that $f^{-1}[0, a] \cap \gamma_1 \neq \emptyset$ and such that if $\{y, z\} \subset \gamma_i \cap \text{Domain}(f)$, then $|f(y) - f(z)| < \delta$. For each $i \leq k$, choose $x_i \in \gamma_i \cap \text{Domain}(f)$ such that $x_1 \in f^{-1}[0, a]$ and let $\gamma_i^* = \gamma_i \times (f(x_i) - \delta, f(x_i) + \delta)$. $\Gamma = \langle \gamma_1^*, \dots, \gamma_k^* \rangle$ is an open set in $\mathcal{C}_K(X)$ containing f .

Let $g \in \Gamma$. We will show that $\varphi(g^{-1}[0, a'], g^{-1}[b', 1], x') = 0$ for all $x' \in X$. First, $g^{-1}[b', 1] = \emptyset$. To see that this is true, let $(y, g(y)) \in \gamma_i^*$. Then $g(y) < f(x_i) + \delta \leq c + \delta < b + \varepsilon < b'$. Now, we show that $g^{-1}[0, a'] \neq \emptyset$. To this end, let $(y, g(y)) \in \gamma_1^*$. Then $g(y) < f(x_1) + \delta \leq a + \delta < a'$ and $g^{-1}[0, a'] \neq \emptyset$. We have that $\varphi(g^{-1}[0, a'], g^{-1}[b', 1], x') = 0$ for all $x' \in X$. Case I follows.

Case II: $f^{-1}[b, 1] \neq \emptyset$. According to Observation 1, since $f^{-1}[0, a] \cap f^{-1}[b, 1] = \emptyset$, there are mutually exclusive open sets in $K(X)$, $\mathcal{A} = \langle \alpha_1, \dots, \alpha_n \rangle$ containing $f^{-1}[0, a]$ and $\mathcal{B} = \langle \beta_1, \dots, \beta_n \rangle$ containing $f^{-1}[b, 1]$, and O , open in X containing x , such that if $(A', B', x') \in \mathcal{A} \times \mathcal{B} \times O$, $(A, B) \in \mathcal{M}(X)$, and $A' \subset A$, and $B \subset B'$, then $\varphi(A, B, x') < \zeta$.

Let $\delta > 0$ be such that $2\delta < \varepsilon$, $f^{-1}[0, a + 2\delta] \in \mathcal{A}$, and $f^{-1}[b - 2\delta, 1] \in \mathcal{B}$. Choose open sets in X , $\{\gamma_1, \dots, \gamma_{k_1}, \dots, \gamma_{k_2}, \dots, \gamma_k\}$, such that (1) $\text{Domain}(f) \in \langle \gamma_1, \dots, \gamma_k \rangle$, (2) $f^{-1}[0, a] \in \langle \gamma_1, \dots, \gamma_{k_1} \rangle \subset \mathcal{A}$, (3) $f^{-1}[b, 1] \in \langle \gamma_{k_2}, \dots, \gamma_{k_n} \rangle \subset \mathcal{B}$, and (4) if t_1 and t_2 are in $\text{Domain}(f) \cap \gamma_i$, then $|f(t_1) - f(t_2)| < \delta/2$.

Let $g \in \Gamma$. For each $i \leq k$, choose x_i in $\text{Domain}(f) \cap \gamma_i$ such that if $i \leq k_1$, then $f(x_i) \leq a$ and if $i \geq k_2$, then $f(x_i) \geq b$, and let $\gamma_i^* = \gamma_i \times (f(x_i) - \delta, f(x_i) + \delta)$. Then $\Gamma = \langle \gamma_1^*, \dots, \gamma_k^* \rangle$ is an open set in $\mathcal{C}_K(X)$ containing f . For each $i \leq k_1$,

let $y_i \in \text{Domain}(g)$ such that $(y_i, g(y_i)) \in \gamma_i^*$ and let $A' = \{y_1, \dots, y_{k_1}\}$. Then $g(y_i) < f(x_i) + \delta < a'$. Thus, $A' \subset g^{-1}[0, a']$ and $A' \in \mathcal{A}$. If $g^{-1}[b', 1] = \emptyset$, then $\varphi(g^{-1}[0, a'], g^{-1}[b', 1], x') = 0$ for all $x' \in X$, and we are done. So let $y \in g^{-1}[b', 1]$. There is an i such that $(y, g(y)) \in \gamma_i^*$. Since $f(x_i) > g(y) - \delta \geq b' - \delta > b$, $i \geq k_2$, and there is an integer j such that $y \in \beta_j$. Therefore, if we let $B' = \{x_{k_2}, \dots, x_k\} \cup g^{-1}[b', 1]$, then $g^{-1}[b', 1] \subset B' \in \mathcal{B}$. We have that if $g \in \Gamma$ and $x' \in O$, then

$$\varphi(g^{-1}[0, a'], g^{-1}[b', 1], x') < \zeta.$$

The proof of Lemma 3(2) follows in the same way.

A construction: Let $\varphi : \mathcal{M}(X) \times X \rightarrow [0, 1]$ be as given in Lemma 2. For convenience, if $H \neq \emptyset$, then we define $\varphi(\emptyset, H, x) = 1$, and $\varphi(H, \emptyset, x) = 0$. For $f \in \mathcal{C}_K^*(X)$, $z \in [0, 1]$, and $n \in \mathcal{N}$, define

- $\underline{f}_{n,z}(x) = \begin{cases} \varphi(f^{-1}[0, z], f^{-1}[z + 1/n, 1], x), & \text{if } f^{-1}[0, z] \neq \emptyset, \\ 1 & \text{otherwise,} \end{cases}$
- $\overline{f}_{n,z}(x) = \begin{cases} \varphi(f^{-1}[0, z - 1/n], f^{-1}[z, 1], x), & \text{if } f^{-1}[z, 1] \neq \emptyset, \\ 0 & \text{otherwise,} \end{cases}$
- $U_{f,n,z} = \{x \mid \underline{f}_{n,z}(x) < z/n\}$, and
- $V_{f,n,z} = \{x \mid \overline{f}_{n,z}(x) > 1 - 1/n + z/n\}$.

We define $W_{f,z} = \bigcup_{i \in \mathcal{N}} (U_{f,i,z} \setminus \overline{V}_{f,i,z})$. Then $W_{f,z}$ is an open set containing $f^{-1}[0, z]$ and $\overline{W}_{f,z} \cap f^{-1}(z, 1] = \emptyset$.

Note that (1) if $z_1 < z_2$, then $\underline{f}_{n,z_2}(x) \leq \underline{f}_{n,z_1}(x)$ and $\overline{f}_{n,z_2}(x) \leq \overline{f}_{n,z_1}(x)$. Thus, $\overline{U}_{f,n,z_1} \subset U_{f,n,z_2}$ and $\overline{V}_{f,n,z_2} \subset V_{f,n,z_1}$. (2) If $z \in [0, 1]$, then $\overline{U}_{f,n+1,z} \subset U_{f,n,z}$ and $\overline{V}_{f,n+1,z} \subset V_{f,n,z}$. (3) If $z > \sup(f)$, then $W_{f,z} = X$. (4) If $z < \inf(f)$, then $W_{f,z} = \emptyset$.

Lemma 4. Suppose that $0 < z_1 < z_2 < 1$ and $1/N < (z_2 - z_1)/3$. Then $V_{f,N,z_2} \cap U_{f,N,z_1} = \emptyset$.

Proof. We are given that $z_1 + 1/N < z_2 - 1/N$. If $f^{-1}[0, z_1] = \emptyset$, then $U_{f,N,z_1} = \emptyset$. If $f^{-1}[z_2, 1] = \emptyset$, then $V_{f,N,z_2} = \emptyset$. If $f^{-1}[0, z_1] \neq \emptyset$ and $f^{-1}[z_2, 1] \neq \emptyset$, then $\{\eta \mid \varphi(f^{-1}[0, z_1], f^{-1}[z_1 + 1/N, 1], \eta) < 1\} \subset \{\eta \mid \varphi(f^{-1}[0, z_2 - 1/N], f^{-1}[z_2, 1], \eta) = 0\}$. Thus, $\{\eta \mid \varphi(f^{-1}[0, z_1], f^{-1}[z_1 + 1/N, 1], \eta) < z_1/N\} \cap \{\eta \mid \varphi(f^{-1}[0, z_2 - 1/N], f^{-1}[z_2, 1], \eta) > 1 - 1/N + z_2/N\} = \emptyset$.

Lemma 5. If $z_1 < z_2$, then $\overline{W}_{f,z_1} \subset W_{f,z_2}$.

Proof. Suppose that $x \in \overline{W}_{f,z_1} \setminus W_{f,z_2}$. By Lemma 4, we may let N denote the first integer such that $x \notin U_{f,N,z_1} \cap V_{f,N,z_2}$. Since $x \in \overline{W}_{f,z_1}$, $x \in \overline{U}_{f,1,z_1}$, and $f^{-1}[0, z_1] \neq \emptyset$. But, $f^{-1}[z_1 + 1, 1] = \emptyset$; so, $U_{f,1,z_1} = X$. Since $x \in \overline{U}_{f,1,z_1} \subset U_{f,1,z_2}$ and $x \notin W_{f,z_2}$, it must be that $x \in \overline{V}_{f,1,z_2}$ and $f^{-1}[z_2, 1] \neq \emptyset$. Since $f^{-1}[0, z_2 - 1] = \emptyset$, $V_{f,1,z_2} = X$. Thus, $N > 1$.

If $x \in \overline{V}_{f,N,z_2}$, then $x \notin U_{f,N,z_1}$, $x \in V_{f,N,z_1}$, and $x \notin \overline{U}_{f,N+1,z_1}$. Then $x \in (V_{f,N,z_1} \setminus \overline{U}_{f,N+1,z_1})$ and $(V_{f,N,z_1} \setminus \overline{U}_{f,N+1,z_1}) \cap W_{f,z_1} = \emptyset$, which would be a contradiction. So, $x \notin \overline{V}_{f,N,z_2}$. Since $x \notin W_{f,z_2}$, $x \notin U_{f,N,z_2}$, and so, $x \notin \overline{U}_{f,N,z_1}$. But then $x \in (V_{f,N-1,z_1} \setminus \overline{U}_{f,N,z_1}) \cap W_{f,z_1} = \emptyset$, which is a contradiction from which the lemma follows.

Lemma 6. *If X is weakly continuously Urysohn, then there is a continuous function $e : \mathcal{C}_K^*(X) \times X \rightarrow [0, 1]$ such that*

- (1) $ef(x) = f(x)$ for all $x \in \text{Domain}(f)$, and
- (2) if f and g have a common domain and if $f(x) \leq g(x)$ for all $x \in \text{Domain}(f)$, then $ef(x) \leq eg(x)$ for all $x \in X$.

Proof. For $f \in \mathcal{C}_K^*(X)$ and $x \in X$, define $ef(x) = \text{glb}\{z \in [0, 1] \mid x \in W_{f,z}\}$. It follows from the construction that $ef : X \rightarrow [0, 1]$ is an extension of f such that if f and g share a common domain, then $f \leq g$ implies that $ef \leq eg$. Furthermore, the image of ef is contained in the convex hull of the image of f . It is a standard argument that ef is continuous. It remains to show that $e : (\mathcal{C}_K^*(X)) \times X \rightarrow [0, 1]$ is continuous. To this end, let $f \in \mathcal{C}_K^*(X)$, $x \in X$, and $\varepsilon > 0$. Let $z = ef(x)$. We will obtain an open set $\Gamma \times O$ in $\mathcal{C}_K^*(X) \times X$ containing (f, x) such that if $(g, x') \in \Gamma \times O$, then $|ef(x) - eg(x')| < \varepsilon$.

Part I: First, we obtain an open set $\Gamma_1 \times O_1$ in $\mathcal{C}_K^*(X) \times X$ containing (f, x) such that if $(g, x') \in \Gamma_1 \times O_1$, then $eg(x) \leq z + \varepsilon$; that is, if $1 > z_0 > z + \varepsilon$, then $x' \in W_{g,z_0}$. To this end, choose z' and z'' in $[0, 1]$ such that $z < z' < z'' < z + \varepsilon$. Since $x \in W_{f,z'}$, there is an $N > 0$ such that $x \in U_{f,N,z'} \setminus \overline{V}_{f,N,z'}$.

Step A: Let $\varepsilon' = \varepsilon + z - z''$. Since $V_{f,N,z'} \neq X$, $f^{-1}[0, z' - 1/N] \neq \emptyset$ and $f^{-1}[0, z'' - 1/N] \neq \emptyset$.

$$\begin{aligned} \varphi(f^{-1}[0, z'' - 1/N], f^{-1}[z'', 1], x) &\leq \varphi(f^{-1}[0, z' - 1/N], f^{-1}[z', 1], x) \\ &\leq 1 - 1/N + z'/N < 1 - 1/N + z''/N. \end{aligned}$$

By Lemma 3, there is an open set $\Lambda_1 \times W_1$ in $\mathcal{C}_K^*(X) \times X$ containing (f, x) such that if $(g, x') \in \Lambda_1 \times W_1$ and $z_0 > z'' + \varepsilon' = z + \varepsilon$, then $\varphi(g^{-1}[0, z_0 - 1/N], g^{-1}[z_0, 1], x') < 1 - 1/N + z'' < 1 - 1/N + z_0/N$ and $x' \notin \overline{V}_{g,N,z_0}$.

Step B: Let $\varepsilon' = \varepsilon + z - z'$. Since $f^{-1}[0, z' - 1/N] \neq \emptyset$, $f^{-1}[0, z'] \neq \emptyset$. Since $\varphi(f^{-1}[0, z'], f^{-1}[z' + 1/N, 1], x) < z'/N$, we may employ Lemma 3 to obtain an open set $\Lambda_2 \times U_2$ in $\mathcal{C}_K^*(X) \times X$ containing (f, x) such that if $z_0 > z' + \varepsilon' = z + \varepsilon$ and $(g, x') \in \Lambda_2 \times U_2$, then $\varphi(g^{-1}[0, z_0], g^{-1}[z_0 + 1/N, 1], x') < z'/N < z_0/N$ and $x' \in U_{g,N,z_0}$.

To complete the argument for Part I, let $\Gamma_1 = \Lambda_1 \cap \Lambda_2$ and $O_1 = U_1 \cap U_2$.

Part II: Second, we find an open set $\Gamma_2 \times O_2$ in $\mathcal{C}_K^*(X) \times X$ containing (f, x) such that if $(g, x') \in \Gamma_2 \times O_2$, then $eg(x') \geq z - \varepsilon$; that is, if $z_0 < z - \varepsilon$, then $x' \notin W_{g,z_0}$. We consider three cases.

Case A: $f^{-1}[0, z - \varepsilon] = \emptyset$.

Choose z' so that $z - \varepsilon < z' < z$ and $f^{-1}[0, z'] = \emptyset$. Set $2\varepsilon' = z' - (z - \varepsilon)$. Let $\{U_1 \dots U_n\}$ be an irreducible open cover of $f^{-1}[z', 1] = \text{Domain}(f)$ such that if $1 \leq i \leq n$ and $\{a, b\} \subset \text{Domain}(f) \cap U_i$, then $|f(a) - f(b)| < \varepsilon'$. For each $1 \leq i \leq n$, choose $x_i \in U_i \cap \text{Domain}(f)$ and let $U_i^* = U_i \times (x_i - \varepsilon', x_i + \varepsilon')$. Then $\Gamma_2 = \langle U_1^*, \dots, U_n^* \rangle$ is an open set containing f such that if $g \in \Gamma_2$, $z_0 < z' - \varepsilon' = z - \varepsilon$, and $x' \in X$, then $g^{-1}[0, z_0] = \emptyset$. Letting $O_2 = X$, we have an open set $\Gamma_2 \times O_2$ in $\mathcal{C}_K^*(X) \times X$ containing (f, x) such that if $(g, x') \in \Gamma_2 \times O_2$, then $eg(x') \geq z - \varepsilon$.

In both Cases B and C, we will choose z' , z'' and z''' in $(0, 1)$ such that $z > z' > z'' > z''' > z - \varepsilon$. Lemma 4 gives us a first integer N such that $x \notin U_{f,N,z''} \cap V_{f,N,z'}$.

Case B: $f^{-1}[0, z - \varepsilon] \neq \emptyset$ and $x \in \overline{V}_{f,N,z'}$. Then $x \notin U_{f,N,z''}$ so

$$\varphi(f^{-1}[0, z''], f^{-1}[z'' + 1/N, 1], x) \geq z''/N$$

and $f^{-1}[z'' + 1/N, 1] \neq \emptyset$. It follows that $f^{-1}[z''' + 1/N, 1] \neq \emptyset$ and

$$\varphi(f^{-1}[0, z'''], f^{-1}[z''' + 1/N, 1], x) > z''/N > z'''/N.$$

We let $\varepsilon' = z''' - (z - \varepsilon)$ and apply Lemma 3 to obtain an open set $\Omega_1 \times V_1$ in $\mathcal{C}_K^*(X) \times X$ containing (f, x) such that if $(g, x') \in \Omega_1 \times V_1$ and $z_0 < z''' - \varepsilon' = z - \varepsilon$, then

$$\varphi(g^{-1}[0, z_0], g^{-1}[z_0 + 1/N, 1], x') > z'''/N > z_0/N.$$

Now, since $x \in \overline{V}_{f,N,z'}$, $x \in V_{f,N,z''}$ and $\varphi(f^{-1}[0, z''' - 1/N], f^{-1}[z''', 1], x) > 1 - 1/N + z'''/N$. Letting $\varepsilon' = z''' - (z - \varepsilon)$, we apply Lemma 3 to obtain an open set $\Omega_2 \times V_2 \in \mathcal{C}_K^*(X) \times X$ containing (f, x) such that if $(g, x') \in \mathcal{C}_K^*(X) \times X$ and $z_0 < z''' - \varepsilon' = z - \varepsilon$, then $x' \notin U_{g,i,z_0}$ for all $i \geq N$.

To complete Case B, we let $\Gamma_2 = \Omega_1 \cap \Omega_2$ and $O_2 = V_1 \cap V_2$.

Case C: $f^{-1}[0, z - \varepsilon] \neq \emptyset$ and $x \notin \overline{V}_{f,N,z'}$. Then since $x \notin W_{f,z'}$, $x \notin U_{f,N,z'}$. Thus, $\varphi(f^{-1}[0, z'], f^{-1}[z' + 1/N, 1], x) \geq z'/N$. It follows that $f^{-1}[z' + 1/N, 1] \neq \emptyset$ and $\varphi(f^{-1}[0, z''], f^{-1}[z'' + 1/N, 1], x) \geq z'/N > z''/N$. Again, Lemma 3 gives us an open set $\Omega_1 \times V_1 \in \mathcal{C}_K^*(X) \times X$ such that if $(g, x') \in \Omega_1 \times V_1$, then $x' \notin U_{g,N,z_0}$.

Since $f^{-1}[0, z - \varepsilon] \neq \emptyset$, $f^{-1}[0, z''] \neq \emptyset$, and $U_{f,1,z''} = X$. Similarly, since $f^{-1}[z' + 1/N, 1] \neq \emptyset$, $V_{f,1,z'} = X$. Therefore, $N > 1$. By hypothesis, $x' \in V_{f,N-1,z'} \subset V_{f,N-1,z''}$. Letting $\varepsilon' = z'' - (z - \varepsilon)$, we employ Lemma 3 to obtain an open set $\Omega_2 \times V_2$ in $\mathcal{C}_K^*(X) \times X$ such that if $(g, x') \in \Omega_2 \times V_2$ and $z_0 < z'' - \varepsilon' = z - \varepsilon$, then $z' \notin W_{g,z_0}$.

To complete Case C, we let $\Gamma_2 = \Omega_1 \cap \Omega_2$ and $O_2 = V_1 \cap V_2$. To complete the proof of the lemma, we let $\Gamma = \Gamma_1 \cap \Gamma_2$ and $O = O_1 \cap O_2$.

We are now in a position to obtain our main result.

Theorem 1. *If X is a Hausdorff space, then the following conditions are equivalent:*

- (1) X is weakly continuously Urysohn.
- (2) There is a continuous extender $e : \mathcal{C}_K(X) \times X \rightarrow \mathcal{R}$.
- (3) There is a continuous extender $e : \mathcal{C}_K(X) \times X \rightarrow \mathcal{R}$ such that if f and g have a common domain and if $f(x) \leq g(x)$ for all $x \in \text{Domain}(f)$, then $ef(x) \leq eg(x)$ for all $x \in X$.

Proof. Clearly, (3) $\Rightarrow$ (2). We will use a standard argument to show that (1) $\Rightarrow$ (3). Let $e^* : \mathcal{C}_K^*(X) \times X \rightarrow [0, 1]$ be the extender given in Lemma 6. Let $h : \mathcal{R} \rightarrow (0, 1)$ be an order preserving homeomorphism taking $\mathcal{R}$ onto $(0, 1)$. For $f \in \mathcal{C}_K(X)$, define $ef = h^{-1} \circ e^* \circ h \circ f$. To obtain (2) $\Rightarrow$ (1), for each $(x, y) \in X^2 \setminus \Delta$ let $\psi_{(x,y)} : \{x, y\} \rightarrow [0, 1]$ be defined by $\psi_{(x,y)}(x) = 0$ and $\psi_{(x,y)}(y) = 1$. Let $\varphi(x, y, t) = e\psi_{(x,y)}(t)$.

The author would like to express his appreciation to the referees for their patience with the initial versions of this paper.

REFERENCES

- [1] A. V. Arhangel'skiĭ, *On a class of spaces containing all metric and all locally bicomact spaces* (Russian) Dokl. Akad. Nauk SSSR 151 (1963), 751–754. MR0152988 (27:2959)
- [2] Harold R. Bennett and David J. Lutzer, *Continuous separating families in ordered spaces and strong base conditions*, Topology Appl. 119 (2002), no. 3, 305–314. MR1888675 (2002m:54034)
- [3] Lorenz Halbeisen and Norbert Hungerbühler, *On continuously Urysohn and strongly separating spaces*, Topology Appl. 118 (2002), no. 3, 329–335. MR1874554 (2002j:54041)

- [4] H. P. Künzi and L. B. Shapiro, *On simultaneous extension of continuous partial functions*, Proc. Amer. Math. Soc. 125 (1997), 1853-1859. MR1415348 (98g:54015)
- [5] K. Kuratowski, *Sur l'espace des fonctions partielles*, Ann. Mat. Pura Appl. 40 (1955), 61-67. MR0074807 (17:650b)
- [6] K. Kuratowski, *Sur une méthode de métrisation complète de certains espaces d'ensembles compacts*, Fund. Math. 43 (1956), 114-138. MR0079258 (18:58a)
- [7] E. N. Stepanova, *Continuation of continuous functions and the metrizability of paracompact p -spaces* (Russian), Mat. Zametki 53 (1993), no. 3, 92-101; translation in Math. Notes 53 (1993), no. 3-4, 308-314. MR1220188 (94k:54031)
- [8] P. L. Zenor, *Extending continuous functions in compact metric spaces*, Topology Conference (Virginia Polytech. Inst. and State Univ., Blacksburg, Va., 1973), pp. 277-283. Lecture Notes in Math., Vol. 375, Springer, Berlin, 1974. MR0355950 (50:8423)

DEPARTMENT OF MATHEMATICS, AUBURN UNIVERSITY, ALABAMA 36849