EXACT MULTIPLICITY RESULT FOR THE PERTURBED SCALAR CURVATURE PROBLEM IN $\mathbb{R}^N$ ($N \geq 3$)

S. PRASHANTH

(Communicated by David S. Tartakoff)

ABSTRACT. Let $D^{1,2}(\mathbb{R}^N)$ denote the closure of $C_0^\infty(\mathbb{R}^N)$ in the norm $\|u\|_{D^{1,2}(\mathbb{R}^N)}^2 = \int_{\mathbb{R}^N} |\nabla u|^2$. Let $N \geq 3$ and define the constants $\alpha_N = N(N-2)$ and $C_N = (N(N-2))^{\frac{N-2}{4}}$. Let $K \in C^2(\mathbb{R}^N)$. We consider the following problem for $\varepsilon \geq 0$:

$$(P_\varepsilon) \quad \left\{ \begin{array}{l} \text{Find } u \in D^{1,2}(\mathbb{R}^N) \text{ solving :} \\ -\Delta u = \alpha_N(1 + \varepsilon K(x))u^{\frac{N+2}{N-2}}, \\ u > 0 \end{array} \right\} \text{ in } \mathbb{R}^N.$$

We show an exact multiplicity result for (P_ε) for all small $\varepsilon > 0$.

1. INTRODUCTION

Let $D^{1,2}(\mathbb{R}^N)$ denote the closure of $C_0^\infty(\mathbb{R}^N)$ in the norm

$$\|u\|_{D^{1,2}(\mathbb{R}^N)} = \left(\int_{\mathbb{R}^N} |\nabla u|^2 \right)^{1/2}.$$

Let $N \geq 3$ and define the constants $\alpha_N = N(N-2)$ and $C_N = (N(N-2))^{\frac{N-2}{4}}$. Let $K \in C^2(\mathbb{R}^N)$. We consider the following problem for $\varepsilon \geq 0$:

$$(P_\varepsilon) \quad \left\{ \begin{array}{l} \text{Find } u \in D^{1,2}(\mathbb{R}^N) \text{ solving :} \\ -\Delta u = \alpha_N(1 + \varepsilon K(x))u^{\frac{N+2}{N-2}}, \\ u > 0 \end{array} \right\} \text{ in } \mathbb{R}^N.$$

We are interested in showing an exact multiplicity result for (P_ε) for all small $\varepsilon > 0$ (see Theorem 1.4 below).

The above problem is a “perturbed” version of the well-known scalar curvature problem which arises in differential geometry. More precisely, the problem is to find out if a given smooth function R on the N -dimensional unit sphere S^N is the scalar curvature function of a metric g on S^N which is conformal to the standard

---

Received by the editors April 14, 2005 and, in revised form, July 30, 2005.

2000 *Mathematics Subject Classification*. Primary 35B32, 35B33.

Key words and phrases. Yamabe problem, exact multiplicity, scalar curvature.

©2006 American Mathematical Society
 Reverts to public domain 28 years from publication

metric. This gives rise to the following problem:

$$(P_\varepsilon)_g \quad \left\{ \begin{array}{l} \text{Find } u \in C^2(S^N) \text{ solving:} \\ \frac{4(N-1)}{N-2} \Delta_g u + R u^{\frac{N+2}{N-2}} = N(N-1)u, \\ u > 0 \end{array} \right\} \text{ in } S^N.$$

The above problem has been extensively studied using the background of differential geometry; see the book of T. Aubin [2] for a survey of the available results.

We now assume that R is a perturbation of the constant, viz, $R = 1 + \varepsilon K$ for a smooth function K on S^N and $\varepsilon > 0$ small. Then, using the standard stereographic projection from S^N to $\mathbb{R}^N$, it can be checked that $(P_\varepsilon)_g$ is transformed to (P_ε) .

Existence of solutions to (P_ε) was done in [1] using variational methods and finite-dimensional reduction techniques. To describe their result, we make the following assumptions on K :

- (K1) $K \in C^2(\mathbb{R}^N)$, $\|K\|_{L^\infty(\mathbb{R}^N)} + \|\nabla K\|_{L^\infty(\mathbb{R}^N; \mathbb{R}^N)} + \|D^2 K\|_{L^\infty(\mathbb{R}^N; \mathbb{R}^N \times \mathbb{R}^N)} < \infty$.
- (K2) (a) There exists $\rho > 0$ such that $\langle \nabla K(x), x \rangle < 0 \quad \forall |x| \geq \rho$,
 (b) $\langle \nabla K(x), x \rangle \in L^1(\mathbb{R}^N)$, $\int_{\mathbb{R}^N} \langle \nabla K(x), x \rangle dx < 0$.
- (K3) The set of all critical points of K , denoted by $\text{crit}(K)$, is finite.
- (K4) $\forall \xi \in \text{crit}(K)$, there exists $\beta = \beta_\xi \in [1, N]$ and $Q_\xi : \mathbb{R}^N \rightarrow \mathbb{R}$ depending continuously on y locally near ξ such that

$$A_\xi \stackrel{\text{def}}{=} \frac{N-2}{2N} \int_{\mathbb{R}^N} Q_\xi(y) U_{1,0}^{\frac{2N}{N-2}}(y) dy \neq 0$$

and the following relations hold:

$$(K5) \quad \left\{ \begin{array}{l} Q_\xi(\lambda x) = \lambda^\beta Q_\xi(x), \quad \forall \lambda \geq 0, \\ K(x) = K(y) + Q_\xi(x-y) + o(|x-y|^\beta) \text{ as } x \rightarrow y, \end{array} \right.$$

$$\sum_{\xi \in \text{crit}(K)} \deg_{loc}(\nabla K, \xi) \neq (-1)^N.$$

We also recall the following well-known classification result for solutions of (P_0) :

Theorem 1.1. *Solutions of (P_0) form an $(N+1)$ -dimensional manifold given by*

$$\mathcal{M} = \{U_{\delta,y}(x) \stackrel{\text{def}}{=} C_N \delta^{\frac{N-2}{2}} (\delta^2 + |x-y|^2)^{\frac{2-N}{2}} : (\delta, y) \in \mathbb{R}^+ \times \mathbb{R}^N\}.$$

We can now state the following existence result.

Theorem 1.2 ([1]). *Let K satisfy the assumptions (K1)–(K5). Then there exists $\varepsilon_0 > 0$ such that the following hold:*

- (i) (P_ε) has a solution $\forall \varepsilon \in (0, \varepsilon_0)$,
- (ii) $\forall \varepsilon \in (0, \varepsilon_0)$, there exists $(\delta_\varepsilon, y_\varepsilon) \in \mathbb{R}^+ \times \mathbb{R}^N$ such that for any compact set $A \subset \mathbb{R}^+ \times \mathbb{R}^N$, we may find a constant $c(A) > 0$ such that $\|u_\varepsilon - U_{\delta_\varepsilon, y_\varepsilon}\|_{D^{1,2}(\mathbb{R}^N)} \leq c(A)\varepsilon$.

We now define what we mean by a stable zero of a vector field.

Definition 1.3. Let $\mathbf{G} : \mathbb{R}^+ \times \mathbb{R}^N \rightarrow \mathbb{R}^{N+1}$ be a C^1 vector field. We say that a point $(\delta, y) \in \mathbb{R}^+ \times \mathbb{R}^N$ is a stable zero for $\mathbf{G}$ if $\mathbf{G}(\delta, y) = \mathbf{0}$ and its derivative $D\mathbf{G}(\delta, y)$ is an invertible matrix.

We define the following functional:

$$(1.1) \quad \Gamma(\delta, y) = \frac{N-2}{2N} \int_{\mathbb{R}^N} K(\delta x + y) U_{1,0}^{\frac{2N}{N-2}}(x) dx.$$

For a set $A \subset \mathbb{R}^+ \times \mathbb{R}^N$, we let $\mathcal{M}_A = \{U_{\delta,y} : (\delta, y) \in A\}$. For a function $u \in D^{1,2}(\mathbb{R}^N)$, let $d(u, \mathcal{M}_A) = \inf_{(\delta,y) \in A} \|u - U_{\delta,y}\|_{D^{1,2}(\mathbb{R}^N)}$. We can now state the following exact multiplicity result which we will prove later in §4.

Theorem 1.4. *Let K satisfy the assumptions (K1)–(K5). We further suppose that $\nabla \Gamma$ has finitely many zeroes in $\mathbb{R}^+ \times \mathbb{R}^N$, all of which are stable. Let $A \subset \mathbb{R}^+ \times \mathbb{R}^N$ be any compact set containing the zeroes of $\nabla \Gamma$. Then there exist $\rho_0 = \rho_0(A) > 0$ and $\varepsilon_0 = \varepsilon_0(\rho) > 0$ such that for all $\varepsilon \in (0, \varepsilon_0)$, the problem (P_ε) has exactly the same number of solutions u with $d(u, \mathcal{M}_A) < \rho_0$ as the number of zeroes of $\nabla \Gamma$.*

2. PRELIMINARY RESULTS

In this section, we recall some of the well-known results concerning the problem (P_ε) and its linearized version. Given a solution u_ε of (P_ε) we consider the following linearization of (P_ε) about u_ε :

$$((LP)_\varepsilon) \quad \begin{cases} -\Delta w &= N(N+2)u_\varepsilon^{4/N-2}w \text{ in } \mathbb{R}^N, \\ w &\in D^{1,2}(\mathbb{R}^N). \end{cases}$$

Let $(LP_0)_{\delta,y}$ denote the above linearized problem when $\varepsilon = 0$ and $u_\varepsilon = U_{\delta,y}$ for some $(\delta, y) \in \mathbb{R}^+ \times \mathbb{R}^N$. Then, we have the following characterization of solutions of $(LP_0)_{\delta,y}$.

Theorem 2.1. *Every solution of $(LP_0)_{\delta,y}$ is of the form*

$$(2.1) \quad w = c_0 \frac{\delta U_{\delta,y}}{\partial \delta} + \sum_{k=1}^N c_k \frac{\partial U_{\delta,y}}{\partial y_i}$$

for some $c_i \in \mathbb{R}$, $i = 0, 1, \dots, N$.

Proof. See ([3]). □

We now recall the following natural decay estimates for solutions of (P_ε) and $(LP)_\varepsilon$.

Theorem 2.2. *Let $\{u_\varepsilon\}_{\varepsilon>0}$ be a sequence of solutions of (P_ε) with $\sup_{\varepsilon>0} \|u_\varepsilon\|_{D^{1,2}(\mathbb{R}^N)} < \infty$. Let $\{w_\varepsilon\}_{\varepsilon>0}$ be a sequence of solutions of $(LP)_\varepsilon$ with either $\sup_{\varepsilon>0} \|w_\varepsilon\|_{L^\infty(\mathbb{R}^N)} < \infty$ or $\sup_{\varepsilon>0} \|w_\varepsilon\|_{D^{1,2}(\mathbb{R}^N)} < \infty$. Then we have the following decay estimates:*

$$\begin{aligned} \sup_{\varepsilon>0} \sup_{x \in \mathbb{R}^N} \{|x|^{N-2} u_\varepsilon(x) + |x|^{N-1} |\nabla u_\varepsilon(x)|\} &< \infty, \\ \sup_{\varepsilon>0} \sup_{x \in \mathbb{R}^N} \{|x|^{N-2} |w_\varepsilon(x)| + |x|^{N-1} |\nabla w_\varepsilon(x)|\} &< \infty. \end{aligned}$$

Proof. Follows in a standard way by using the Kelvin Transform in $\mathbb{R}^N$ and elliptic regularity. □

Finally, we state the very important Pohozaev identity for (P_ε) :

Theorem 2.3. *Let $\{u_\varepsilon\}_{\varepsilon>0}$ be a sequence of solutions of (P_ε) with $\sup_{\varepsilon>0} \|u_\varepsilon\|_{D^{1,2}(\mathbb{R}^N)} < \infty$. Then the following identities hold:*

$$(2.2) \quad \int_{\mathbb{R}^N} K(x) u_\varepsilon^{\frac{N+2}{N-2}} \frac{\partial u_\varepsilon}{\partial x_i} = 0, \quad i = 1, \dots, N,$$

$$(2.3) \quad \int_{\mathbb{R}^N} K(x) u_\varepsilon^{\frac{N+2}{N-2}} [(x-y) \cdot \nabla u_\varepsilon + \left(\frac{N-2}{2}\right) u_\varepsilon] = 0.$$

Proof. See [4]. □

3. LOCAL UNIQUENESS OF SOLUTIONS

We note that if $\{u_\varepsilon\}_{\varepsilon>0}$ is a sequence of solutions of (P_ε) converging (as $\varepsilon \rightarrow 0$) to $U_{\delta,y}$ in $D^{1,2}(\mathbb{R}^N)$, then thanks to Theorems 2.2 and 2.3 we obtain that necessarily $\nabla \Gamma(\delta, y) = \mathbf{0}$. We show in this section that for all small enough $\varepsilon > 0$ there is at most one sequence of solutions $\{u_\varepsilon\}$ “bifurcating” from $U_{\delta,y}$ when (δ, y) is a stable zero of $\nabla \Gamma$.

Theorem 3.1. *Let $(\delta, y) \in \mathbb{R}^+ \times \mathbb{R}^N$ be a stable zero of $\nabla \Gamma$. Let $\{u_{1,\varepsilon}\}_{\varepsilon>0}$ and $\{u_{2,\varepsilon}\}_{\varepsilon>0}$ be two sequences of solutions to (P_ε) with $\|u_{i,\varepsilon} - U_{\delta,y}\|_{D^{1,2}(\mathbb{R}^N)} \rightarrow 0$ as $\varepsilon \rightarrow 0, i = 1, 2$. Then, there exists $\varepsilon_0 > 0$ such that $u_{1,\varepsilon} \equiv u_{2,\varepsilon} \forall \varepsilon \in (0, \varepsilon_0)$.*

Proof. We suppose that (P_ε) admits two distinct sequences of solutions $\{u_{1,\varepsilon_n}\}$ and $\{u_{2,\varepsilon_n}\}$ for some sequence $\varepsilon_n \rightarrow 0$ for $i = 1, 2$ with $\|u_{i,\varepsilon_n} - U_{\delta,y}\|_{D^{1,2}(\mathbb{R}^N)} \rightarrow 0$ as $\varepsilon_n \rightarrow 0$ and arrive at a contradiction. For notational ease, we let $u_{i,n} = u_{i,\varepsilon_n}$. Define $\tilde{w}_n = u_{1,n} - u_{2,n}$. Note that $\|\tilde{w}_n\|_{D^{1,2}(\mathbb{R}^N)} \rightarrow 0$ as $n \rightarrow \infty$. By Theorem 2.2, we note that $\{\tilde{w}_n\}$ is a bounded sequence in $L^\infty(\mathbb{R}^N)$. We let $w_n = \frac{\tilde{w}_n}{\|\tilde{w}_n\|_{L^\infty(\mathbb{R}^N)}}$. Let $x_n \in \mathbb{R}^N$ be such that $|w_n(x_n)| \geq \frac{1}{2}$. We note that w_n satisfies

$$-\Delta w_n = N(N+2)(1 + \varepsilon_n K) c_n w_n,$$

where

$$c_n(x) = \int_0^1 (t u_{1,n}(x) + (1-t) u_{2,n}(x))^{4/N-2} dt.$$

Using standard regularity theory, we get that $w_n \rightarrow w$ in $C_{loc}^2(\mathbb{R}^N)$ and that w satisfies

$$-\Delta w = N(N+2) U_{\delta,y}^{4/N-2} w \text{ in } \mathbb{R}^N.$$

Thanks to Theorem 2.1 we obtain that

$$(3.1) \quad w = c_0 \frac{\partial U_{\delta,y}}{\partial \delta} + \sum_{k=1}^N c_k \frac{\partial U_{\delta,y}}{\partial y_k}$$

for some $c_k \in \mathbb{R}, k = 0, 1, \dots, N$. We now

Claim. $c_k = 0, k = 0, 1, \dots, N$.

Proof of Claim. From the Pohozaev identity (2.2) we obtain that

$$\int_{\mathbb{R}^N} K(x) \frac{\partial}{\partial x_j} \left(u_{2,n}^{\frac{2N}{N-2}} - u_{1,n}^{\frac{2N}{N-2}} \right) = 0, j = 1, \dots, N.$$

Using integration by parts and the decay estimates in Theorem 2.2, we obtain from the above equation that

$$\int_{\mathbb{R}^N} \frac{\partial K}{\partial x_j} \left(u_{2,n}^{\frac{2N}{N-2}} - u_{1,n}^{\frac{2N}{N-2}} \right) = 0, j = 1, \dots, N.$$

Using the Taylor series, we can write the above equation as

$$\int_{\mathbb{R}^N} \frac{\partial K}{\partial x_j} \left(\int_0^1 (tu_{2,n}(x) + (1-t)u_{1,n}(x))^{\frac{N+2}{N-2}} dt \right) w_n(x) dx = 0.$$

Passing to the limit, as $n \rightarrow \infty$, in the above equation we obtain

$$\int_{\mathbb{R}^N} \frac{\partial K}{\partial x_j} U_{\delta,y}^{\frac{N+2}{N-2}} w = 0.$$

Once again, integrating by parts in the above equation, we obtain

$$\left(\frac{N+2}{N-2} \right) \int_{\mathbb{R}^N} K(x) U_{\delta,y}^{4/N-2} \frac{\partial U_{\delta,y}}{\partial x_j} w dx + \int_{\mathbb{R}^N} K(x) U_{\delta,y}^{\frac{N+2}{N-2}} \frac{\partial w}{\partial x_j} dx = 0.$$

Noting that $\frac{\partial U_{\delta,y}}{\partial x_j} = -\frac{\partial U_{\delta,y}}{\partial y_j}$, we obtain from the above equation, for $1 \leq j \leq N$,

$$(3.2) \quad \left(\frac{N+2}{N-2} \right) \int_{\mathbb{R}^N} K(x) U_{\delta,y}^{4/N-2} \frac{\partial U_{\delta,y}}{\partial y_j} w + \int_{\mathbb{R}^N} K(x) U_{\delta,y}^{\frac{N+2}{N-2}} \frac{\partial w}{\partial y_j} = 0.$$

Integrating by parts on the ball $B_R(y)$, we have for $i = 1, 2$,

$$\begin{aligned} \int_{B_R(y)} (x-y) \cdot \nabla (K(x) u_{i,n}^{\frac{2N}{N-2}}) &= NR \int_{\partial B_R(y)} K(\sigma) u_{i,n}^{\frac{2N}{N-2}}(\sigma) d\sigma \\ &\quad - N \int_{B_R(y)} K(x) u_{i,n}^{\frac{2N}{N-2}} dx. \end{aligned}$$

Letting $R \rightarrow \infty$ in the above equation and using the decay estimates in Theorem 2.2 we obtain

$$\int_{\mathbb{R}^N} (x-y) \cdot \nabla (K(x) u_{i,n}^{\frac{2N}{N-2}}(x)) dx = -N \int_{\mathbb{R}^N} K(x) u_{i,n}^{\frac{2N}{N-2}}(x) dx, \quad i = 1, 2.$$

The above equation for $i = 1, 2$ can be rewritten as

$$\begin{aligned} \int_{\mathbb{R}^N} (x-y) \cdot \nabla K(x) u_{i,n}^{\frac{2N}{N-2}}(x) dx + \left(\frac{2N}{N-2} \right) \int_{\mathbb{R}^N} K(x) u_{i,n}^{\frac{N+2}{N-2}}(x) (x-y) \cdot \nabla u_{i,n}(x) dx \\ + N \int_{\mathbb{R}^N} K(x) u_{i,n}^{\frac{2N}{N-2}}(x) dx = 0. \end{aligned}$$

Using the Pohozaev identity (2.3) in the above equation, we obtain

$$\int_{\mathbb{R}^N} (x-y) \cdot \nabla K(x) u_{i,n}^{\frac{2N}{N-2}} = 0, \quad i = 1, 2.$$

Subtracting the above identity for $u_{1,n}$ from that for $u_{2,n}$, dividing by the appropriate norm of $u_{2,n} - u_{1,n}$ and finally using Taylor's expansion, we obtain

$$\int_{\mathbb{R}^N} (x-y) \cdot \nabla K(x) \left(\int_0^1 (tu_{1,n}(x) + (1-t)u_{2,n}(x))^{\frac{N+2}{N-2}} dt \right) w_n(x) dx = 0.$$

Letting $n \rightarrow \infty$ in the above equation, we obtain

$$\int_{\mathbb{R}^N} (x-y) \cdot \nabla K(x) U_{\delta,y}^{\frac{N+2}{N-2}}(x) w(x) dx = 0.$$

We again integrate by parts in the above equation to obtain

$$(3.3) \quad \int_{\mathbb{R}^N} K(x) U_{\delta,y}^{\frac{N+2}{N-2}} [Nw + (x-y) \cdot \nabla w] + \left(\frac{N+2}{N-2} \right) \int_{\mathbb{R}^N} K(x) U_{\delta,y}^{4/N-2} (x-y) \cdot \nabla U_{\delta,y} w = 0.$$

A simple calculation (see the Appendix) using the Pohozaev identities in (2.2)–(2.3) shows that, if we let $\mathbf{c} = (c_0, \dots, c_N)^T$, then (3.2) and (3.3) imply that

$$D^2\Gamma(\delta, y)\mathbf{c} = \mathbf{0}.$$

Since $D^2\Gamma(\delta, y)$ was assumed to be an invertible matrix, this means that $\mathbf{c} = \mathbf{0}$, thereby proving the Claim.

From the above Claim, it follows that $w \equiv 0$ in $\mathbb{R}^N$. Hence, $w_n \rightarrow 0$ in $C_{loc}^2(\mathbb{R}^N)$ which implies that $|x_n| \rightarrow \infty$ as $n \rightarrow \infty$. Define the following Kelvin Transforms:

$$\left. \begin{aligned} \widehat{u}_{i,n}(x) &= |x|^{2-N} u_{i,n}\left(\frac{x}{|x|^2}\right), \quad i = 1, 2, \\ \widehat{w}_n(x) &= |x|^{2-N} w_n\left(\frac{x}{|x|^2}\right), \\ \widehat{c}_n(x) &= \int_0^1 (t\widehat{u}_{1,n}(x) + (1-t)\widehat{u}_{2,n}(x))^{\frac{4}{N-2}} dt \end{aligned} \right\} x \in \mathbb{R}^n \setminus \{0\}.$$

Then, it can be checked that $\widehat{w}_n$ satisfies the following equation:

$$-\Delta \widehat{w}_n(x) = N(N+2)(1 + \varepsilon_n K(\frac{x}{|x|^2})) \widehat{c}_n(x) \widehat{w}_n(x) \text{ in } \mathbb{R}^N \setminus \{0\}.$$

Using the decay estimates in Theorem 2.2 we obtain that

$$\sup_n \sup_{x \in B_1(0) \setminus \{0\}} |\widehat{w}_n(x)| < \infty.$$

Since $\widehat{w}_n \rightarrow 0$ in $C_{loc}^2(B_1(0) \setminus \{0\})$, by the dominated convergence theorem, we obtain that $\widehat{w}_n \rightarrow 0$ in $L^p(B_1(0)) \forall p \geq 1$. It is also easy to see that $\{\widehat{c}_n\}$ is a bounded sequence in $L^2(B_1(0))$. Since the capacity of one point set is zero, we can apply the regularity theory to $\widehat{w}_n$ (see Theorem 8.17 of Gilbarg-Trudinger, "Elliptic PDEs of Second Order") and get that

$$\|\widehat{w}_n\|_{L^\infty(B_{1/2}(0))} \leq c \|\widehat{w}_n\|_{L^p(B_1(0))} \quad \forall p > 1.$$

But this implies $\|\widehat{w}_n\|_{L^\infty(B_{1/2}(0))} \rightarrow 0$ as $n \rightarrow \infty$, contradicting the fact that $|\widehat{w}_n(\frac{x_n}{|x_n|^2})| \geq \frac{1}{2}$ for all large n . This proves the theorem. $\square$

4. THE EXACT MULTIPLICITY RESULT

We are now ready to prove the exact multiplicity result stated in Theorem 1.4.

Proof of Theorem 1.4. Let M be the number of zeroes of $\nabla\Gamma$. Appealing to Theorem 1.2 we obtain that there exists $\varepsilon_1 = \varepsilon_1(A) > 0$ such that for any $\varepsilon \in (0, \varepsilon_1)$, the problem (P_ε) has at least M solutions $\{u_\varepsilon^i\}_{i=1}^M$ and M points $\{(\delta_i, y_i)\}_{i=1}^M \subset A$ such that $u_\varepsilon^i - U_{\delta_i, y_i} \rightarrow 0$ in $D^{1,2}(\mathbb{R}^N)$, $i = 1, \dots, M$, as $\varepsilon \rightarrow 0$. For $\mu > 0$ define

$$\mathcal{S}_\mu = \{u : u \text{ solves } (P_\varepsilon) \text{ for } \varepsilon \in (0, \mu)\} \setminus \{u_\varepsilon^i\}_{0 < \varepsilon < \mu, 1 \leq i \leq M}.$$

Now define the quantity

$$\theta_\mu = \inf_{u \in \mathcal{S}_\mu} d(u, \mathcal{M}_A).$$

We now have the following claim.

Claim. $\theta_0 = \lim_{\mu \rightarrow 0} \theta_\mu > 0$.

We suppose that $\theta_0 = 0$ and derive a contradiction. We may then find sequences $\mu_n \rightarrow 0$, $\{u_n\} \subset \mathcal{S}_{\mu_n}$ and $\{(\delta_n, y_n)\} \subset A$ such that $\|u_n - U_{\delta_n, y_n}\|_{D^{1,2}(\mathbb{R}^N)} \rightarrow 0$ as $n \rightarrow \infty$. Let $(\delta_n, y_n) \rightarrow (\delta, y) \in A$. Clearly we have $\nabla\Gamma(\delta, y) = \mathbf{0}$ which means that $\{u_n\}$ is a sequence of solutions “bifurcating” from (δ, y) . But using the uniqueness result in Theorem 3.1 we obtain a contradiction since $\{u_n\} \subset \mathcal{S}_{\mu_n}$ for all n . This proves the Claim.

Therefore, we may choose $\mu_0 > 0$ small enough (but fixed) so that $\theta_\mu \geq \frac{\theta_0}{2}$ for all $0 < \mu < \mu_0$. Also from Theorem 1.2 we obtain that for some constant $c > 1$ and $\varepsilon_2 > 0$ we have $d(u_\varepsilon^i, \mathcal{M}_A) \leq c\varepsilon$, $i = 1, \dots, M$, $\varepsilon \in (0, \varepsilon_2)$. The theorem now follows by taking $\rho_0 = \frac{\theta_0}{2}$ and $\varepsilon_0 = \min\{\frac{\theta_0}{2c}, \mu_0, \varepsilon_2\}$. $\square$

5. APPENDIX

Let $(\delta_0, y_0) \in \mathbb{R}^+ \times \mathbb{R}^N$ be a stable critical point of Γ . This in particular means that (see Theorem 1.2) there exists a sequence of solutions $\{u_\varepsilon\}$ of (P_ε) converging to U_{δ_0, y_0} in $D^{1,2}(\mathbb{R}^N)$ as $\varepsilon \rightarrow 0$. As a consequence we have that U_{δ_0, y_0} satisfies the Pohozaev identities (2.2)–(2.3).

Let

$$(5.1) \quad \Lambda(\delta, y) = \int_{\mathbb{R}^N} K(\delta x + y) U_{1,0}^{\frac{2N}{N-2}} dx, \quad (\delta, y) \in \mathbb{R}^+ \times \mathbb{R}^N.$$

Note that $\Lambda = (\frac{N-2}{2N})\Gamma$. We can now state

Proposition A.1. *Let K satisfy assumptions (K1)–(K5). Let $(\delta_0, y_0) \in \mathbb{R}^+ \times \mathbb{R}^N$ be a stable critical point of Γ . Let*

$$(5.2) \quad w = c_0 \frac{\partial U_{\delta_0, y_0}}{\partial \delta} + \sum_{k=1}^N c_k \frac{\partial U_{\delta_0, y_0}}{\partial y_k}.$$

Then (3.2)–(3.3) imply that $D^2\Gamma(\delta_0, y_0) \cdot (c_0, c_1, \dots, c_N)^T = \mathbf{0}$.

Proof. We note that $U_{\delta,y}$ satisfies the relations:

$$(5.3) \quad U_{\delta,y}(x) = \delta^{\frac{2-N}{2}} U_{1,0} \left(\frac{x-y}{\delta} \right),$$

$$(5.4) \quad \nabla U_{\delta,y}(x) = \delta^{-\frac{N}{2}} \nabla U_{1,0} \left(\frac{x-y}{\delta} \right),$$

$$(5.5) \quad (x-y) \cdot \nabla U_{\delta,y} = - \left[\delta \frac{\partial U_{\delta,y}}{\partial \delta} + \left(\frac{N-2}{2} \right) U_{\delta,y} \right].$$

We further note that since (δ_0, y_0) is a critical point of Γ , we have that U_{δ_0, y_0} satisfies the Pohozaev identities in (2.2)–(2.3). We now compute $D^2\Lambda(\delta_0, y_0)$. We have,

$$(5.6) \quad \frac{\partial \Lambda}{\partial \delta}(\delta, y) = \int_{\mathbb{R}^N} x \cdot \nabla K(\delta x + y) U_{1,0}^{\frac{2N}{N-2}}(x) dx,$$

$$(5.7) \quad \frac{\partial \Lambda}{\partial y_i}(\delta, y) = \frac{1}{\delta} \int_{\mathbb{R}^N} \frac{\partial K}{\partial x_i}(\delta x + y) U_{1,0}^{\frac{2N}{N-2}}(x) dx, \quad 1 \leq i \leq N.$$

In (5.6)–(5.7) we make the change of variable $z = \delta x + y$, use (5.3), integrate by parts and again change back to the x variable to get

$$(5.8) \quad \begin{aligned} -\delta \frac{\partial \Lambda}{\partial \delta}(\delta, y) &= N \int_{\mathbb{R}^N} K(\delta x + y) U_{1,0}^{\frac{2N}{N-2}}(x) dx \\ &\quad + \left(\frac{2N}{N-2} \right) \int_{\mathbb{R}^N} K(\delta x + y) U_{1,0}^{\frac{N+2}{N-2}}(x) (x \cdot \nabla U_{1,0}(x)) dx, \end{aligned}$$

$$(5.9) \quad -\delta \frac{\partial \Lambda}{\partial y_i}(\delta, y) = \left(\frac{2N}{N-2} \right) \int_{\mathbb{R}^N} K(\delta x + y) U_{1,0}^{\frac{N+2}{N-2}}(x) \frac{\partial U_{1,0}}{\partial x_i}(x) dx, \quad 1 \leq i \leq N.$$

Differentiating (5.8) on both sides with respect to δ, y_i ($1 \leq i \leq N$), noting that $\frac{\partial \Lambda}{\partial \delta}(\delta_0, y_0) = \frac{\partial \Lambda}{\partial y_i}(\delta_0, y_0) = 0$, changing variables as $z = \delta x + y$, using (5.3)–(5.5) and integrating by parts, we get

$$(5.10) \quad \begin{aligned} -\delta_0 \frac{\partial^2 \Lambda}{\partial \delta^2}(\delta_0, y_0) &= \left(\frac{2N}{N-2} \right) \left[N \int_{\mathbb{R}^N} K(z) U_{\delta_0, y_0}^{\frac{N+2}{N-2}}(z) \frac{\partial U_{\delta_0, y_0}}{\partial \delta}(z) dz \right. \\ &\quad + \int_{\mathbb{R}^N} K(z) U_{\delta_0, y_0}^{\frac{N+2}{N-2}}(z) (z - y_0) \cdot \nabla \left(\frac{\partial U_{\delta_0, y_0}}{\partial \delta} \right) dz \\ &\quad \left. + \left(\frac{N+2}{N-2} \right) \int_{\mathbb{R}^N} K(z) U_{\delta_0, y_0}^{\frac{4}{N-2}}(z) (z - y_0) \cdot \nabla U_{\delta_0, y_0}(z) \frac{\partial U_{\delta_0, y_0}}{\partial \delta}(z) dz \right], \end{aligned}$$

$$(5.11) \quad \begin{aligned} -\frac{\partial^2 \Lambda}{\partial \delta \partial y_i}(\delta_0, y_0) &= \left(\frac{2N}{N-2} \right) \left[\int_{\mathbb{R}^N} K(z) U_{\delta_0, y_0}^{\frac{N+2}{N-2}}(z) \frac{\partial}{\partial z_i} \left(\frac{\partial U_{\delta_0, y_0}}{\partial \delta}(z) \right) dz \right. \\ &\quad \left. + \left(\frac{N+2}{N-2} \right) \int_{\mathbb{R}^N} K(z) U_{\delta_0, y_0}^{\frac{4}{N-2}}(z) \frac{\partial U_{\delta_0, y_0}}{\partial \delta}(z) \frac{\partial U_{\delta_0, y_0}}{\partial z_i} dz \right]. \end{aligned}$$

In deriving (5.11) we need to additionally use that the Pohozaev identity (2.2) holds for U_{δ_0, y_0} in the last step.

Again, for $1 \leq i, j \leq N$, differentiating (5.9) with respect to y_j , using (5.3)–(5.5) and integrating by parts, we get

$$(5.12) \quad -\frac{\partial^2 \Lambda}{\partial y_i \partial y_j}(\delta_0, y_0) = \left(\frac{2N}{N-2} \right) \left[\int_{\mathbb{R}^N} K(z) U_{\delta_0, y_0}^{\frac{N+2}{N-2}}(z) \frac{\partial}{\partial z_j} \left(\frac{\partial U_{\delta_0, y_0}}{\partial z_i}(z) \right) dz \right. \\ \left. + \left(\frac{N+2}{N-2} \right) \int_{\mathbb{R}^N} K(z) U_{\delta_0, y_0}^{\frac{4}{N-2}}(z) \frac{\partial U_{\delta_0, y_0}}{\partial z_j^0}(z) \frac{\partial U_{\delta_0, y_0}}{\partial z_i}(z) dz \right].$$

Let w be as in (5.2). Using (3.2)–(3.3) we can easily check that (5.10)–(5.12) imply that $D^2\Gamma(\delta_0, y_0) \cdot (c_0, c_1, \dots, c_N)^T = \mathbf{0}$. $\square$

ACKNOWLEDGMENT

I would like to thank Professor Massimo Grossi for helpful discussions and the anonymous referee for suggestions that improved the presentation of the paper.

REFERENCES

- [1] A. Ambrosetti, J. Garcia Azorero and I. Peral, Perturbation of $\Delta u + u^{\frac{N+2}{N-2}} = 0$, the scalar curvature problem in $\mathbb{R}^N$ and related topics, *J. Funct. Anal.* **165** (1999), 117-149. MR1696454 (2000d:35047)
- [2] T. Aubin, Some nonlinear problems in Riemannian Geometry, Springer Monographs in Mathematics, *Springer-Verlag, Berlin* (1998). MR1636569 (99i:58001)
- [3] G. Bianchi and H. Egnell, A note on Sobolev inequality, *J. Funct. Anal.* **100** (1991), 10-24. MR1124290 (92i:46033)
- [4] J. Kazdan and F. Warner, Existence and conformal deformation of metrics with prescribed Gaussian and Scalar curvature, *Ann. of Math.* **101** (1975), 317-331. MR0375153 (51:11349)

TIFR CENTRE, INDIAN INSTITUTE OF SCIENCE CAMPUS, P.B. NO. 1234, BANGALORE - 560 012, INDIA